

Massachusetts
Highly Erodible Soils
and
Potentially Highly Erodible Soils

Bristol County – Northern Part

Publication Map Symbol	SSURGO Map Symbol	<i>Map Units Highly Erodible by Water</i> (Only those units considered tillable)* Map Unit Name
HfC	245C	Hinckley sandy loam 8 to 15 percent slopes
HfD	245D	Hinckley sandy loam, 15 to 25 percent slopes
HfE	245E	Hinckley sandy loam, 25 to 35 percent slopes
MeC	254C	Merrimac fine sandy loam, 8 to 15 percent slopes
PaC	305C	Paxton fine sandy loam, 8 to 15 percent slopes
WnD	255D	Windsor loamy sand, 15 to 25 percent slopes

Publication Map Symbol	SSURGO Map Symbol	<i>Map Units <u>POTENTIALLY</u> Highly Erodible by Water</i> (Only those units considered tillable)* Map Unit Name	Critical LS Factor
AgB	275B	Agawam fine sandy loam, 3 to 8 percent slopes	.71
CoB	409B	Charlton-Paxton fine sandy loams, 3 to 8 percent slopes	.71
HfB	245B	Hinckley sandy loam, 3 to 8 percent slopes	.85
MeB	254B	Merrimac fine sandy loam, 3 to 8 percent slopes	.71
PaB	305B	Paxton fine sandy loam, 3 to 8 percent slopes	.71
RdB	70B	Ridgebury fine sandy loam, 3 to 8 percent slopes	.71
ScB	223B	Scio silt loam, 3 to 8 percent slopes	.35
StB	260B	Sudbury fine sandy loam, 3 to 8 percent slopes	.71
UnB	230B	Unadilla very fine sandy loam, 3 to 8 percent slopes	.35
WnC	255C	Windsor loamy sand, 8 to 15 percent slopes	1.4
WrB	310B	Woodbridge fine sandy loam, 3 to 8 percent slopes	.71

5/12/2006

* Stony and rocky units, urban land, “soil” – urban land complexes, pits, Udorthents, udipsamments, etc., are not considered tillable. Should you encounter an instance where such areas are converted to cropland or are farmed, contact the local soil scientist or state soils staff for assistance.