

Massachusetts  
**Highly Erodible Soils**  
**and**  
**Potentially Highly Erodible Soils**

**Hampshire County, Central Part**

<b>Publication Map Symbol</b>	<b>SSURGO Map Symbol</b>	<i>Map Units Highly Erodible by Water</i> (Only those units considered tillable)*
		<b>Map Unit Name</b>
AgC	275C	Agawam fine sandy loam, 8 to 15 percent slopes
BoC	220C	Boxford silt loam, 8 to 15 percent slopes
CkC	405C	Charlton fine sandy loam, 8 to 15 percent slopes
GfC	440C	Gloucester fine sandy loam, 8 to 15 percent slopes
HfC	251C	Haven very fine sandy loam, 8 to 20 percent slopes
HgC	253C	Hinckley loamy sand, 8 to 15 percent slopes
HgD	253D	Hinckley loamy sand, 15 to 25 percent slopes
HgE	253E	Hinckley loamy sand, 25 to 35 percent slopes
MeC	254C	Merrimac fine sandy loam, 8 to 15 percent slopes
MeD	254D	Merrimac fine sandy loam, 15 to 25 percent slopes
MoC	300C	Montauk fine sandy loam, 8 to 15 percent slopes
PaC	305C	Paxton fine sandy loam, 8 to 15 percent slopes
PaD	305D	Paxton fine sandy loam, 15 to 25 percent slopes
PuC	250C	Pollux fine sandy loam, 8 to 15 percent slopes
WeC	397C	Wethersfield fine sandy loam, 8 to 15 percent slopes
WnD	255D	Windsor loamy sand, 15 to 25 percent slopes
WtC	310C	Woodbridge fine sandy loam, 8 to 15 percent slopes

7/13/2006

\* Stony and rocky units, urban land, “soil” – urban land complexes, pits, Udorthents, udipsamments, etc., are not considered tillable. Should you encounter an instance where such areas are converted to cropland or are farmed, contact the local soil scientist or state soils staff for assistance.

Massachusetts  
**Highly Erodible Soils**  
**and**  
**Potentially Highly Erodible Soils**

**Hampshire County, Central Part**

<b>Publication Map Symbol</b>	<b>SSURGO Map Symbol</b>	<b>Map Units <u>POTENTIALLY</u> Highly Erodible by Water</b> (Only those units considered tillable)*  <b>Map Unit Name</b>	<b>Critical LS Factor</b>
AgB	275B	Agawam fine sandy loam, 3 to 8 percent slopes	.66
AmB	258B	Amostown fine sandy loam, 3 to 8 percent slopes	.57
BaB	225B	Belgrade silt loam, 3 to 8 percent slopes	.33
BoB	220B	Boxford silt loam, 3 to 8 percent slopes	.5
CkB	405B	Charlton fine sandy loam, 3 to 8 percent slopes	.66
EsB	727B	Enosburg fine sandy loam, 3 to 8 percent slopes	.66
GfB	440B	Gloucester fine sandy loam, 3 to 8 percent slopes	.66
HfB	251B	Haven very fine sandy loam, 3 to 8 percent slopes	.5
HgB	253B	Hinckley loamy sand, 3 to 8 percent slopes	.8
MeB	254B	Merrimac fine sandy loam, 3 to 8 percent slopes	.66
MoB	300B	Montauk fine sandy loam, 3 to 8 percent slopes	.5
NgB	276B	Ninigret fine sandy loam, 3 to 8 percent slopes	.57
PaB	305B	Paxton fine sandy loam, 3 to 8 percent slopes	.66
PuB	250B	Pollux fine sandy loam, 3 to 8 percent slopes	.57
RdB	70B	Ridgebury fine sandy loam, 3 to 8 percent slopes	.66
SgB	315B	Scituate fine sandy loam, 3 to 8 percent slopes	.66
SrB	260B	Sudbury fine sandy loam, 3 to 8 percent slopes	.66
WeB	397B	Wethersfield fine sandy loam, 3 to 8 percent slopes	.57
WnC	255C	Windsor loamy sand, 8 to 15 percent slopes	1.57
WoC	749C	Windsor loamy sand, silty substratum, 8 to 15 percent slopes	1.57
Wp	750C	Windsor-Scitico-Amostown complex	.76
WtB	310B	Woodbridge fine sandy loam, 3 to 8 percent slopes	.66

7/13/2006

\* Stony and rocky units, urban land, “soil” – urban land complexes, pits, Udorthents, udipsamments, etc., are not considered tillable. Should you encounter an instance where such areas are converted to cropland or are farmed, contact the local soil scientist or state soils staff for assistance.