

Massachusetts
Highly Erodible Soils
and
Potentially Highly Erodible Soils

Plymouth County

Publication Map Symbol	No SSURGO Map Symbol available	<i>Map Units Highly Erodible by Water</i> (Only those units considered tillable)*
		Map Unit Name
BbC		Bernardston silt loam, 8 to 15 percent slopes
EnC		Enfield very fine sandy loam, 8 to 15 percent slopes
EsC		Essex coarse sandy loam, 8 to 15 percent slopes
GaC		Gloucester fine sandy loam, firm substratum, 8 to 15 percent slopes
HaE		Hinckley gravelly loamy sand, 15 to 35 percent slopes
MeC		Merrimac fine sandy loam, 8 to 15 percent slopes
MfC		Merrimac sandy loam, 8 to 15 percent slopes
MfE		Merrimac sandy loam, 15 to 35 percent slopes
QuC		Quonset sandy loam, 8 to 15 percent slopes
QuE		Quonset sandy loam, 15 to 35 percent slopes
WbC		Warwick fine sandy loam, 8 to 15 percent slopes
WnE		Windsor loamy sand, 15 to 35 percent slopes

7/13/2006

* Stony and rocky units, urban land, “soil” – urban land complexes, pits, Udorthents, udipsamments, etc., are not considered tillable. Should you encounter an instance where such areas are converted to cropland or are farmed, contact the local soil scientist or state soils staff for assistance.

Massachusetts
Highly Erodible Soils
and
Potentially Highly Erodible Soils

Plymouth County

Publication Map Symbol	No SSURGO Map Symbol available	<i>Map Units <u>POTENTIALLY</u> Highly Erodible by Water</i> (Only those units considered tillable)* Map Unit Name	Critical LS Factor
AfB		Agawam fine sandy loam, 3 to 8 percent slopes	.73
AgB		Agawam fine sandy loam, silty subsoil variant, 3 to 8 percent slopes	.73
BaB		Belgrade silt loam, 3 to 8 percent slopes	.36
BbB		Bernardston silt loam, 3 to 8 percent slopes	.63
EnB		Enfield very fine sandy loam, 3 to 8 percent slopes	.36
EsB		Essex coarse sandy loam, 3 to 8 percent slopes	1.0
GaB		Gloucester fine sandy loam, firm substratum, 3 to 8 percent slopes	.73
GbB		Gloucester loamy sand, 3 to 8 percent slopes	.73
GbC		Gloucester loamy sand, 8 to 15 percent slopes	.73
HaB		Hinckley gravelly loamy sand, 3 to 8 percent slopes	1.0
HaC		Hinckley gravelly loamy sand, 8 to 15 percent slopes	1.0
HoB		Hollis-Charlton fine sandy loams, 3 to 8 percent slopes	.49
MeB		Merrimac fine sandy loam, 3 to 8 percent slopes	.73
MfB		Merrimac sandy loam, 3 to 8 percent slopes	.73
NnB		Ninigret sandy loam, silty subsoil variant, 3 to 8 percent slopes	.42
NoB		Norwell sandy loam, 3 to 8 percent slopes	.73
PtA		Pittstown silt loam, 0 to 8 percent slopes	.63
QuB		Quonset sandy loam, 3 to 8 percent slopes	.87
SeB		Scituate sandy loam, 3 to 8 percent slopes	.73
TsA		Tisbury very fine sandy loam, 0 to 8 percent slopes	.36
WbB		Warwick fine sandy loam, 3 to 8 percent slopes	.73
WnC		Windsor loamy sand, 8 to 15 percent slopes	1.72

7/13/2006

* Stony and rocky units, urban land, “soil” – urban land complexes, pits, Udorthents, udipsamments, etc., are not considered tillable. Should you encounter an instance where such areas are converted to cropland or are farmed, contact the local soil scientist or state soils staff for assistance.