

NONTECHNICAL SOIL DESCRIPTIONS

These descriptions describe soil properties or management considerations specific to a soil map unit and components of map units. These reports are generated from the National Soil Information System soil database for distribution to land users.

Cb--Coastal Beaches

Coastal Beaches component makes up 70 percent of the map unit. The assigned Kw erodibility factor is .05. This soil is poorly drained. The slowest permeability within 60 inches is rapid. Available water capacity is moderate and shrink swell potential is low. This soil is frequently flooded and is not ponded. The top of the seasonal high water table is at 36 inches. The soil has a moderately saline horizon. It is in nonirrigated land capability class 8w. This component is a hydric soil.

DoA--Downer Loamy Sand, 0 To 2 Percent Slopes

Downer component makes up 100 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .20. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 2s. This component is not a hydric soil.

DoB--Downer Loamy Sand, 2 To 5 Percent Slopes

Downer component makes up 100 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .20. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 2s. This component is not a hydric soil.

DoC--Downer Loamy Sand, 5 To 10 Percent Slopes

Downer component makes up 100 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .20. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 3e. This component is not a hydric soil.

DoC3--Downer Loamy Sand, 5 To 10 Percent Slopes, Severely Eroded

Downer component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .20. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 3e. This component is not a hydric soil.

Fa--Fallsington Loam

Fallsington component makes up 100 percent of the map unit. Prime farmland if drained. The assigned Kw erodibility factor is .32. This soil is poorly drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 6 inches. There are no saline horizons. It is in nonirrigated land capability class 4w. This component is a hydric soil.

Fb--Fallsington Sandy Loam

Fallsington component makes up 100 percent of the map unit. Prime farmland if drained. The assigned Kw erodibility factor is .24. This soil is poorly drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 6 inches. There are no saline horizons. It is in nonirrigated land capability class 4w. This component is a hydric soil.

FdA--Fallsington And Dragston Fine Sandy Loams, 0 To 2 Percent Slopes

Fallsington component makes up 50 percent of the map unit. Prime farmland if drained. The assigned Kw erodibility factor is .24. This soil is poorly drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 6 inches. There are no saline horizons. It is in nonirrigated land capability class 4w. This component is a hydric soil.

Dragston component makes up 50 percent of the map unit. Prime farmland if drained. The assigned Kw erodibility factor is .20. This soil is somewhat poorly drained. The slowest permeability within 60 inches is moderately rapid. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 21 inches. There are no saline horizons. It is in nonirrigated land capability class 3w. This component is not a hydric soil.

NONTECHNICAL SOIL DESCRIPTIONS--Continued

FdB--Fallsington And Dragston Fine Sandy Loams, 2 To 5 Percent Slopes

Fallsington component makes up 50 percent of the map unit. Prime farmland if drained. The assigned Kw erodibility factor is .24. This soil is poorly drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 6 inches. There are no saline horizons. It is in nonirrigated land capability class 4w. This component is a hydric soil.

Dragston component makes up 50 percent of the map unit. Prime farmland if drained. The assigned Kw erodibility factor is .20. This soil is somewhat poorly drained. The slowest permeability within 60 inches is moderately rapid. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 21 inches. There are no saline horizons. It is in nonirrigated land capability class 3w. This component is not a hydric soil.

FgA--Fallsington And Dragston Loams, 0 To 2 Percent Slopes

Fallsington component makes up 50 percent of the map unit. Prime farmland if drained. The assigned Kw erodibility factor is .32. This soil is poorly drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 6 inches. There are no saline horizons. It is in nonirrigated land capability class 4w. This component is a hydric soil.

Dragston component makes up 50 percent of the map unit. Prime farmland if drained. The assigned Kw erodibility factor is .20. This soil is somewhat poorly drained. The slowest permeability within 60 inches is moderately rapid. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 21 inches. There are no saline horizons. It is in nonirrigated land capability class 3w. This component is not a hydric soil.

FgB--Fallsington And Dragston Loams, 2 To 5 Percent Slopes

Fallsington component makes up 50 percent of the map unit. Prime farmland if drained. The assigned Kw erodibility factor is .32. This soil is poorly drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 6 inches. There are no saline horizons. It is in nonirrigated land capability class 4w. This component is a hydric soil.

Dragston component makes up 50 percent of the map unit. Prime farmland if drained. The assigned Kw erodibility factor is .20. This soil is somewhat poorly drained. The slowest permeability within 60 inches is moderately rapid. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 21 inches. There are no saline horizons. It is in nonirrigated land capability class 3w. This component is not a hydric soil.

GcB--Galestown Loamy Sand, Clayey Substratum, 0 To 5 Percent Slopes

Galestown component makes up 100 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .17. This soil is somewhat excessively drained. The slowest permeability within 60 inches is rapid. Available water capacity is high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 3s. This component is not a hydric soil.

GlB--Galestown-Lakeland Sands, 0 To 5 Percent Slopes

Galestown component makes up 50 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .17. This soil is somewhat excessively drained. The slowest permeability within 60 inches is rapid. Available water capacity is high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 7s. This component is not a hydric soil.

Lakeland component makes up 30 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .10. This soil is excessively drained. The slowest permeability within 60 inches is rapid. Available water capacity is high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 4s. This component is not a hydric soil.

GlC--Galestown-Lakeland Sands, 5 To 10 Percent Slopes

Galestown component makes up 50 percent of the map unit. The assigned Kw erodibility factor is .17. This soil is somewhat excessively drained. The slowest permeability within 60 inches is rapid. Available water capacity is high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 7s. This component is not a hydric soil.

NONTECHNICAL SOIL DESCRIPTIONS--Continued

Lakeland component makes up 30 percent of the map unit. The assigned Kw erodibility factor is .10. This soil is excessively drained. The slowest permeability within 60 inches is rapid. Available water capacity is high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 6s. This component is not a hydric soil.

Gp--Gravel And Borrow Pits

Gravel And Borrow Pi component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .02. The slowest permeability within 60 inches is rapid. Available water capacity is low and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 8s. This component is not a hydric soil.

Jo--Johnston Loam

Johnston Loam component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .20. This soil is very poorly drained. The slowest permeability within 60 inches is moderately rapid. Available water capacity is very high and shrink swell potential is low. This soil is frequently flooded and is not ponded. The top of the seasonal high water table is at 0 inches. There are no saline horizons. It is in nonirrigated land capability class 7w. This component is a hydric soil.

KfA--Keyport Fine Sandy Loam, 0 To 2 Percent Slopes

Keyport component makes up 95 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .37. This soil is moderately well drained. The slowest permeability within 60 inches is slow. Available water capacity is very high and shrink swell potential is moderate. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 33 inches. There are no saline horizons. It is in nonirrigated land capability class 2w. This component is not a hydric soil.

KmA--Keyport Silt Loam, 0 To 2 Percent Slopes

Keyport component makes up 95 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .43. This soil is moderately well drained. The slowest permeability within 60 inches is slow. Available water capacity is very high and shrink swell potential is moderate. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 33 inches. There are no saline horizons. It is in nonirrigated land capability class 2w. This component is not a hydric soil.

KnA--Klej Loamy Sand, 0 To 2 Percent Slopes

Klej component makes up 95 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .17. This soil is somewhat poorly drained. The slowest permeability within 60 inches is very slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 18 inches. There are no saline horizons. It is in nonirrigated land capability class 3w. This component is not a hydric soil.

KnB--Klej Loamy Sand, 2 To 5 Percent Slopes

Klej component makes up 95 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .17. This soil is somewhat poorly drained. The slowest permeability within 60 inches is very slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 18 inches. There are no saline horizons. It is in nonirrigated land capability class 3w. This component is not a hydric soil.

LaB--Lakeland Loamy Sand, Clayey Substratum 0 To 5 Percent Slopes

Lakeland component makes up 100 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .10. This soil is excessively drained. The slowest permeability within 60 inches is rapid. Available water capacity is high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 4s. This component is not a hydric soil.

LgB--Lakeland-Galestown Loamy Sands, Clayey Substratum, 2 To 5 Percent Slopes

Lakeland component makes up 50 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .10. This soil is excessively drained. The slowest permeability within 60 inches is rapid. Available water capacity is high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 4s. This component is not a hydric soil.

Galestown component makes up 30 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .17. This soil is somewhat excessively drained. The slowest permeability within 60 inches is rapid. Available water capacity is high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 3s. This component is not a hydric soil.

NONTECHNICAL SOIL DESCRIPTIONS--Continued

LmC--Lakeland-Galestown Loamy Sands, 5 To 10 Percent Slopes

Lakeland component makes up 50 percent of the map unit. The assigned Kw erodibility factor is .10. This soil is excessively drained. The slowest permeability within 60 inches is rapid. Available water capacity is high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 6s. This component is not a hydric soil.

Galestown component makes up 30 percent of the map unit. The assigned Kw erodibility factor is .17. This soil is somewhat excessively drained. The slowest permeability within 60 inches is rapid. Available water capacity is high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 7s. This component is not a hydric soil.

Lo--Leon Loamy Sand

Leon component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .10. This soil is poorly drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 12 inches. There are no saline horizons. It is in nonirrigated land capability class 4w. This component is a hydric soil.

Ma--Made Land

Made Land component makes up 100 percent of the map unit. The assigned Kw erodibility factor is . Available water capacity is very low and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 8s. This component is not a hydric soil.

MfA--Matapeake Fine Sandy Loam, 0 To 2 Percent Slopes

Matapeake component makes up 100 percent of the map unit. All areas are prime farmland. The assigned Kw erodibility factor is .37. This soil is well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 1. This component is not a hydric soil.

MfB2--Matapeake Fine Sandy Loam, 2 To 5 Percent Slopes, Moderately Eroded

Matapeake component makes up 100 percent of the map unit. All areas are prime farmland. The assigned Kw erodibility factor is .37. This soil is well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 2e. This component is not a hydric soil.

MfC--Matapeake Fine Sandy Loam, 5 To 10 Percent Slopes

Matapeake component makes up 100 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .37. This soil is well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 3e. This component is not a hydric soil.

MkA--Matapeake Silt Loam, 0 To 2 Percent Slopes

Matapeake component makes up 100 percent of the map unit. All areas are prime farmland. The assigned Kw erodibility factor is .49. This soil is well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 1. This component is not a hydric soil.

MkB2--Matapeake Silt Loam, 2 To 5 Percent Slopes, Moderately Eroded

Matapeake component makes up 100 percent of the map unit. All areas are prime farmland. The assigned Kw erodibility factor is .49. This soil is well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 2e. This component is not a hydric soil.

MkC2--Matapeake Silt Loam, 5 To 10 Percent Slopes, Moderately Eroded

Matapeake component makes up 100 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .49. This soil is well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 3e. This component is not a hydric soil.

NONTECHNICAL SOIL DESCRIPTIONS--Continued

MkC3--Matapeake Silt Loam, 5 To 10 Percent Slopes, Severely Eroded

Matapeake component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .49. This soil is well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 4e. This component is not a hydric soil.

MkD--Matapeake Silt Loam, 10 To 15 Percent Slopes

Matapeake component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .49. This soil is well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 4e. This component is not a hydric soil.

MpA--Mattapex Fine Sandy Loam, 0 To 2 Percent Slopes

Mattapex component makes up 95 percent of the map unit. All areas are prime farmland. The assigned Kw erodibility factor is .37. This soil is moderately well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 27 inches. There are no saline horizons. It is in nonirrigated land capability class 2w. This component is not a hydric soil.

MpB2--Mattapex Fine Sandy Loam, 2 To 5 Percent Slopes, Moderately Eroded

Mattapex component makes up 95 percent of the map unit. All areas are prime farmland. The assigned Kw erodibility factor is .37. This soil is moderately well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 27 inches. There are no saline horizons. It is in nonirrigated land capability class 2e. This component is not a hydric soil.

MsA--Mattapex Silt Loam, 0 To 2 Percent Slopes

Mattapex component makes up 95 percent of the map unit. All areas are prime farmland. The assigned Kw erodibility factor is .43. This soil is moderately well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 27 inches. There are no saline horizons. It is in nonirrigated land capability class 2w. This component is not a hydric soil.

MsB2--Mattapex Silt Loam, 2 To 5 Percent Slopes, Moderately Eroded

Mattapex component makes up 95 percent of the map unit. All areas are prime farmland. The assigned Kw erodibility factor is .43. This soil is moderately well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 27 inches. There are no saline horizons. It is in nonirrigated land capability class 2e. This component is not a hydric soil.

Mx--Mixed Alluvial Land

Mixed Alluvial Land component makes up 70 percent of the map unit. The assigned Kw erodibility factor is .37. This soil is poorly drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is frequently flooded and is not ponded. The top of the seasonal high water table is at 6 inches. There are no saline horizons. It is in nonirrigated land capability class 7s. This component is a hydric soil.

My--Muck And Peat

Muck And Peat component makes up 100 percent of the map unit. The assigned Kw erodibility factor is . This soil is very poorly drained. The slowest permeability within 60 inches is very rapid. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 6 inches. There are no saline horizons. It is in nonirrigated land capability class 5w. This component is a hydric soil.

OhA--Othello Silt Loam, 0 To 2 Percent Slopes

Othello component makes up 100 percent of the map unit. Prime farmland if drained. The assigned Kw erodibility factor is .37. This soil is poorly drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 6 inches. There are no saline horizons. It is in nonirrigated land capability class 3w. This component is a hydric soil.

NONTECHNICAL SOIL DESCRIPTIONS--Continued

OhB2--Othello Silt Loam, 2 To 5 Percent Slopes, Moderately Eroded

Othello component makes up 95 percent of the map unit. Prime farmland if drained. The assigned Kw erodibility factor is .37. This soil is poorly drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 6 inches. There are no saline horizons. It is in nonirrigated land capability class 3w. This component is a hydric soil.

Om--Othello Silt Loam, Low

Othello component makes up 100 percent of the map unit. Prime farmland if drained. The assigned Kw erodibility factor is .37. This soil is poorly drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 6 inches. There are no saline horizons. It is in nonirrigated land capability class 4w. This component is a hydric soil.

Oo--Othello Silt Loam, Silty Substratum

Othello component makes up 100 percent of the map unit. Prime farmland if drained. The assigned Kw erodibility factor is .37. This soil is poorly drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 6 inches. There are no saline horizons. It is in nonirrigated land capability class 3w. This component is a hydric soil.

Os--Othello Silty Clay Loam

Othello component makes up 100 percent of the map unit. Prime farmland if drained. The assigned Kw erodibility factor is .37. This soil is poorly drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 6 inches. There are no saline horizons. It is in nonirrigated land capability class 3w. This component is a hydric soil.

Ot--Othello Silty Clay Loam, Silty Substratum

Othello component makes up 100 percent of the map unit. Prime farmland if drained. The assigned Kw erodibility factor is .37. This soil is poorly drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 6 inches. There are no saline horizons. It is in nonirrigated land capability class 3w. This component is a hydric soil.

Pd--Plummer Loamy Sand

Plummer component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .10. This soil is poorly drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 6 inches. There are no saline horizons. It is in nonirrigated land capability class 4w. This component is a hydric soil.

Pk--Pocomoke Loam

Pocomoke component makes up 100 percent of the map unit. Prime farmland if drained. The assigned Kw erodibility factor is .20. This soil is very poorly drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 0 inches. There are no saline horizons. It is in nonirrigated land capability class 4w. This component is a hydric soil.

Pm--Pocomoke Sandy Loam

Pocomoke component makes up 100 percent of the map unit. Prime farmland if drained. The assigned Kw erodibility factor is .20. This soil is very poorly drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 0 inches. There are no saline horizons. It is in nonirrigated land capability class 4w. This component is a hydric soil.

Po--Portsmouth Loam

Portsmouth component makes up 100 percent of the map unit. Prime farmland if drained. The assigned Kw erodibility factor is .24. This soil is very poorly drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 6 inches. There are no saline horizons. It is in nonirrigated land capability class 6w. This component is a hydric soil.

NONTECHNICAL SOIL DESCRIPTIONS--Continued

Pr--Portsmouth Silt Loam

Portsmouth component makes up 100 percent of the map unit. Prime farmland if drained. The assigned Kw erodibility factor is .24. This soil is very poorly drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 6 inches. There are no saline horizons. It is in nonirrigated land capability class 6w. This component is a hydric soil.

Sa--St Johns Loamy Sand

St Johns component makes up 100 percent of the map unit. Prime farmland if drained. The assigned Kw erodibility factor is .10. This soil is poorly drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 3 inches. There are no saline horizons. It is in nonirrigated land capability class 3w. This component is a hydric soil.

SfA--Sassafras Sandy Loam, 0 To 2 Percent Slopes

Sassafras component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .28. This soil is well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 1. This component is not a hydric soil.

SfB2--Sassafras Sandy Loam, 2 To 5 Percent Slopes, Moderately Eroded

Sassafras component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .28. This soil is well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 2e. This component is not a hydric soil.

SfC2--Sassafras Sandy Loam, 5 To 10 Percent Slopes, Moderately Eroded

Sassafras component makes up 100 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .28. This soil is well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 3e. This component is not a hydric soil.

SfC3--Sassafras Sandy Loam, 5 To 10 Percent Slopes, Severely Eroded

Sassafras component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .28. This soil is well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 4e. This component is not a hydric soil.

SfD--Sassafras Sandy Loam, 10 To 15 Percent Slopes

Sassafras component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .28. This soil is well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 4e. This component is not a hydric soil.

St--Steep Sandy Land

Steep Sandy Land component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .28. This soil is well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 7e. This component is not a hydric soil.

Sw--Swamp

Swamp component makes up 100 percent of the map unit. The assigned Kw erodibility factor is . This soil is very poorly drained. The slowest permeability within 60 inches is very rapid. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 6 inches. There are no saline horizons. It is in nonirrigated land capability class 5w. This component is a hydric soil.

Tm--Tidal Marsh

Tidal Marsh component makes up 100 percent of the map unit. The assigned Kw erodibility factor is . This soil is poorly drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is frequently flooded and is not ponded. The top of the seasonal high water table is at 0 inches. The soil has a moderately saline horizon. It is in nonirrigated land capability class 8w. This component is a hydric soil.

NONTECHNICAL SOIL DESCRIPTIONS--Continued

WdA--Woodstown Loam, 0 To 2 Percent Slopes

Woodstown component makes up 95 percent of the map unit. All areas are prime farmland. The assigned Kw erodibility factor is .32. This soil is moderately well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 30 inches. There are no saline horizons. It is in nonirrigated land capability class 2w. This component is not a hydric soil.

WdB2--Woodstown Loam, 2 To 5 Percent Slopes, Moderately Eroded

Woodstown component makes up 95 percent of the map unit. All areas are prime farmland. The assigned Kw erodibility factor is .32. This soil is moderately well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 30 inches. There are no saline horizons. It is in nonirrigated land capability class 2w. This component is not a hydric soil.

WoA--Woodstown Sandy Loam, 0 To 2 Percent Slopes

Woodstown component makes up 95 percent of the map unit. All areas are prime farmland. The assigned Kw erodibility factor is .24. This soil is moderately well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 30 inches. There are no saline horizons. It is in nonirrigated land capability class 2w. This component is not a hydric soil.

WoB2--Woodstown Sandy Loam, 2 To 5 Percent Slopes, Moderately Eroded

Woodstown component makes up 95 percent of the map unit. All areas are prime farmland. The assigned Kw erodibility factor is .24. This soil is moderately well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 30 inches. There are no saline horizons. It is in nonirrigated land capability class 2w. This component is not a hydric soil.

