

## NONTECHNICAL SOIL DESCRIPTIONS

These descriptions describe soil properties or management considerations specific to a soil map unit and components of map units. These reports are generated from the National Soil Information System soil database for distribution to land users.

---

**AdA--Aldino Silt Loam, 0 To 3 Percent Slopes**

Aldino component makes up 95 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .43. The depth to a restrictive feature is 40 inches to bedrock (paralithic). This soil is moderately well drained. The slowest permeability within 60 inches is slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 24 inches. There are no saline horizons. It is in nonirrigated land capability class 2w. This component is not a hydric soil.

**AdB--Aldino Silt Loam, 3 To 8 Percent Slopes**

Aldino component makes up 95 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .43. The depth to a restrictive feature is 40 inches to bedrock (paralithic). This soil is moderately well drained. The slowest permeability within 60 inches is slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 24 inches. There are no saline horizons. It is in nonirrigated land capability class 2e. This component is not a hydric soil.

**AdC--Aldino Silt Loam, 8 To 15 Percent Slopes**

Aldino component makes up 95 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .43. The depth to a restrictive feature is 40 inches to bedrock (paralithic). This soil is moderately well drained. The slowest permeability within 60 inches is slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 24 inches. There are no saline horizons. It is in nonirrigated land capability class 3e. This component is not a hydric soil.

**AsB--Aldino Very Stony Silt Loam, 0 To 8 Percent Slopes**

Aldino component makes up 95 percent of the map unit. The assigned Kw erodibility factor is .37. The depth to a restrictive feature is 40 inches to bedrock (paralithic). This soil is moderately well drained. The slowest permeability within 60 inches is slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 24 inches. There are no saline horizons. It is in nonirrigated land capability class 6s. This component is not a hydric soil.

**Av--Alluvial Land**

Alluvial Land component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .43. This soil is poorly drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 6 inches. There are no saline horizons. It is in nonirrigated land capability class 7s. This component is a hydric soil.

**BaA--Baile Silt Loam, 0 To 3 Percent Slopes**

Baile component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .43. This soil is poorly drained. The slowest permeability within 60 inches is slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 3 inches. There are no saline horizons. It is in nonirrigated land capability class 5w. This component is a hydric soil.

**BaB--Baile Silt Loam, 3 To 8 Percent Slopes**

Baile component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .43. This soil is poorly drained. The slowest permeability within 60 inches is slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 3 inches. There are no saline horizons. It is in nonirrigated land capability class 6w. This component is a hydric soil.

**BeA--Beltsville Silt Loam, 0 To 2 Percent Slopes**

Beltsville component makes up 95 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .43. This soil is moderately well drained. The slowest permeability within 60 inches is slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 24 inches. There are no saline horizons. It is in nonirrigated land capability class 2w. This component is not a hydric soil.

## NONTECHNICAL SOIL DESCRIPTIONS--Continued

---

**BeB--Beltsville Silt Loam, 2 To 5 Percent Slopes**

Beltsville component makes up 95 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .43. This soil is moderately well drained. The slowest permeability within 60 inches is slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 24 inches. There are no saline horizons. It is in nonirrigated land capability class 2e. This component is not a hydric soil.

**BeC--Beltsville Silt Loam, 5 To 10 Percent Slopes**

Beltsville component makes up 95 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .43. This soil is moderately well drained. The slowest permeability within 60 inches is slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 24 inches. There are no saline horizons. It is in nonirrigated land capability class 3e. This component is not a hydric soil.

**BrC2--Brandywine Gravelly Loam, 8 To 15 Percent Slopes, Moderately Eroded**

Brandywine component makes up 100 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .20. This soil is somewhat excessively drained. The slowest permeability within 60 inches is moderately rapid. Available water capacity is high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 3e. This component is not a hydric soil.

**BrD3--Brandywine Gravelly Loam, 15 To 25 Percent Slopes, Severely Eroded**

Brandywine component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .20. This soil is somewhat excessively drained. The slowest permeability within 60 inches is moderately rapid. Available water capacity is high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 6e. This component is not a hydric soil.

**BrE3--Brandywine Gravelly Loam, 25 To 45 Percent Slopes, Severely Eroded**

Brandywine component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .20. This soil is somewhat excessively drained. The slowest permeability within 60 inches is moderately rapid. Available water capacity is high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 7e. This component is not a hydric soil.

**CcA--Chester Silt Loam, 0 To 3 Percent Slopes**

Chester component makes up 100 percent of the map unit. All areas are prime farmland. The assigned Kw erodibility factor is .32. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 1. This component is not a hydric soil.

**CcB2--Chester Silt Loam, 3 To 8 Percent Slopes Moderately Eroded**

Chester component makes up 100 percent of the map unit. All areas are prime farmland. The assigned Kw erodibility factor is .32. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 2e. This component is not a hydric soil.

**CcC2--Chester Silt Loam, 8 To 15 Percent Slopes Moderately Eroded**

Chester component makes up 100 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .32. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 3e. This component is not a hydric soil.

**CgB2--Chester Gravelly Silt Loam, 3 To 8 Percent Slopes Moderately Eroded**

Chester component makes up 100 percent of the map unit. All areas are prime farmland. The assigned Kw erodibility factor is .28. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 2e. This component is not a hydric soil.

**CgC2--Chester Gravelly Silt Loam, 8 To 15 Percent Slopes Moderately Eroded**

Chester component makes up 100 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .28. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 3e. This component is not a hydric soil.

## NONTECHNICAL SOIL DESCRIPTIONS--Continued

---

CgD2--Chester Gravelly Silt Loam, 15 To 25 Percent Slopes, Moderately Eroded  
Chester component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .28. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 4e. This component is not a hydric soil.

ChB2--Chillum Silt Loam, 2 To 5 Percent Slopes, Moderately Eroded  
Chillum component makes up 100 percent of the map unit. All areas are prime farmland. The assigned Kw erodibility factor is .43. This soil is well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 2e. This component is not a hydric soil.

CkC2--Chillum-Neshaminy Silt Loams, 5 To 10 Percent Slopes, Moderately Eroded  
Chillum component makes up 50 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .43. This soil is well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 3e. This component is not a hydric soil.

Neshaminy component makes up 30 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .32. The depth to a restrictive feature is greater than 60 inches to bedrock. This soil is well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 2e. This component is not a hydric soil.

CrE--Chrome Channery Silty Clay Loam, 15 To 45 Percent Slopes  
Chrome component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .24. The depth to a restrictive feature is 20 to 40 inches to bedrock (lithic). This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is high and shrink swell potential is moderate. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 6e. This component is not a hydric soil.

Cu--Codorus Silt Loam  
Codorus component makes up 95 percent of the map unit. Prime farmland if protected from flooding or not frequently flooded during the growing season. The assigned Kw erodibility factor is .49. This soil is moderately well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is occasionally flooded and is not ponded. The top of the seasonal high water table is at 18 inches. There are no saline horizons. It is in nonirrigated land capability class 2w. This component is not a hydric soil.

Cv--Comus Silt Loam  
Comus component makes up 100 percent of the map unit. Prime farmland if protected from flooding or not frequently flooded during the growing season. The assigned Kw erodibility factor is .43. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is occasionally flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 2w. This component is not a hydric soil.

Cx--Cut And Fill Land  
Cut And Fill Land component makes up 100 percent of the map unit. The assigned Kw erodibility factor is . Available water capacity is very low and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 8s. This component is not a hydric soil.

DcA--Delanco Silt Loam, 0 To 3 Percent Slopes  
Delanco component makes up 95 percent of the map unit. All areas are prime farmland. The assigned Kw erodibility factor is .37. This soil is moderately well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is moderate. This soil is rarely flooded and is not ponded. The top of the seasonal high water table is at 21 inches. There are no saline horizons. It is in nonirrigated land capability class 2w. This component is not a hydric soil.

## NONTECHNICAL SOIL DESCRIPTIONS--Continued

---

**DcB--Delanco Silt Loam, 3 To 8 Percent Slopes**

Delanco component makes up 95 percent of the map unit. All areas are prime farmland. The assigned Kw erodibility factor is .37. This soil is moderately well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is moderate. This soil is rarely flooded and is not ponded. The top of the seasonal high water table is at 21 inches. There are no saline horizons. It is in nonirrigated land capability class 2e. This component is not a hydric soil.

**EhB2--Elioak Silt Loam, 3 To 8 Percent Slopes, Moderately Eroded**

Elioak component makes up 100 percent of the map unit. All areas are prime farmland. The assigned Kw erodibility factor is .32. This soil is well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 2e. This component is not a hydric soil.

**EhC2--Elioak Silt Loam, 8 To 15 Percent Slopes Moderately Eroded**

Elioak component makes up 100 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .32. This soil is well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 3e. This component is not a hydric soil.

**En--Elkton Silt Loam**

Elkton component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .43. This soil is poorly drained. The slowest permeability within 60 inches is very slow. Available water capacity is very high and shrink swell potential is moderate. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 6 inches. There are no saline horizons. It is in nonirrigated land capability class 3w. This component is a hydric soil.

**EsA--Elsinboro Loam, 0 To 2 Percent Slopes**

Elsinboro component makes up 100 percent of the map unit. All areas are prime farmland. The assigned Kw erodibility factor is .37. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 60 inches. There are no saline horizons. It is in nonirrigated land capability class 1. This component is not a hydric soil.

**EsB2--Elsinboro Loam, 2 To 5 Percent Slopes Moderately Eroded**

Elsinboro component makes up 100 percent of the map unit. All areas are prime farmland. The assigned Kw erodibility factor is .37. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 60 inches. There are no saline horizons. It is in nonirrigated land capability class 2e. This component is not a hydric soil.

**EsC2--Elsinboro Loam, 5 To 10 Percent Slopes, Moderately Eroded**

Elsinboro component makes up 100 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .37. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 60 inches. There are no saline horizons. It is in nonirrigated land capability class 2e. This component is not a hydric soil.

**EvC--Evesboro Loamy Sand, 5 To 15 Percent Slopes**

Evesboro component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .17. This soil is excessively drained. The slowest permeability within 60 inches is moderately rapid. Available water capacity is high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 7s. This component is not a hydric soil.

**Fs--Fallsington Loam**

Fallsington component makes up 100 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .32. This soil is poorly drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 6 inches. There are no saline horizons. It is in nonirrigated land capability class 4w. This component is a hydric soil.

## NONTECHNICAL SOIL DESCRIPTIONS--Continued

---

**GcB2--Glenelg Loam, 3 To 8 Percent Slopes, Moderately Eroded**

Glenelg component makes up 100 percent of the map unit. All areas are prime farmland. The assigned Kw erodibility factor is .32. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 2e. This component is not a hydric soil.

**GcC2--Glenelg Loam, 8 To 15 Percent Slopes Moderately Eroded**

Glenelg component makes up 100 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .32. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 3e. This component is not a hydric soil.

**GcC3--Glenelg Loam, 8 To 15 Percent Slopes, Severely Eroded**

Glenelg component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .32. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 4e. This component is not a hydric soil.

**GcD2--Glenelg Loam, 15 To 25 Percent Slopes, Moderately Eroded**

Glenelg component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .32. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 4e. This component is not a hydric soil.

**GcD3--Glenelg Loam, 15 To 25 Percent Slopes, Severely Eroded**

Glenelg component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .32. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 6e. This component is not a hydric soil.

**GgB2--Glenelg Gravelly Loam, 3 To 8 Percent Slopes, Moderately Eroded**

Glenelg component makes up 100 percent of the map unit. All areas are prime farmland. The assigned Kw erodibility factor is .32. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 2e. This component is not a hydric soil.

**GgC2--Glenelg Gravelly Loam, 8 To 15 Percent Slopes, Moderately Eroded**

Glenelg component makes up 100 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .32. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 3e. This component is not a hydric soil.

**GgC3--Glenelg Gravelly Loam, 8 To 15 Percent Slopes Severely Eroded**

Glenelg component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .32. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 4e. This component is not a hydric soil.

**GgD2--Glenelg Gravelly Loam, 15 To 25 Percent Slopes Moderately Eroded**

Glenelg component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .32. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 4e. This component is not a hydric soil.

**GgD3--Glenelg Gravelly Loam, 15 To 25 Percent Slopes, Severely Eroded**

Glenelg component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .32. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 1. This component is not a hydric soil.

## NONTECHNICAL SOIL DESCRIPTIONS--Continued

---

**GnA--Glenville Silt Loam, 0 To 3 Percent Slopes**

Glenville component makes up 95 percent of the map unit. All areas are prime farmland. The assigned Kw erodibility factor is .32. This soil is moderately well drained. The slowest permeability within 60 inches is slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 21 inches. There are no saline horizons. It is in nonirrigated land capability class 2e. This component is not a hydric soil.

**GnB--Glenville Silt Loam, 3 To 8 Percent Slopes**

Glenville component makes up 95 percent of the map unit. All areas are prime farmland. The assigned Kw erodibility factor is .32. This soil is moderately well drained. The slowest permeability within 60 inches is slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 21 inches. There are no saline horizons. It is in nonirrigated land capability class 2w. This component is not a hydric soil.

**Hb--Hatboro Silt Loam**

Hatboro component makes up 100 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .49. This soil is poorly drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is frequently flooded and is not ponded. The top of the seasonal high water table is at 3 inches. There are no saline horizons. It is in nonirrigated land capability class 3w. This component is a hydric soil.

**JpB--Joppa Gravelly Sandy Loam, 2 To 5 Percent Slopes**

Joppa component makes up 100 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .28. This soil is well drained. The slowest permeability within 60 inches is moderately rapid. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 2s. This component is not a hydric soil.

**JpC--Joppa Gravelly Sandy Loam, 5 To 10 Percent Slopes**

Joppa component makes up 100 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .28. This soil is well drained. The slowest permeability within 60 inches is moderately rapid. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 3e. This component is not a hydric soil.

**KeB--Kelly Silt Loam, 3 To 8 Percent Slopes**

Kelly component makes up 95 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .37. The depth to a restrictive feature is 40 to 60 inches to bedrock (lithic). This soil is moderately well drained. The slowest permeability within 60 inches is slow. Available water capacity is very high and shrink swell potential is moderate. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 24 inches. There are no saline horizons. It is in nonirrigated land capability class 4w. This component is not a hydric soil.

**KeC2--Kelly Silt Loam, 8 To 15 Percent Slopes, Moderately Eroded**

Kelly component makes up 95 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .43. The depth to a restrictive feature is 42 to 60 inches to bedrock (lithic). This soil is moderately well drained. The slowest permeability within 60 inches is slow. Available water capacity is very high and shrink swell potential is moderate. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 21 inches. There are no saline horizons. It is in nonirrigated land capability class 3e. This component is not a hydric soil.

**KfD--Kelly Very Stony Silt Loam, 3 To 25 Percent Slopes**

Kelly component makes up 95 percent of the map unit. The assigned Kw erodibility factor is .24. The depth to a restrictive feature is 40 to 60 inches to bedrock (lithic). This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 6s. This component is not a hydric soil.

**KpA--Keyport Silt Loam, 0 To 2 Percent Slopes**

Keyport component makes up 95 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .43. This soil is moderately well drained. The slowest permeability within 60 inches is slow. Available water capacity is very high and shrink swell potential is moderate. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 33 inches. There are no saline horizons. It is in nonirrigated land capability class 2w. This component is not a hydric soil.

## NONTECHNICAL SOIL DESCRIPTIONS--Continued

---

**KpB--Keyport Silt Loam, 2 To 5 Percent Slopes**

Keyport component makes up 95 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .43. This soil is moderately well drained. The slowest permeability within 60 inches is slow. Available water capacity is very high and shrink swell potential is moderate. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 33 inches. There are no saline horizons. It is in nonirrigated land capability class 2e. This component is not a hydric soil.

**KrA--Kinkora Silt Loam, 0 To 3 Percent Slopes**

Kinkora component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .43. This soil is poorly drained. The slowest permeability within 60 inches is slow. Available water capacity is very high and shrink swell potential is moderate. This soil is rarely flooded and is not ponded. The top of the seasonal high water table is at 3 inches. There are no saline horizons. It is in nonirrigated land capability class 3w. This component is a hydric soil.

**KrB--Kinkora Silt Loam, 3 To 8 Percent Slopes**

Kinkora component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .43. This soil is poorly drained. The slowest permeability within 60 inches is slow. Available water capacity is very high and shrink swell potential is moderate. This soil is rarely flooded and is not ponded. The top of the seasonal high water table is at 3 inches. There are no saline horizons. It is in nonirrigated land capability class 3w. This component is a hydric soil.

**LeB2--Legore Silt Loam, 3 To 8 Percent Slopes Moderately Eroded**

Legore component makes up 100 percent of the map unit. All areas are prime farmland. The assigned Kw erodibility factor is .32. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 2e. This component is not a hydric soil.

**LeC2--Legore Silt Loam, 8 To 15 Percent Slopes, Moderately Eroded**

Legore component makes up 100 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .32. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 3e. This component is not a hydric soil.

**LeD2--Legore Silt Loam, 15 To 25 Percent Slopes, Moderately Eroded**

Legore component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .32. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 4e. This component is not a hydric soil.

**LeE--Legore Silt Loam, 25 To 45 Percent Slopes**

Legore component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .32. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 7e. This component is not a hydric soil.

**LfC--Legore Very Stony Silt Loam, 0 To 15 Percent Slopes**

Legore component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .24. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 6s. This component is not a hydric soil.

**LfD--Legore Very Stony Silt Loam, 15 To 25 Percent Slopes**

Legore component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .24. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 6s. This component is not a hydric soil.

**LfE--Legore Very Stony Silt Loam, 25 To 45 Percent Slopes**

Legore component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .24. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 7s. This component is not a hydric soil.

## NONTECHNICAL SOIL DESCRIPTIONS--Continued

---

**LgC3--Legore Silty Clay Loam, 8 To 15 Percent Slopes, Severely Eroded**

Legore component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .32. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 4e. This component is not a hydric soil.

**LgD3--Legore Silty Clay Loam 15 To 25 Percent Slopes, Severely Eroded**

Legore component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .32. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 6e. This component is not a hydric soil.

**Lr--Leonardtown Silt Loam**

Leonardtown component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .43. This soil is poorly drained. The slowest permeability within 60 inches is slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 6 inches. There are no saline horizons. It is in nonirrigated land capability class 4w. This component is a hydric soil.

**LyB--Loamy And Clayey Land, 0 To 5 Percent Slopes**

Loamy And Clayey Lan component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .17. This soil is well drained. The slowest permeability within 60 inches is slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. It is in nonirrigated land capability class 2s. This component is not a hydric soil.

**LyD--Loamy And Clayey Land, 5 To 15 Percent Slopes**

Loamy And Clayey Lan component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .17. This soil is well drained. The slowest permeability within 60 inches is slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. It is in nonirrigated land capability class 4e. This component is not a hydric soil.

**LyE--Loamy And Clayey Land, 15 To 30 Percent Slopes**

Loamy And Clayey Lan component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .17. This soil is well drained. The slowest permeability within 60 inches is slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. It is in nonirrigated land capability class 7e. This component is not a hydric soil.

**MbB2--Manor Loam, 3 To 8 Percent Slopes, Moderately Eroded**

Manor component makes up 100 percent of the map unit. All areas are prime farmland. The assigned Kw erodibility factor is .37. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 2e. This component is not a hydric soil.

**MbC2--Manor Loam, 8 To 15 Percent Slopes, Moderately Eroded**

Manor component makes up 100 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .37. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 3e. This component is not a hydric soil.

**MbC3--Manor Loam, 8 To 15 Percent Slopes, Severely Eroded**

Manor component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .37. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 4e. This component is not a hydric soil.

**MbD2--Manor Loam, 15 To 25 Percent Slopes, Moderately Eroded**

Manor component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .37. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 4e. This component is not a hydric soil.

## NONTECHNICAL SOIL DESCRIPTIONS--Continued

---

**MbD3--Manor Loam, 15 To 25 Percent Slopes, Severely Eroded**

Manor component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .37. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 6e. This component is not a hydric soil.

**McB2--Manor Channery Loam, 3 To 8 Percent Slopes Moderately Eroded**

Manor component makes up 100 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .37. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 2e. This component is not a hydric soil.

**McC2--Manor Channery Loam, 8 To 15 Percent Slopes, Moderately Eroded**

Manor component makes up 100 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .37. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 3e. This component is not a hydric soil.

**McC3--Manor Channery Loam, 8 To 15 Percent Slopes, Severely Eroded**

Manor component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .37. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 4e. This component is not a hydric soil.

**McD2--Manor Channery Loam, 15 To 25 Percent Slopes Moderately Eroded**

Manor component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .37. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 4e. This component is not a hydric soil.

**McD3--Manor Channery Loam, 15 To 25 Percent Slopes, Severely Eroded**

Manor component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .37. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 6e. This component is not a hydric soil.

**MdE--Manor Very Stony Loam, 25 To 45 Percent Slopes**

Manor component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .32. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 7s. This component is not a hydric soil.

**MfE--Manor Soils, 25 To 45 Percent Slopes**

Manor component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .37. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 7e. This component is not a hydric soil.

**MgC--Manor And Glenelg Very Stony Loams, 3 To 15 Percent Slopes**

Manor component makes up 50 percent of the map unit. The assigned Kw erodibility factor is .32. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 6s. This component is not a hydric soil.

Glenelg component makes up 50 percent of the map unit. The assigned Kw erodibility factor is .32. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 6s. This component is not a hydric soil.

## NONTECHNICAL SOIL DESCRIPTIONS--Continued

---

**MgD--Manor And Glenelg Very Stony Loams, 15 To 25 Percent Slopes**

Manor component makes up 50 percent of the map unit. The assigned Kw erodibility factor is .32. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 6s. This component is not a hydric soil.

Glenelg component makes up 50 percent of the map unit. The assigned Kw erodibility factor is .32. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 6s. This component is not a hydric soil.

**MkA--Matapeake Silt Loam, 0 To 2 Percent Slopes**

Matapeake component makes up 100 percent of the map unit. All areas are prime farmland. The assigned Kw erodibility factor is .49. This soil is well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 1. This component is not a hydric soil.

**MkB--Matapeake Silt Loam, 2 To 5 Percent Slopes**

Matapeake component makes up 100 percent of the map unit. All areas are prime farmland. The assigned Kw erodibility factor is .49. This soil is well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 2e. This component is not a hydric soil.

**MLA--Mattapex Silt Loam, 0 To 2 Percent Slopes**

Mattapex component makes up 95 percent of the map unit. All areas are prime farmland. The assigned Kw erodibility factor is .43. This soil is moderately well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 27 inches. There are no saline horizons. It is in nonirrigated land capability class 2w. This component is not a hydric soil.

**MLB--Mattapex Silt Loam, 2 To 5 Percent Slopes**

Mattapex component makes up 95 percent of the map unit. All areas are prime farmland. The assigned Kw erodibility factor is .43. This soil is moderately well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 27 inches. There are no saline horizons. It is in nonirrigated land capability class 2e. This component is not a hydric soil.

**MsA--Montalto Silt Loam, 0 To 3 Percent Slopes**

Montalto component makes up 100 percent of the map unit. All areas are prime farmland. The assigned Kw erodibility factor is .32. This soil is well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is high. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 1. This component is not a hydric soil.

**MsB2--Montalto Silt Loam, 3 To 8 Percent Slopes, Moderately Eroded**

Montalto component makes up 100 percent of the map unit. All areas are prime farmland. The assigned Kw erodibility factor is .32. This soil is well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is high. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 2e. This component is not a hydric soil.

**MsC2--Montalto Silt Loam, 8 To 15 Percent Slopes, Moderately Eroded**

Montalto component makes up 100 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .32. This soil is well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is high. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 3e. This component is not a hydric soil.

**NeA--Neshaminy Silt Loam, 0 To 3 Percent Slopes**

Neshaminy component makes up 100 percent of the map unit. All areas are prime farmland. The assigned Kw erodibility factor is .32. The depth to a restrictive feature is greater than 60 inches to bedrock. This soil is well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 1. This component is not a hydric soil.

## NONTECHNICAL SOIL DESCRIPTIONS--Continued

---

**NeB2--Neshaminy Silt Loam, 3 To 8 Percent Slopes, Moderately Eroded**

Neshaminy component makes up 100 percent of the map unit. All areas are prime farmland. The assigned Kw erodibility factor is .32. The depth to a restrictive feature is greater than 60 inches to bedrock. This soil is well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 2e. This component is not a hydric soil.

**NeC2--Neshaminy Silt Loam, 8 To 15 Percent Slopes, Moderately Eroded**

Neshaminy component makes up 100 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .32. The depth to a restrictive feature is greater than 60 inches to bedrock. This soil is well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 3e. This component is not a hydric soil.

**NsC--Neshaminy And Montalto Very Stony Silt Loams 0 To 15 Percent Slopes**

Neshaminy component makes up 50 percent of the map unit. The assigned Kw erodibility factor is .24. The depth to a restrictive feature is 48 inches to bedrock (lithic). This soil is well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 6s. This component is not a hydric soil.

Montalto component makes up 50 percent of the map unit. The assigned Kw erodibility factor is .32. This soil is well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is high. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 6s. This component is not a hydric soil.

**NsD--Neshaminy And Montalto Very Stony Silt Loams, 15 To 25 Percent Slopes**

Neshaminy component makes up 50 percent of the map unit. The assigned Kw erodibility factor is .24. The depth to a restrictive feature is 48 inches to bedrock (lithic). This soil is well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 6s. This component is not a hydric soil.

Montalto component makes up 50 percent of the map unit. The assigned Kw erodibility factor is .32. This soil is well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is high. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 7s. This component is not a hydric soil.

**NsE--Neshaminy And Montalto Very Stony Silt Loams, 25 To 45 Percent Slopes**

Neshaminy component makes up 50 percent of the map unit. The assigned Kw erodibility factor is .24. The depth to a restrictive feature is 48 inches to bedrock (lithic). This soil is well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 7s. This component is not a hydric soil.

Montalto component makes up 50 percent of the map unit. The assigned Kw erodibility factor is .32. This soil is well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is high. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 7s. This component is not a hydric soil.

**Ot--Othello Silt Loam**

Othello component makes up 100 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .37. This soil is poorly drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 6 inches. There are no saline horizons. It is in nonirrigated land capability class 3w. This component is a hydric soil.

**Sa--Sand And Gravel Pits**

Sand And Gravel Pits component makes up 95 percent of the map unit. The assigned Kw erodibility factor is .02. The slowest permeability within 60 inches is rapid. Available water capacity is low and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 8s. This component is not a hydric soil.

## NONTECHNICAL SOIL DESCRIPTIONS--Continued

---

ShB2--Sassafras Sandy Loam, 2 To 5 Percent Slopes, Moderately Eroded

Sassafras component makes up 100 percent of the map unit. All areas are prime farmland. The assigned Kw erodibility factor is .28. This soil is well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 2e. This component is not a hydric soil.

## ShC2--Sassafras Sandy Loam, 5 To 10 Percent Slopes, Moderately Eroded

Sassafras component makes up 100 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .28. This soil is well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 3e. This component is not a hydric soil.

## SlB2--Sassafras Loam, 2 To 5 Percent Slopes, Moderately Eroded

Sassafras component makes up 100 percent of the map unit. All areas are prime farmland. The assigned Kw erodibility factor is .28. This soil is well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 2e. This component is not a hydric soil.

## SlC2--Sassafras Loam, 5 To 10 Percent Slopes, Moderately Eroded

Sassafras component makes up 100 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .28. This soil is well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 3e. This component is not a hydric soil.

## SsD--Sassafras And Joppa Soils, 10 To 15 Percent Slopes

Sassafras component makes up 50 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .28. This soil is well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 4e. This component is not a hydric soil.

Joppa component makes up 50 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .28. This soil is well drained. The slowest permeability within 60 inches is moderately rapid. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 4e. This component is not a hydric soil.

## SsE--Sassafras And Joppa Soils, 15 To 30 Percent Slopes

Sassafras component makes up 50 percent of the map unit. The assigned Kw erodibility factor is .28. This soil is well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 7e. This component is not a hydric soil.

Joppa component makes up 50 percent of the map unit. The assigned Kw erodibility factor is .28. This soil is well drained. The slowest permeability within 60 inches is moderately rapid. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 6e. This component is not a hydric soil.

## St--Stony Land, Steep

Stony Land component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .32. This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 7s. This component is not a hydric soil.

## Sw--Swamp

Swamp component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .05. This soil is very poorly drained. The slowest permeability within 60 inches is moderately rapid. Available water capacity is very high and shrink swell potential is low. This soil is frequently flooded and is not ponded. The top of the seasonal high water table is at 0 inches. There are no saline horizons. It is in nonirrigated land capability class 7w. This component is a hydric soil.

## NONTECHNICAL SOIL DESCRIPTIONS--Continued

---

**Tm--Tidal Marsh**

Tidal Marsh component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .43. This soil is poorly drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is frequently flooded and is not ponded. The top of the seasonal high water table is at 0 inches. The soil has a moderately saline horizon. It is in nonirrigated land capability class 8w. This component is a hydric soil.

**WaA--Watchung Silt Loam, 0 To 3 Percent Slopes**

Watchung component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .43. This soil is poorly drained. The slowest permeability within 60 inches is slow. Available water capacity is very high and shrink swell potential is moderate. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 6 inches. There are no saline horizons. It is in nonirrigated land capability class 4w. This component is a hydric soil.

**WaB--Watchung Silt Loam, 3 To 8 Percent Slopes**

Watchung component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .43. This soil is poorly drained. The slowest permeability within 60 inches is slow. Available water capacity is very high and shrink swell potential is moderate. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 6 inches. There are no saline horizons. It is in nonirrigated land capability class 6w. This component is a hydric soil.

**WcB--Watchung Very Stony Silt Loam, 0 To 8 Percent Slopes**

Watchung component makes up 100 percent of the map unit. The assigned Kw erodibility factor is .43. This soil is poorly drained. The slowest permeability within 60 inches is slow. Available water capacity is very high and shrink swell potential is moderate. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 6 inches. There are no saline horizons. It is in nonirrigated land capability class 7s. This component is a hydric soil.

**WhB--Whiteford Silt Loam, 3 To 8 Percent Slopes**

Whiteford component makes up 100 percent of the map unit. All areas are prime farmland. The assigned Kw erodibility factor is .32. The depth to a restrictive feature is 40 to 60 inches to bedrock (lithic). This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 2e. This component is not a hydric soil.

**WhC2--Whiteford Silt Loam, 8 To 15 Percent Slopes, Moderately Eroded**

Whiteford component makes up 100 percent of the map unit. Farmland of statewide importance. The assigned Kw erodibility factor is .32. The depth to a restrictive feature is 40 to 60 inches to bedrock (lithic). This soil is well drained. The slowest permeability within 60 inches is moderate. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The water table is deeper than 6 feet. There are no saline horizons. It is in nonirrigated land capability class 3e. This component is not a hydric soil.

**WoB--Woodstown Loam, 0 To 5 Percent Slopes**

Woodstown component makes up 95 percent of the map unit. All areas are prime farmland. The assigned Kw erodibility factor is .32. This soil is moderately well drained. The slowest permeability within 60 inches is moderately slow. Available water capacity is very high and shrink swell potential is low. This soil is not flooded and is not ponded. The top of the seasonal high water table is at 30 inches. There are no saline horizons. It is in nonirrigated land capability class 2w. This component is not a hydric soil.

