

Map Unit Description (MN)

Martin County, Minnesota

[Data apply to the entire extent of the map unit within the survey area. Map unit and soil properties for a specific parcel of land may vary somewhat and should be determined by onsite investigation]

8B--Sparta loamy fine sand, 1 to 6 percent slopes

Sparta

Extent: 90 percent of the unit

Landform(s): outwash plains

Slope gradient: 1 to 6 percent

Parent material: sandy outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: excessively drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .17

Land capability, nonirrigated 4s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 14 in	loamy fine sand	moderately rapid	1.28 to 1.70 in	5.1 to 7.3
Bw -- 14 to 30 in	loamy fine sand	rapid	0.79 to 1.73 in	5.1 to 7.3
C -- 30 to 60 in	fine sand	rapid	1.20 to 2.09 in	5.1 to 7.8

Farrer

Extent: 4 percent of the unit

Landform(s): outwash plains

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

8B--Sparta loamy fine sand, 1 to 6 percent slopes

Hanska

Extent: 3 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>

Litchfield

Extent: 3 percent of the unit

Landform(s): outwash plains

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>

Map Unit Description (MN)

Martin County, Minnesota

27B--Dickinson fine sandy loam, 1 to 6 percent slopes

Dickinson

Extent: 90 percent of the unit

Landform(s): outwash plains

Slope gradient: 1 to 6 percent

Parent material: coarse-loamy outwash over sandy outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .24

Land capability, nonirrigated 3e

Hydric soil: no

Hydrologic group: A

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 19 in	fine sandy loam	moderately rapid	2.27 to 2.83 in	5.6 to 7.3
Bw -- 19 to 34 in	sandy loam	moderately rapid	1.80 to 2.24 in	5.1 to 6.5
C -- 34 to 60 in	fine sand	rapid	2.08 to 2.60 in	5.1 to 6.5

Hanska

Extent: 5 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

27B--Dickinson fine sandy loam, 1 to 6 percent slopes

Clarion

Extent: 5 percent of the unit

Landform(s): moraines

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

Representative soil profile:

Texture

Permeability

*Available water
capacity*

pH

Map Unit Description (MN)

Martin County, Minnesota

35--Blue Earth mucky silty clay loam

Blue Earth

Extent: 85 percent of the unit

Landform(s): relict lakebeds on moraines

Slope gradient: 0 to 1 percent

Parent material: fine-silty coprogenic material

Restrictive feature(s): greater than 60 inches

Flooding: rare

Ponding: frequent

Drainage class: very poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .24

Land capability, nonirrigated 3w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap --	0 to 10 in mucky silty clay loam	moderate	1.77 to 2.36 in	7.4 to 8.4
C --	10 to 60 in mucky silty clay loam	moderate	9.00 to 12.00 in	7.4 to 8.4

Lemond

Extent: 10 percent of the unit

Landform(s): flats

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>

Map Unit Description (MN)

Martin County, Minnesota

35--Blue Earth mucky silty clay loam

Linder

Extent: 5 percent of the unit

Landform(s): outwash plains

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

Representative soil profile:

Texture

Permeability

*Available water
capacity*

pH

Map Unit Description (MN)

Martin County, Minnesota

37B--Farrar fine sandy loam, 1 to 6 percent slopes

Farrar

Extent: 90 percent of the unit

Landform(s): moraines

Slope gradient: 1 to 6 percent

Parent material: coarse-loamy eolian deposits over fine-loamy till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .15

Land capability, nonirrigated 2e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 20 in	fine sandy loam	moderately rapid	3.21 to 3.61 in	5.6 to 7.3
2C -- 20 to 60 in	loam	moderate	6.76 to 7.56 in	6.1 to 8.4

Estherville

Extent: 4 percent of the unit

Landform(s): outwash plains

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>

Map Unit Description (MN)

Martin County, Minnesota

37B--Farrar fine sandy loam, 1 to 6 percent slopes

Clarion

Extent: 3 percent of the unit

Landform(s): moraines

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Litchfield

Extent: 3 percent of the unit

Landform(s): outwash plains

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

39B--Wadena loam, 1 to 6 percent slopes

Wadena

Extent: 90 percent of the unit

Landform(s): outwash plains

Slope gradient: 1 to 6 percent

Parent material: fine-loamy outwash over sandy and gravelly outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 2e

Hydric soil: no

Hydrologic group: B

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 20 in	loam	moderate	4.02 to 4.42 in	6.1 to 7.3
Bw,BC -- 20 to 38 in	loam	moderate	2.48 to 3.37 in	5.6 to 7.3
2C -- 38 to 60 in	sand	very rapid	0.44 to 0.88 in	6.6 to 8.4

Dickinson

Extent: 4 percent of the unit

Landform(s): outwash plains

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

39B--Wadena loam, 1 to 6 percent slopes

Hanska

Extent: 3 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>

Biscay

Extent: 3 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>

Map Unit Description (MN)

Martin County, Minnesota

41B--Estherville sandy loam, 1 to 6 percent slopes

Estherville

Extent: 90 percent of the unit

Landform(s): outwash plains

Slope gradient: 1 to 6 percent

Parent material: coarse-loamy outwash over sandy and gravelly outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat excessively drained

Soil loss tolerance (T factor): 2

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .24

Land capability, nonirrigated 3s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 13 in	sandy loam	moderately rapid	1.69 to 2.34 in	5.6 to 7.3
Bw -- 13 to 18 in	sandy loam	moderately rapid	0.67 to 0.92 in	5.6 to 7.3
2C -- 18 to 60 in	gravelly coarse sand	rapid	0.83 to 1.67 in	6.6 to 8.4

Dickinson

Extent: 4 percent of the unit

Landform(s): outwash plains

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

41B--Estherville sandy loam, 1 to 6 percent slopes

Hanska

Extent: 3 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Biscay

Extent: 3 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

41C--Estherville sandy loam, 6 to 12 percent slopes

Estherville

Extent: 90 percent of the unit
Landform(s): outwash plains
Slope gradient: 6 to 12 percent
Parent material: coarse-loamy outwash over sandy and gravelly outwash
Restrictive feature(s): greater than 60 inches
Flooding: none
Ponding: none
Drainage class: somewhat excessively drained

Soil loss tolerance (T factor): 2
Wind erodibility group (WEG): 3
Wind erodibility index (WEI): 86
Kw factor (surface layer) .24

Land capability, nonirrigated 4s
Hydric soil: no
Hydrologic group: A
Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 13 in	sandy loam	moderately rapid	1.69 to 2.34 in	5.6 to 7.3
Bw -- 13 to 18 in	sandy loam	moderately rapid	0.67 to 0.92 in	5.6 to 7.3
2C -- 18 to 60 in	gravelly coarse sand	rapid	0.83 to 1.67 in	6.6 to 8.4

Dickinson

Extent: 4 percent of the unit
Landform(s): outwash plains
Slope gradient:
Parent material:
Restrictive feature(s): greater than 60 inches
Flooding:
Ponding:
Drainage class:

Soil loss tolerance (T factor):
Wind erodibility group (WEG):
Wind erodibility index (WEI):
Kw factor (surface layer)
Land capability, nonirrigated
Hydric soil: no
Hydrologic group:
Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

41C--Estherville sandy loam, 6 to 12 percent slopes

Clarion

Extent: 3 percent of the unit

Landform(s): moraines

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Swanlake

Extent: 3 percent of the unit

Landform(s): moraines

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

84--Brownton silty clay

Brownton

Extent: 90 percent of the unit

Landform(s): rims on moraines

Slope gradient: 0 to 1 percent

Parent material: clayey lacustrine deposits over fine-loamy till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4

Wind erodibility index (WEI): 86

Kw factor (surface layer) .28

Land capability, nonirrigated 2w

Hydric soil: yes

Hydrologic group: C/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 20 in	silty clay	slow	3.61 to 4.42 in	7.4 to 8.4
C -- 20 to 60 in	clay	slow	5.17 to 6.36 in	7.4 to 8.4

Waldorf

Extent: 5 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>

Map Unit Description (MN)

Martin County, Minnesota

84--Brownton silty clay

Okoboji

Extent: 5 percent of the unit

Landform(s): depressions

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

Representative soil profile:

Texture

Permeability

*Available water
capacity*

pH

Map Unit Description (MN)

Martin County, Minnesota

86--Canisteo clay loam

Canisteo

Extent: 90 percent of the unit

Landform(s): rims on depressions on moraines, flats on moraines

Slope gradient: 0 to 2 percent

Parent material: fine-loamy till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .24

Land capability, nonirrigated 2w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 9 in	clay loam	moderate	1.63 to 1.99 in	7.4 to 8.4
A -- 9 to 23 in	clay loam	moderate	2.07 to 2.62 in	7.4 to 8.4
Bk -- 23 to 36 in	clay loam	moderate	1.56 to 2.34 in	7.4 to 8.4
C -- 36 to 60 in	loam	moderate	3.36 to 3.84 in	7.4 to 8.4

Webster

Extent: 5 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

86--Canisteo clay loam

Glencoe

Extent: 5 percent of the unit

Landform(s): depressions

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

Representative soil profile:

Texture

Permeability

*Available water
capacity*

pH

Map Unit Description (MN)

Martin County, Minnesota

94B--Terril loam, 2 to 6 percent slopes

Terril

Extent: 90 percent of the unit
Landform(s): hills on moraines
Slope gradient: 2 to 6 percent
Parent material: fine-loamy colluvium
Restrictive feature(s): greater than 60 inches
Flooding: none
Ponding: none
Drainage class: moderately well drained

Soil loss tolerance (T factor): 5
Wind erodibility group (WEG): 6
Wind erodibility index (WEI): 48
Kw factor (surface layer) .28
Land capability, nonirrigated 2e
Hydric soil: no
Hydrologic group: C
Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A1 -- 0 to 15 in	loam	moderate	2.99 to 3.29 in	6.1 to 7.3
A2 -- 15 to 36 in	clay loam	moderate	3.55 to 3.96 in	6.1 to 7.3
Bw -- 36 to 60 in	loam	moderate	3.84 to 4.32 in	6.1 to 7.8

Clarion

Extent: 4 percent of the unit
Landform(s): moraines
Slope gradient:
Parent material:
Restrictive feature(s): greater than 60 inches
Flooding:
Ponding:
Drainage class:

Soil loss tolerance (T factor):
Wind erodibility group (WEG):
Wind erodibility index (WEI):
Kw factor (surface layer)
Land capability, nonirrigated
Hydric soil: no
Hydrologic group:
Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

94B--Terril loam, 2 to 6 percent slopes

Swanlake

Extent: 3 percent of the unit

Landform(s): moraines

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Webster

Extent: 3 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

96--Collinwood silty clay

Collinwood

Extent: 90 percent of the unit

Landform(s): hills on lake plains

Slope gradient: 0 to 3 percent

Parent material: clayey lacustrine deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4

Wind erodibility index (WEI): 86

Kw factor (surface layer) .24

Land capability, nonirrigated 2w

Hydric soil: no

Hydrologic group: C

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 15 in	silty clay	moderately slow	2.09 to 2.54 in	5.6 to 7.3
Bw -- 15 to 35 in	clay	moderately slow	2.61 to 3.21 in	5.6 to 7.3
C -- 35 to 60 in	silty clay	moderately slow	2.73 to 3.72 in	7.4 to 8.4

Waldorf

Extent: 5 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

96--Collinwood silty clay

Truman

Extent: 5 percent of the unit

Landform(s): lake plains

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

Representative soil profile:

Texture

Permeability

*Available water
capacity*

pH

Map Unit Description (MN)

Martin County, Minnesota

101B--Truman silt loam, 1 to 6 percent slopes

Truman

Extent: 90 percent of the unit

Landform(s): hills on lake plains

Slope gradient: 1 to 6 percent

Parent material: fine-silty lacustrine deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .37

Land capability, nonirrigated 2e

Hydric soil: no

Hydrologic group: B

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 13 in	silt loam	moderate	2.60 to 2.99 in	5.6 to 7.3
Bw -- 13 to 33 in	silt loam	moderate	3.61 to 4.22 in	5.6 to 7.8
C -- 33 to 60 in	silt loam	moderate	4.82 to 5.35 in	7.4 to 8.4

Bold

Extent: 4 percent of the unit

Landform(s): lake plains

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

101B--Truman silt loam, 1 to 6 percent slopes

Madelia

Extent: 3 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Kingston

Extent: 3 percent of the unit

Landform(s): lake plains

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

102B--Clarion loam, 1 to 6 percent slopes

Clarion

Extent: 90 percent of the unit

Landform(s): hills on moraines

Slope gradient: 1 to 6 percent

Parent material: fine-loamy till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 2e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 14 in	loam	moderate	2.83 to 3.12 in	5.6 to 7.3
Bw -- 14 to 33 in	loam	moderate	3.21 to 3.59 in	5.6 to 7.8
C -- 33 to 60 in	loam	moderate	4.55 to 5.09 in	7.4 to 8.4

Swanlake

Extent: 4 percent of the unit

Landform(s): moraines

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

102B--Clarion loam, 1 to 6 percent slopes

Webster

Extent: 3 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Storden

Extent: 3 percent of the unit

Landform(s): moraines

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

106B--Lester loam, 2 to 6 percent slopes

Lester

Extent: 90 percent of the unit

Landform(s): hills on moraines

Slope gradient: 2 to 5 percent

Parent material: fine-loamy till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 2e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 19 in	loam	moderate	3.78 to 4.16 in	5.6 to 7.3
Bt -- 19 to 41 in	clay loam	moderate	3.31 to 4.19 in	5.6 to 7.3
C -- 41 to 60 in	loam	moderate	2.65 to 3.40 in	7.4 to 7.8

Swanlake

Extent: 4 percent of the unit

Landform(s): moraines

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

106B--Lester loam, 2 to 6 percent slopes

Webster

Extent: 3 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>

Nicollet

Extent: 3 percent of the unit

Landform(s): moraines

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>

Map Unit Description (MN)

Martin County, Minnesota

106C--Lester loam, 6 to 12 percent slopes

Lester

Extent: 90 percent of the unit

Landform(s): hills on moraines

Slope gradient: 6 to 12 percent

Parent material: fine-loamy till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .32

Land capability, nonirrigated 3e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 9 in	loam	moderate	1.81 to 1.99 in	5.6 to 7.3
Bt -- 9 to 41 in	clay loam	moderate	4.78 to 6.06 in	5.1 to 7.3
C -- 41 to 60 in	loam	moderate	2.65 to 3.59 in	7.4 to 8.4

Storden

Extent: 4 percent of the unit

Landform(s): moraines

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

106C--Lester loam, 6 to 12 percent slopes

Webster

Extent: 3 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>

Nicollet

Extent: 3 percent of the unit

Landform(s): moraines

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>

Map Unit Description (MN)

Martin County, Minnesota

112--Harps clay loam

Harps

Extent: 90 percent of the unit

Landform(s): rims on moraines

Slope gradient: 0 to 2 percent

Parent material: fine-loamy till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .24

Land capability, nonirrigated 2w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,Ak -- 0 to 14 in	clay loam	moderate	2.69 to 2.98 in	7.9 to 8.4
Bgk -- 14 to 33 in	clay loam	moderate	3.21 to 3.59 in	7.9 to 8.4
C -- 33 to 60 in	loam	moderate	3.75 to 5.09 in	7.4 to 8.4

Glencoe

Extent: 4 percent of the unit

Landform(s): depressions

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

112--Harps clay loam

Crippen

Extent: 3 percent of the unit

Landform(s): moraines

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Okoboji

Extent: 3 percent of the unit

Landform(s): depressions

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

113--Webster clay loam

Webster

Extent: 90 percent of the unit

Landform(s): flats on moraines

Slope gradient: 0 to 2 percent

Parent material: fine-loamy till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .24

Land capability, nonirrigated 2w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 19 in	clay loam	moderate	3.59 to 3.97 in	6.6 to 7.3
Bg -- 19 to 34 in	clay loam	moderate	2.39 to 2.69 in	6.6 to 7.8
Cg -- 34 to 60 in	loam	moderate	3.64 to 4.94 in	7.4 to 8.4

Glencoe

Extent: 5 percent of the unit

Landform(s): depressions

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

113--Webster clay loam

Canisteo

Extent: 5 percent of the unit

Landform(s): rims

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

Representative soil profile:

Texture

Permeability

*Available water
capacity*

pH

Map Unit Description (MN)

Martin County, Minnesota

114--Glencoe clay loam

Glencoe

Extent: 85 percent of the unit

Landform(s): depressions on moraines

Slope gradient: 0 to 1 percent

Parent material: fine-loamy alluvium

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: frequent

Drainage class: very poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .24

Land capability, nonirrigated 3w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 41 in	clay loam	moderate	7.37 to 9.01 in	6.1 to 7.8
Bg -- 41 to 53 in	silty clay loam	moderate	2.20 to 2.69 in	6.1 to 7.8
Cg -- 53 to 60 in	silty clay loam	moderate	1.00 to 1.27 in	6.6 to 7.8

Canisteo

Extent: 10 percent of the unit

Landform(s): rims

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

114--Glencoe clay loam

Webster

Extent: 5 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

Representative soil profile:

Texture

Permeability

*Available water
capacity*

pH

Map Unit Description (MN)

Martin County, Minnesota

118--Crippin loam

Crippin

Extent: 90 percent of the unit

Landform(s): rises on moraines

Slope gradient: 0 to 3 percent

Parent material: fine-loamy till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .24

Land capability, nonirrigated 1

Hydric soil: no

Hydrologic group: B/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 15 in	loam	moderate	2.99 to 3.29 in	6.6 to 8.4
Bw -- 15 to 27 in	loam	moderate	2.01 to 2.24 in	7.4 to 8.4
C -- 27 to 60 in	loam	moderate	5.62 to 6.28 in	7.9 to 8.4

Swanlake

Extent: 4 percent of the unit

Landform(s): moraines

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

118--Crippin loam

Clarion

Extent: 3 percent of the unit

Landform(s): moraines

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Harps

Extent: 3 percent of the unit

Landform(s): rims

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

128B--Grogan silt loam, 1 to 6 percent slopes

Grogan

Extent: 90 percent of the unit

Landform(s): lake plains

Slope gradient: 1 to 6 percent

Parent material: sandy and silty glaciolacustrine deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .43

Land capability, nonirrigated 2e

Hydric soil: no

Hydrologic group: A

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 18 in	silt loam	moderately rapid	3.98 to 4.35 in	5.6 to 7.3
Bw -- 18 to 38 in	silt loam	moderately rapid	3.35 to 3.74 in	6.1 to 7.8
C -- 38 to 60 in	stratified very fine sandy loam to silt loam	moderately rapid	3.75 to 4.19 in	7.4 to 8.4

Clarion

Extent: 4 percent of the unit

Landform(s): moraines

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

128B--Grogan silt loam, 1 to 6 percent slopes

Ocheyedan

Extent: 3 percent of the unit

Landform(s): lake plains

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Madelia

Extent: 3 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

130--Nicollet clay loam

Nicollet

Extent: 90 percent of the unit
Landform(s): rises on moraines
Slope gradient: 1 to 3 percent
Parent material: fine-loamy till
Restrictive feature(s): greater than 60 inches
Flooding: none
Ponding: none
Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 5
Wind erodibility group (WEG): 6
Wind erodibility index (WEI): 48
Kw factor (surface layer) .24
Land capability, nonirrigated 1
Hydric soil: no
Hydrologic group: B/D
Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 17 in	clay loam	moderate	2.88 to 3.72 in	5.6 to 7.3
Bw -- 17 to 36 in	clay loam	moderate	2.83 to 3.59 in	5.6 to 7.8
C -- 36 to 60 in	loam	moderate	3.36 to 4.56 in	7.4 to 8.4

Clarion

Extent: 5 percent of the unit
Landform(s): moraines
Slope gradient:
Parent material:
Restrictive feature(s): greater than 60 inches
Flooding:
Ponding:
Drainage class:

Soil loss tolerance (T factor):
Wind erodibility group (WEG):
Wind erodibility index (WEI):
Kw factor (surface layer)
Land capability, nonirrigated
Hydric soil: no
Hydrologic group:
Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

130--Nicollet clay loam

Webster

Extent: 5 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

Representative soil profile:

Texture

Permeability

*Available water
capacity*

pH

Map Unit Description (MN)

Martin County, Minnesota

134--Okoboji silty clay loam

Okoboji

Extent: 85 percent of the unit

Landform(s): depressions on moraines

Slope gradient: 0 to 1 percent

Parent material: fine-silty alluvium

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: frequent

Drainage class: very poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .28

Land capability, nonirrigated 3w

Hydric soil: yes

Hydrologic group: C/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 28 in	silty clay loam	moderately slow	5.87 to 6.43 in	6.1 to 7.8
Bg -- 28 to 42 in	silty clay loam	moderately slow	2.55 to 2.83 in	6.6 to 7.8
Cg -- 42 to 60 in	clay loam	moderately slow	3.19 to 3.54 in	6.6 to 8.4

Canisteo

Extent: 5 percent of the unit

Landform(s): rims

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

134--Okoboji silty clay loam

Harps

Extent: 5 percent of the unit

Landform(s): rims

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

Representative soil profile:

Texture

Permeability

*Available water
capacity*

pH

Palms

Extent: 5 percent of the unit

Landform(s): depressions

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

Representative soil profile:

Texture

Permeability

*Available water
capacity*

pH

Map Unit Description (MN)

Martin County, Minnesota

136--Madelia silty clay loam

Madelia

Extent: 90 percent of the unit

Landform(s): flats on lake plains

Slope gradient: 0 to 2 percent

Parent material: fine-silty lacustrine deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .32

Land capability, nonirrigated 2w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 22 in	silty clay loam	moderate	3.97 to 5.29 in	6.1 to 7.3
Bg -- 22 to 37 in	silty clay loam	moderate	2.39 to 3.29 in	6.6 to 7.8
Cg -- 37 to 60 in	silt loam	moderate	3.65 to 5.02 in	7.4 to 8.4

Harps

Extent: 5 percent of the unit

Landform(s): rims

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

136--Madelia silty clay loam

Kingston

Extent: 5 percent of the unit

Landform(s): lake plains

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

Representative soil profile:

Texture

Permeability

*Available water
capacity*

pH

Map Unit Description (MN)

Martin County, Minnesota

140--Spicer silty clay loam

Spicer

Extent: 90 percent of the unit

Landform(s): flats on lake plains

Slope gradient: 0 to 2 percent

Parent material: fine-silty lacustrine deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .32

Land capability, nonirrigated 2w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 24 in	silty clay loam	moderate	4.32 to 5.76 in	7.4 to 8.4
Bg -- 24 to 34 in	silty clay loam	moderate	1.57 to 2.17 in	7.4 to 8.4
Cg -- 34 to 60 in	silty clay loam	moderate	4.16 to 5.72 in	7.4 to 8.4

Kingston

Extent: 5 percent of the unit

Landform(s): lake plains

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

140--Spicer silty clay loam

Madelia

Extent: 5 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

Representative soil profile:

Texture

Permeability

*Available water
capacity*

pH

Map Unit Description (MN)

Martin County, Minnesota

181--Litchfield sandy loam

Litchfield

Extent: 90 percent of the unit

Landform(s): outwash plains

Slope gradient: 0 to 3 percent

Parent material: coarse-loamy outwash over sandy outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .17

Land capability, nonirrigated 3s

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 16 in	sandy loam	moderately rapid	1.94 to 2.42 in	5.6 to 7.3
Bw -- 16 to 37 in	stratified loamy sand to fine sandy loam	moderately rapid	1.46 to 3.34 in	5.6 to 7.8
C -- 37 to 60 in	sand	rapid	1.83 to 2.28 in	7.4 to 8.4

Dickinson

Extent: 4 percent of the unit

Landform(s): outwash plains

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

181--Litchfield sandy loam

Hanska

Extent: 3 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

Representative soil profile:

Texture

Permeability

*Available water
capacity*

pH

Lemond

Extent: 3 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

Representative soil profile:

Texture

Permeability

*Available water
capacity*

pH

Map Unit Description (MN)

Martin County, Minnesota

197--Kingston silt loam

Kingston

Extent: 90 percent of the unit

Landform(s): rises on lake plains

Slope gradient: 0 to 3 percent

Parent material: fine-silty lacustrine deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .37

Land capability, nonirrigated 1

Hydric soil: no

Hydrologic group: C

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 11 in	silt loam	moderate	1.98 to 2.65 in	5.6 to 7.3
Bw -- 11 to 29 in	silty clay loam	moderate	2.90 to 3.62 in	5.6 to 7.8
C -- 29 to 60 in	silt loam	moderate	4.91 to 6.14 in	7.4 to 8.4

Bold

Extent: 5 percent of the unit

Landform(s): lake plains

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

197--Kingston silt loam

Madelia

Extent: 5 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

Representative soil profile:

Texture

Permeability

*Available water
capacity*

pH

Map Unit Description (MN)

Martin County, Minnesota

227--Lemond sandy loam

Lemond

Extent: 90 percent of the unit

Landform(s): flats on outwash plains

Slope gradient: 0 to 2 percent

Parent material: coarse-loamy outwash over sandy outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .17

Land capability, nonirrigated 2w

Hydric soil: yes

Hydrologic group: A/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 20 in	sandy loam	moderately rapid	3.01 to 3.61 in	7.4 to 8.4
Bg -- 20 to 30 in	sandy loam	moderately rapid	0.98 to 1.28 in	7.4 to 8.4
2Cg -- 30 to 60 in	sand	rapid	1.50 to 2.09 in	7.4 to 8.4

Hanska

Extent: 4 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

227--Lemond sandy loam

Litchfield

Extent: 3 percent of the unit

Landform(s): outwash plains

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>

Webster

Extent: 3 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>

Map Unit Description (MN)

Martin County, Minnesota

229--Waldorf silty clay loam

Waldorf

Extent: 90 percent of the unit

Landform(s): flats on lake plains

Slope gradient: 0 to 2 percent

Parent material: clayey lacustrine deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4

Wind erodibility index (WEI): 86

Kw factor (surface layer) .24

Land capability, nonirrigated 2w

Hydric soil: yes

Hydrologic group: C/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 24 in	silty clay loam	moderate	4.32 to 6.00 in	6.1 to 7.3
Bg -- 24 to 36 in	silty clay	moderately slow	1.54 to 1.89 in	6.6 to 7.8
Cg -- 36 to 60 in	silty clay loam	moderate	4.80 to 5.28 in	7.4 to 8.4

Brownnton

Extent: 4 percent of the unit

Landform(s): rims

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

229--Waldorf silty clay loam

Okoboji

Extent: 3 percent of the unit

Landform(s): depressions

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>

Collinwood

Extent: 3 percent of the unit

Landform(s): lake plains

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>

Map Unit Description (MN)

Martin County, Minnesota

247--Linder loam

Linder

Extent: 90 percent of the unit

Landform(s): outwash plains

Slope gradient: 0 to 2 percent

Parent material: coarse-loamy outwash over sandy outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .32

Land capability, nonirrigated 2s

Hydric soil: no

Hydrologic group: B/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 15 in	loam	moderate	2.99 to 3.29 in	5.6 to 7.8
Bw -- 15 to 29 in	sandy loam	moderately rapid	2.13 to 2.41 in	6.1 to 7.8
2C -- 29 to 60 in	coarse sand	very rapid	0.61 to 1.23 in	7.4 to 8.4

Biscay

Extent: 4 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

247--Linder loam

Dickinson

Extent: 2 percent of the unit

Landform(s): outwash plains

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Wadena

Extent: 2 percent of the unit

Landform(s): outwash plains

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

247--Linder loam

Mayer

Extent: 2 percent of the unit

Landform(s): rims

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

Representative soil profile:

Texture

Permeability

*Available water
capacity*

pH

Map Unit Description (MN)

Martin County, Minnesota

255--Mayer loam

Mayer

Extent: 90 percent of the unit

Landform(s): flats on outwash plains

Slope gradient: 0 to 2 percent

Parent material: fine-loamy outwash over sandy outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .20

Land capability, nonirrigated 2w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 16 in	loam	moderate	3.23 to 3.55 in	7.4 to 8.4
Bg -- 16 to 30 in	loam	moderate	2.20 to 2.62 in	7.4 to 8.4
2Cg -- 30 to 60 in	sand	rapid	0.60 to 1.20 in	7.4 to 8.4

Linder

Extent: 5 percent of the unit

Landform(s): outwash plains

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

255--Mayer loam

Wadena

Extent: 5 percent of the unit

Landform(s): outwash plains

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

Representative soil profile:

Texture

Permeability

*Available water
capacity*

pH

Map Unit Description (MN)

Martin County, Minnesota

269--Millington clay loam, occasionally flooded

Millington, occasionally flooded

Extent: 90 percent of the unit

Landform(s): flood plains

Slope gradient: 0 to 2 percent

Parent material: fine-loamy alluvium

Restrictive feature(s): greater than 60 inches

Flooding: occasional

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .24

Land capability, nonirrigated 2w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 28 in	clay loam	moderate	4.75 to 6.43 in	7.4 to 8.4
Bg -- 28 to 39 in	clay loam	moderate	1.87 to 2.20 in	7.4 to 8.4
Cg -- 39 to 60 in	stratified loam to clay loam	moderate	2.92 to 4.17 in	7.4 to 8.4

Spillville

Extent: 5 percent of the unit

Landform(s): flood plains

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

269--Millington clay loam, occasionally flooded

Coland

Extent: 5 percent of the unit

Landform(s): flood plains

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

Representative soil profile:

Texture

Permeability

*Available water
capacity*

pH

Map Unit Description (MN)

Martin County, Minnesota

275B--Ocheyedan loam, 1 to 5 percent slopes

Ocheyedan

Extent: 90 percent of the unit

Landform(s): hills on moraines

Slope gradient: 1 to 5 percent

Parent material: loamy till over silty glaciolacustrine deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 2e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A,AB -- 0 to 15 in	loam	moderate	2.99 to 3.29 in	5.6 to 7.3
2Bw -- 15 to 22 in	loam	moderate	1.13 to 1.28 in	6.1 to 7.8
2C -- 22 to 60 in	silt loam	moderate	7.18 to 7.94 in	6.6 to 8.4

Kingston

Extent: 5 percent of the unit

Landform(s): lake plains

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

275B--Ocheyedan loam, 1 to 5 percent slopes

Fostoria

Extent: 5 percent of the unit

Landform(s): moraines

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

Representative soil profile:

Texture

Permeability

*Available water
capacity*

pH

Map Unit Description (MN)

Martin County, Minnesota

282--Hanska loam

Hanska

Extent: 90 percent of the unit
Landform(s): flats on outwash plains
Slope gradient: 0 to 1 percent
Parent material: coarse-loamy outwash over sandy outwash
Restrictive feature(s): greater than 60 inches
Flooding: none
Ponding: none
Drainage class: poorly drained

Soil loss tolerance (T factor): 3
Wind erodibility group (WEG): 5
Wind erodibility index (WEI): 56
Kw factor (surface layer) .32
Land capability, nonirrigated 2w
Hydric soil: yes
Hydrologic group: A/D
Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 21 in	loam	moderately rapid	4.17 to 4.59 in	6.1 to 7.8
Bg -- 21 to 34 in	sandy loam	moderately rapid	1.30 to 1.69 in	6.1 to 7.3
2C -- 34 to 60 in	sand	rapid	0.78 to 1.30 in	6.6 to 7.8

Litchfield

Extent: 5 percent of the unit
Landform(s): outwash plains
Slope gradient:
Parent material:
Restrictive feature(s): greater than 60 inches
Flooding:
Ponding:
Drainage class:

Soil loss tolerance (T factor):
Wind erodibility group (WEG):
Wind erodibility index (WEI):
Kw factor (surface layer)
Land capability, nonirrigated
Hydric soil: no
Hydrologic group:
Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

282--Hanska loam

Lemond

Extent: 5 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

Representative soil profile:

Texture

Permeability

*Available water
capacity*

pH

Map Unit Description (MN)

Martin County, Minnesota

313--Spillville loam, occasionally flooded

Spillville, occasionally flooded

Extent: 90 percent of the unit

Landform(s): flood plains

Slope gradient: 0 to 2 percent

Parent material: fine-loamy alluvium

Restrictive feature(s): greater than 60 inches

Flooding: occasional

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 2w

Hydric soil: no

Hydrologic group: B/D

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A,AC -- 0 to 32 in	loam	moderate	6.06 to 6.70 in	5.6 to 7.3
C -- 32 to 60 in	loam	moderately rapid	4.19 to 5.03 in	5.6 to 7.3

Millington

Extent: 5 percent of the unit

Landform(s): flood plains

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

313--Spillville loam, occasionally flooded

Linder

Extent: 5 percent of the unit

Landform(s): outwash plains

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

Representative soil profile:

Texture

Permeability

*Available water
capacity*

pH

Map Unit Description (MN)

Martin County, Minnesota

336--Delft loam

Delft

Extent: 90 percent of the unit

Landform(s): drainageways on moraines

Slope gradient: 1 to 3 percent

Parent material: fine-loamy colluvium

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .24

Land capability, nonirrigated 2w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 36 in	loam	moderate	6.45 to 7.17 in	5.6 to 7.8
A -- 36 to 44 in	loam	moderate	1.57 to 1.82 in	5.6 to 7.8
Bg -- 44 to 54 in	loam	moderate	1.87 to 2.17 in	6.6 to 7.8
Cg -- 54 to 60 in	loam	moderate	0.89 to 1.12 in	7.4 to 8.4

Clarion

Extent: 5 percent of the unit

Landform(s): moraines

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

336--Delft loam

Terril

Extent: 5 percent of the unit

Landform(s): moraines

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

Representative soil profile:

Texture

Permeability

*Available water
capacity*

pH

Map Unit Description (MN)

Martin County, Minnesota

350--Canisteo clay loam, depressional

Canisteo, depressional

Extent: 90 percent of the unit
Landform(s): depressions on moraines
Slope gradient: 0 to 1 percent
Parent material: fine-loamy till
Restrictive feature(s): greater than 60 inches
Flooding: rare
Ponding: frequent
Drainage class: very poorly drained

Soil loss tolerance (T factor): 5
Wind erodibility group (WEG): 4L
Wind erodibility index (WEI): 86
Kw factor (surface layer) .24
Land capability, nonirrigated 3w
Hydric soil: yes
Hydrologic group: B/D
Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 9 in	clay loam	moderate	1.63 to 1.99 in	7.4 to 8.4
Bg -- 9 to 27 in	clay loam	moderate	2.66 to 3.37 in	7.4 to 8.4
Cg -- 27 to 60 in	loam	moderate	4.63 to 5.29 in	7.4 to 8.4

Glencoe

Extent: 5 percent of the unit
Landform(s): depressions
Slope gradient:
Parent material:
Restrictive feature(s): greater than 60 inches
Flooding:
Ponding:
Drainage class:

Soil loss tolerance (T factor):
Wind erodibility group (WEG):
Wind erodibility index (WEI):
Kw factor (surface layer)
Land capability, nonirrigated
Hydric soil: yes
Hydrologic group:
Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

350--Canisteo clay loam, depressional

Harps

Extent: 5 percent of the unit

Landform(s): rims

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

Representative soil profile:

Texture

Permeability

*Available water
capacity*

pH

Map Unit Description (MN)

Martin County, Minnesota

392--Biscay sandy clay loam

Biscay

Extent: 90 percent of the unit
Landform(s): flats on outwash plains
Slope gradient: 0 to 2 percent
Parent material: fine-loamy outwash over sandy outwash
Restrictive feature(s): greater than 60 inches
Flooding: none
Ponding: none
Drainage class: poorly drained

Soil loss tolerance (T factor): 3
Wind erodibility group (WEG): 4L
Wind erodibility index (WEI): 86
Kw factor (surface layer) .17
Land capability, nonirrigated 2w
Hydric soil: yes
Hydrologic group: B/D
Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 22 in	sandy clay loam	moderate	3.75 to 4.19 in	6.1 to 7.8
Bg -- 22 to 32 in	sandy clay loam	moderate	1.67 to 1.87 in	6.6 to 7.8
Cg -- 32 to 60 in	stratified sand to loamy sand	rapid	0.56 to 1.12 in	7.4 to 8.4

Lemond

Extent: 4 percent of the unit
Landform(s): drainageways
Slope gradient:
Parent material:
Restrictive feature(s): greater than 60 inches
Flooding:
Ponding:
Drainage class:

Soil loss tolerance (T factor):
Wind erodibility group (WEG):
Wind erodibility index (WEI):
Kw factor (surface layer)
Land capability, nonirrigated
Hydric soil: yes
Hydrologic group:
Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

392--Biscay sandy clay loam

Linder

Extent: 3 percent of the unit

Landform(s): outwash plains

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Wadena

Extent: 3 percent of the unit

Landform(s): outwash plains

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

499--Hanska loam, depressional

Hanska, depressional

Extent: 90 percent of the unit

Landform(s): depressions on outwash plains

Slope gradient: 0 to 1 percent

Parent material: coarse-loamy outwash over sandy outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: frequent

Drainage class: very poorly drained

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .32

Land capability, nonirrigated 3w

Hydric soil: yes

Hydrologic group: A/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 24 in	loam	moderately rapid	4.80 to 5.28 in	6.1 to 7.3
Bg -- 24 to 37 in	sandy loam	moderately rapid	1.30 to 1.69 in	6.1 to 7.3
2C -- 37 to 60 in	sand	rapid	1.14 to 1.60 in	6.6 to 7.8

Litchfield

Extent: 5 percent of the unit

Landform(s): outwash plains

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

499--Hanska loam, depressional

Lemond

Extent: 5 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

Representative soil profile:

Texture

Permeability

*Available water
capacity*

pH

Map Unit Description (MN)

Martin County, Minnesota

539--Palms muck

Palms

<p><i>Extent:</i> 90 percent of the unit</p> <p><i>Landform(s):</i> depressions on moraines</p> <p><i>Slope gradient:</i> 0 to 2 percent</p> <p><i>Parent material:</i> muck herbaceous organic material over fine-loamy till</p> <p><i>Restrictive feature(s):</i> greater than 60 inches</p> <p><i>Flooding:</i> none</p> <p><i>Ponding:</i> frequent</p> <p><i>Drainage class:</i> very poorly drained</p>	<p><i>Soil loss tolerance (T factor):</i> 1</p> <p><i>Wind erodibility group (WEG):</i> 2</p> <p><i>Wind erodibility index (WEI):</i> 134</p> <p><i>Kw factor (surface layer)</i> .02</p> <p><i>Land capability, nonirrigated</i> 3w</p> <p><i>Hydric soil:</i> yes</p> <p><i>Hydrologic group:</i> B/D</p> <p><i>Potential for frost action:</i> high</p>
---	--

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Op -- 0 to 11 in	muck	moderately rapid	3.86 to 5.29 in	
Oa -- 11 to 33 in	muck	moderate	4.85 to 5.73 in	
2C -- 33 to 47 in	silty clay loam	moderate	2.48 to 3.03 in	
2Cg -- 47 to 60 in	clay loam	moderate	1.95 to 2.47 in	

Glencoe

<p><i>Extent:</i> 4 percent of the unit</p> <p><i>Landform(s):</i> depressions</p> <p><i>Slope gradient:</i></p> <p><i>Parent material:</i></p> <p><i>Restrictive feature(s):</i> greater than 60 inches</p> <p><i>Flooding:</i></p> <p><i>Ponding:</i></p> <p><i>Drainage class:</i></p>	<p><i>Soil loss tolerance (T factor):</i></p> <p><i>Wind erodibility group (WEG):</i></p> <p><i>Wind erodibility index (WEI):</i></p> <p><i>Kw factor (surface layer)</i></p> <p><i>Land capability, nonirrigated</i></p> <p><i>Hydric soil:</i> yes</p> <p><i>Hydrologic group:</i></p> <p><i>Potential for frost action:</i></p>
---	--

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>

Map Unit Description (MN)

Martin County, Minnesota

539--Palms muck

Canisteo

Extent: 2 percent of the unit

Landform(s): rims

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Okoboji

Extent: 2 percent of the unit

Landform(s): depressions

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

539--Palms muck

Harps

Extent: 2 percent of the unit

Landform(s): rims

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

548--Palms muck, sandy substratum

Palms, sandy substratum

Extent: 85 percent of the unit
Landform(s): depressions on outwash plains
Slope gradient: 0 to 2 percent
Parent material: muck herbaceous organic material over sandy outwash
Restrictive feature(s): greater than 60 inches
Flooding: frequent
Ponding: frequent
Drainage class: very poorly drained

Soil loss tolerance (T factor): 1
Wind erodibility group (WEG): 2
Wind erodibility index (WEI): 134
Kw factor (surface layer) .02

Land capability, nonirrigated 3w
Hydric soil: yes
Hydrologic group: A/D
Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Op -- 0 to 10 in	muck	moderately rapid	3.44 to 4.43 in	
Oa -- 10 to 21 in	muck	moderately rapid	1.43 to 2.20 in	
2A -- 21 to 34 in	silty clay loam	moderately rapid	1.69 to 2.60 in	
3Cg -- 34 to 60 in	stratified loamy sand to sand	rapid	0.78 to 2.60 in	

Linder

Extent: 5 percent of the unit
Landform(s): beach ridges
Slope gradient:
Parent material:
Restrictive feature(s): greater than 60 inches
Flooding:
Ponding:
Drainage class:

Soil loss tolerance (T factor):
Wind erodibility group (WEG):
Wind erodibility index (WEI):
Kw factor (surface layer)
Land capability, nonirrigated
Hydric soil: no
Hydrologic group:
Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>

Map Unit Description (MN)

Martin County, Minnesota

548--Palms muck, sandy substratum

Blue Earth

Extent: 5 percent of the unit

Landform(s): depressions

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>

Lemond

Extent: 5 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>

Map Unit Description (MN)

Martin County, Minnesota

664--Zook silty clay loam, occasionally flooded

Zook, occasionally flooded

Extent: 90 percent of the unit

Landform(s): flood plains

Slope gradient: 0 to 2 percent

Parent material: clayey alluvium

Restrictive feature(s): greater than 60 inches

Flooding: occasional

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4

Wind erodibility index (WEI): 86

Kw factor (surface layer) .28

Land capability, nonirrigated 2w

Hydric soil: yes

Hydrologic group: C/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 24 in	silty clay loam	moderately slow	5.04 to 5.52 in	5.6 to 7.3
Bg -- 24 to 38 in	silty clay	slow	1.52 to 1.79 in	5.6 to 7.8
Cg -- 38 to 60 in	silty clay loam	moderately slow	2.43 to 4.85 in	5.6 to 7.8

Millington

Extent: 5 percent of the unit

Landform(s): flood plains

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

664--Zook silty clay loam, occasionally flooded

Coland

Extent: 5 percent of the unit

Landform(s): flood plains

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

Representative soil profile:

Texture

Permeability

*Available water
capacity*

pH

Map Unit Description (MN)

Martin County, Minnesota

818--Lemond-Linder complex

Lemond

Extent: 55 percent of the unit

Landform(s): flats on outwash plains

Slope gradient: 0 to 2 percent

Parent material: coarse-loamy outwash over sandy outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .17

Land capability, nonirrigated 2w

Hydric soil: yes

Hydrologic group: A/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 24 in	sandy loam	moderately rapid	3.60 to 4.32 in	7.4 to 8.4
Bg -- 24 to 30 in	sandy loam	moderately rapid	0.59 to 0.77 in	7.4 to 8.4
2Cg -- 30 to 60 in	loamy sand	rapid	1.50 to 2.09 in	7.4 to 8.4

Linder

Extent: 25 percent of the unit

Landform(s): rises

Slope gradient: 0 to 2 percent

Parent material: coarse-loamy outwash over sandy outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .20

Land capability, nonirrigated 2s

Hydric soil: no

Hydrologic group: B/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 20 in	sandy loam	moderate	3.01 to 4.02 in	5.6 to 7.8
Bw -- 20 to 30 in	sandy loam	moderately rapid	1.48 to 1.67 in	6.1 to 7.8
2C -- 30 to 60 in	loamy coarse sand	very rapid	0.60 to 1.20 in	7.4 to 8.4

Map Unit Description (MN)

Martin County, Minnesota

818--Lemond-Linder complex

Blue Earth

Extent: 10 percent of the unit

Landform(s): depressions

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Palms

Extent: 5 percent of the unit

Landform(s): depressions

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

818--Lemond-Linder complex

Crippen

Extent: 5 percent of the unit

Landform(s): moraines

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

Representative soil profile:

Texture

Permeability

*Available water
capacity*

pH

Map Unit Description (MN)

Martin County, Minnesota

886--Nicollet-Crippin complex

Nicollet

Extent: 50 percent of the unit
Landform(s): rises on moraines
Slope gradient: 1 to 3 percent
Parent material: fine-loamy till
Restrictive feature(s): greater than 60 inches
Flooding: none
Ponding: none
Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 5
Wind erodibility group (WEG): 6
Wind erodibility index (WEI): 48
Kw factor (surface layer) .24
Land capability, nonirrigated 1
Hydric soil: no
Hydrologic group: B/D
Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 17 in	clay loam	moderate	3.39 to 3.72 in	6.6 to 8.4
Bw -- 17 to 36 in	clay loam	moderate	3.21 to 3.59 in	7.4 to 8.4
C -- 36 to 60 in	loam	moderate	4.08 to 4.56 in	7.9 to 8.4

Crippin

Extent: 35 percent of the unit
Landform(s): rises on moraines
Slope gradient: 0 to 3 percent
Parent material: fine-loamy till
Restrictive feature(s): greater than 60 inches
Flooding: none
Ponding: none
Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 5
Wind erodibility group (WEG): 4L
Wind erodibility index (WEI): 86
Kw factor (surface layer) .24
Land capability, nonirrigated 1
Hydric soil: no
Hydrologic group: B/D
Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 15 in	loam	moderate	2.99 to 3.29 in	6.6 to 8.4
Bw -- 15 to 27 in	clay loam	moderate	2.01 to 2.24 in	7.4 to 8.4
C -- 27 to 60 in	loam	moderate	4.63 to 6.28 in	7.9 to 8.4

Map Unit Description (MN)

Martin County, Minnesota

886--Nicollet-Crippin complex

Webster

Extent: 5 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Clarion

Extent: 5 percent of the unit

Landform(s): moraines

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

886--Nicollet-Crippin complex

Swanlake

Extent: 5 percent of the unit

Landform(s): moraines

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

Representative soil profile:

Texture

Permeability

*Available water
capacity*

pH

Map Unit Description (MN)

Martin County, Minnesota

887B--Clarion-Swanlake loams, 2 to 6 percent slopes

Clarion

Extent: 60 percent of the unit

Landform(s): hills on moraines

Slope gradient: 2 to 6 percent

Parent material: fine-loamy till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 2e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 14 in	loam	moderate	2.83 to 3.12 in	5.6 to 7.3
Bw -- 14 to 33 in	loam	moderate	3.21 to 3.59 in	5.6 to 7.8
C -- 33 to 60 in	loam	moderate	4.55 to 5.09 in	7.4 to 8.4

Swanlake

Extent: 25 percent of the unit

Landform(s): hills on moraines

Slope gradient: 2 to 6 percent

Parent material: fine-loamy till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .28

Land capability, nonirrigated 2e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 21 in	loam	moderate	4.17 to 5.01 in	7.4 to 8.4
Bw -- 21 to 28 in	loam	moderate	1.20 to 1.35 in	7.4 to 8.4
C -- 28 to 60 in	loam	moderate	5.42 to 6.06 in	7.4 to 8.4

Map Unit Description (MN)

Martin County, Minnesota

887B--Clarion-Swanlake loams, 2 to 6 percent slopes

Webster

Extent: 5 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>

Nicollet

Extent: 5 percent of the unit

Landform(s): moraines

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>

Map Unit Description (MN)

Martin County, Minnesota

887B--Clarion-Swanlake loams, 2 to 6 percent slopes

Canisteo

Extent: 5 percent of the unit

Landform(s): rims

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

Representative soil profile:

Texture

Permeability

*Available water
capacity*

pH

Map Unit Description (MN)

Martin County, Minnesota

909C--Truman-Bold silt loams, 6 to 12 percent slopes

Truman

Extent: 45 percent of the unit

Landform(s): hills on lake plains

Slope gradient: 6 to 12 percent

Parent material: fine-silty lacustrine deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .37

Land capability, nonirrigated 3e

Hydric soil: no

Hydrologic group: B

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 18 in	silt loam	moderate	3.62 to 4.17 in	5.6 to 7.3
Bw -- 18 to 30 in	silt loam	moderate	2.13 to 2.48 in	5.6 to 7.8
C -- 30 to 60 in	silt loam	moderate	5.39 to 5.98 in	7.4 to 8.4

Bold

Extent: 35 percent of the unit

Landform(s): hills on lake plains

Slope gradient: 6 to 12 percent

Parent material: coarse-silty lacustrine deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .43

Land capability, nonirrigated 4e

Hydric soil: no

Hydrologic group: B

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 10 in	silt loam	moderate	2.07 to 2.36 in	7.4 to 8.4
C -- 10 to 60 in	silt loam	moderate	10.00 to 12.00 in	7.4 to 8.4

Map Unit Description (MN)

Martin County, Minnesota

909C--Truman-Bold silt loams, 6 to 12 percent slopes

Madelia

Extent: 10 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>

Kingston

Extent: 5 percent of the unit

Landform(s): lake plains

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>

Map Unit Description (MN)

Martin County, Minnesota

909C--Truman-Bold silt loams, 6 to 12 percent slopes

Terril

Extent: 5 percent of the unit

Landform(s): moraines

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

Representative soil profile:

Texture

Permeability

*Available water
capacity*

pH

Map Unit Description (MN)

Martin County, Minnesota

909D2--Bold-Truman silt loams, 12 to 18 percent slopes, eroded

Bold, eroded

Extent: 45 percent of the unit
Landform(s): hills on lake plains
Slope gradient: 12 to 18 percent
Parent material: coarse-silty lacustrine deposits
Restrictive feature(s): greater than 60 inches
Flooding: none
Ponding: none
Drainage class: well drained

Soil loss tolerance (T factor): 5
Wind erodibility group (WEG): 4L
Wind erodibility index (WEI): 86
Kw factor (surface layer) .43
Land capability, nonirrigated 6e
Hydric soil: no
Hydrologic group: B
Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 8 in	silt loam	moderate	1.65 to 1.89 in	7.4 to 8.4
C -- 8 to 60 in	silt loam	moderate	10.39 to 12.47 in	7.4 to 8.4

Truman, eroded

Extent: 40 percent of the unit
Landform(s): hills on lake plains
Slope gradient: 12 to 18 percent
Parent material: fine-silty lacustrine deposits
Restrictive feature(s): greater than 60 inches
Flooding: none
Ponding: none
Drainage class: well drained

Soil loss tolerance (T factor): 5
Wind erodibility group (WEG): 6
Wind erodibility index (WEI): 48
Kw factor (surface layer) .37
Land capability, nonirrigated 4e
Hydric soil: no
Hydrologic group: B
Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 15 in	silt loam	moderate	2.99 to 3.44 in	5.6 to 7.3
Bw -- 15 to 40 in	silt loam	moderate	4.54 to 5.29 in	5.6 to 7.8
C -- 40 to 60 in	silt loam	moderate	3.54 to 3.94 in	7.4 to 8.4

Map Unit Description (MN)

Martin County, Minnesota

909D2--Bold-Truman silt loams, 12 to 18 percent slopes, eroded

Madelia

Extent: 10 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>

Terril

Extent: 5 percent of the unit

Landform(s): moraines

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>

Map Unit Description (MN)

Martin County, Minnesota

920B--Clarion-Estherville complex, 2 to 6 percent slopes

Clarion

Extent: 50 percent of the unit

Landform(s): hills on moraines

Slope gradient: 2 to 6 percent

Parent material: fine-loamy till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 2e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 14 in	loam	moderate	2.83 to 3.12 in	5.6 to 7.3
Bw -- 14 to 33 in	loam	moderate	3.21 to 3.59 in	5.6 to 7.8
C -- 33 to 60 in	loam	moderate	4.55 to 5.09 in	7.4 to 8.4

Estherville

Extent: 25 percent of the unit

Landform(s): hills on moraines

Slope gradient: 2 to 6 percent

Parent material: coarse-loamy outwash over sandy and gravelly outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat excessively drained

Soil loss tolerance (T factor): 2

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .24

Land capability, nonirrigated 3s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 13 in	sandy loam	moderately rapid	1.69 to 2.34 in	5.6 to 7.3
Bw -- 13 to 18 in	sandy loam	moderately rapid	0.67 to 0.92 in	5.6 to 7.3
2C -- 18 to 60 in	gravelly loamy sand	rapid	0.83 to 1.67 in	6.6 to 8.4

Map Unit Description (MN)

Martin County, Minnesota

920B--Clarion-Estherville complex, 2 to 6 percent slopes

Webster

Extent: 10 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Swanlake

Extent: 10 percent of the unit

Landform(s): moraines

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

920B--Clarion-Estherville complex, 2 to 6 percent slopes

Terril

Extent: 5 percent of the unit

Landform(s): moraines

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

920C2--Clarion-Estherville-Storden complex, 6 to 12 percent slopes, eroded

Clarion, eroded

Extent: 30 percent of the unit

Landform(s): hills on moraines

Slope gradient: 6 to 10 percent

Parent material: fine-loamy till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 3e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 10 in	loam	moderate	1.97 to 2.17 in	5.6 to 7.3
Bw -- 10 to 36 in	loam	moderate	4.42 to 4.94 in	5.6 to 7.3
C -- 36 to 60 in	loam	moderate	4.08 to 4.56 in	7.4 to 8.4

Estherville, eroded

Extent: 20 percent of the unit

Landform(s): hills on moraines

Slope gradient: 6 to 12 percent

Parent material: coarse-loamy outwash over sandy and gravelly outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat excessively drained

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .24

Land capability, nonirrigated 4s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 10 in	sandy loam	moderately rapid	1.28 to 1.77 in	5.6 to 7.3
Bw -- 10 to 30 in	sandy loam	moderately rapid	2.61 to 3.61 in	5.6 to 7.3
2C -- 30 to 60 in	coarse sand	rapid	0.60 to 1.20 in	6.6 to 8.4

Map Unit Description (MN)

Martin County, Minnesota

920C2--Clarion-Estherville-Storden complex, 6 to 12 percent slopes, eroded

Storden, eroded

Extent: 20 percent of the unit

Landform(s): hills on moraines

Slope gradient: 6 to 12 percent

Parent material: fine-loamy till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .32

Land capability, nonirrigated 3e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 9 in	loam	moderate	1.81 to 1.99 in	7.4 to 8.4
C -- 9 to 60 in	loam	moderate	7.62 to 9.65 in	7.4 to 8.4

Delft

Extent: 10 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>

Map Unit Description (MN)

Martin County, Minnesota

920C2--Clarion-Estherville-Storden complex, 6 to 12 percent slopes, eroded

Terril

Extent: 10 percent of the unit

Landform(s): moraines

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Nicollet

Extent: 10 percent of the unit

Landform(s): moraines

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

920D2--Clarion-Estherville-Storden complex, 12 to 18 percent slopes, eroded

Clarion, eroded

Extent: 30 percent of the unit

Landform(s): hills on moraines

Slope gradient: 12 to 18 percent

Parent material: fine-loamy till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 4e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 9 in	loam	moderate	1.81 to 1.99 in	5.6 to 7.3
Bw -- 9 to 18 in	loam	moderate	1.54 to 1.72 in	5.6 to 7.3
C -- 18 to 60 in	loam	moderate	7.09 to 7.93 in	7.4 to 8.4

Estherville, eroded

Extent: 25 percent of the unit

Landform(s): hills on moraines

Slope gradient: 12 to 18 percent

Parent material: coarse-loamy outwash over sandy and gravelly outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat excessively drained

Soil loss tolerance (T factor): 2

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .24

Land capability, nonirrigated 6e

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 9 in	sandy loam	moderately rapid	1.18 to 1.63 in	5.6 to 7.3
Bw -- 9 to 16 in	sandy loam	moderately rapid	0.92 to 1.28 in	5.6 to 7.3
2C -- 16 to 60 in	gravelly coarse sand	rapid	0.87 to 1.75 in	6.6 to 8.4

Map Unit Description (MN)

Martin County, Minnesota

920D2--Clarion-Estherville-Storden complex, 12 to 18 percent slopes, eroded

Storden, eroded

Extent: 20 percent of the unit

Landform(s): hills on moraines

Slope gradient: 12 to 18 percent

Parent material: fine-loamy till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .32

Land capability, nonirrigated 4e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 9 in	loam	moderate	1.81 to 1.99 in	7.4 to 8.4
C -- 9 to 60 in	loam	moderate	7.62 to 9.65 in	7.4 to 8.4

Terril

Extent: 15 percent of the unit

Landform(s): moraines

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

920D2--Clarion-Estherville-Storden complex, 12 to 18 percent slopes, eroded

Delft

Extent: 10 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

Representative soil profile:

Texture

Permeability

*Available water
capacity*

pH

Map Unit Description (MN)

Martin County, Minnesota

921B--Clarion-Storden loams, 2 to 6 percent slopes

Clarion

Extent: 45 percent of the unit

Landform(s): hills on moraines

Slope gradient: 2 to 6 percent

Parent material: fine-loamy till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 2e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 9 in	loam	moderate	1.81 to 1.99 in	5.6 to 7.3
Bw -- 9 to 19 in	loam	moderate	1.67 to 1.87 in	5.6 to 7.8
C -- 19 to 60 in	loam	moderate	6.96 to 7.78 in	7.4 to 8.4

Storden

Extent: 35 percent of the unit

Landform(s): hills on moraines

Slope gradient: 4 to 6 percent

Parent material: fine-loamy till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .32

Land capability, nonirrigated 2e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 9 in	loam	moderate	1.81 to 1.99 in	7.4 to 8.4
C -- 9 to 60 in	loam	moderate	7.62 to 9.65 in	7.4 to 8.4

Map Unit Description (MN)

Martin County, Minnesota

921B--Clarion-Storden loams, 2 to 6 percent slopes

Webster

Extent: 10 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Nicollet

Extent: 10 percent of the unit

Landform(s): moraines

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

921C2--Clarion-Storden loams, 6 to 12 percent slopes, eroded

Clarion, eroded

Extent: 35 percent of the unit

Landform(s): hills on moraines

Slope gradient: 6 to 12 percent

Parent material: fine-loamy till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 3e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 10 in	loam	moderate	1.97 to 2.17 in	5.6 to 7.3
Bw -- 10 to 23 in	loam	moderate	2.21 to 2.47 in	5.6 to 7.3
C -- 23 to 60 in	loam	moderate	6.29 to 7.03 in	7.4 to 8.4

Storden, eroded

Extent: 30 percent of the unit

Landform(s): hills on moraines

Slope gradient: 6 to 12 percent

Parent material: fine-loamy till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .32

Land capability, nonirrigated 3e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 10 in	loam	moderate	1.97 to 2.17 in	7.4 to 8.4
C -- 10 to 60 in	loam	moderate	7.50 to 9.50 in	7.4 to 8.4

Map Unit Description (MN)

Martin County, Minnesota

921C2--Clarion-Storden loams, 6 to 12 percent slopes, eroded

Terril

Extent: 15 percent of the unit

Landform(s): moraines

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Delft

Extent: 10 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

921C2--Clarion-Storden loams, 6 to 12 percent slopes, eroded

Nicollet

Extent: 10 percent of the unit

Landform(s): moraines

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

Representative soil profile:

Texture

Permeability

*Available water
capacity*

pH

Map Unit Description (MN)

Martin County, Minnesota

956--Canisteo-Glencoe clay loams

Canisteo

Extent: 60 percent of the unit

Landform(s): rims on depressions on moraines, flats on moraines

Slope gradient: 0 to 2 percent

Parent material: fine-loamy till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .24

Land capability, nonirrigated 2w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 11 in	clay loam	moderate	1.98 to 2.43 in	7.4 to 8.4
A -- 11 to 22 in	clay loam	moderate	1.65 to 2.09 in	7.4 to 8.4
Bg -- 22 to 28 in	clay loam	moderate	0.71 to 1.06 in	7.4 to 8.4
C -- 28 to 60 in	loam	moderate	4.46 to 5.10 in	7.4 to 8.4

Glencoe

Extent: 30 percent of the unit

Landform(s): depressions on moraines

Slope gradient: 0 to 1 percent

Parent material: fine-loamy alluvium

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: frequent

Drainage class: very poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .24

Land capability, nonirrigated 3w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 26 in	clay loam	moderate	4.68 to 5.72 in	6.1 to 7.8
Bg -- 26 to 38 in	clay loam	moderate	2.20 to 2.69 in	6.1 to 7.8
Cg -- 38 to 60 in	loam	moderate	3.25 to 4.11 in	6.6 to 7.8

Map Unit Description (MN)

Martin County, Minnesota

956--Canisteo-Glencoe clay loams

Webster

Extent: 4 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>

Crippen

Extent: 3 percent of the unit

Landform(s): moraines

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>

Map Unit Description (MN)

Martin County, Minnesota

956--Canisteo-Glencoe clay loams

Nicollet

Extent: 3 percent of the unit

Landform(s): moraines

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

Representative soil profile:

Texture

Permeability

*Available water
capacity*

pH

Map Unit Description (MN)

Martin County, Minnesota

960D2--Storden-Clarion loams, 12 to 18 percent slopes, eroded

Storden, eroded

Extent: 40 percent of the unit

Landform(s): hills on moraines

Slope gradient: 12 to 18 percent

Parent material: fine-loamy till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .32

Land capability, nonirrigated 4e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 8 in	loam	moderate	1.57 to 1.73 in	7.4 to 8.4
C -- 8 to 60 in	loam	moderate	7.80 to 9.87 in	7.4 to 8.4

Clarion, eroded

Extent: 30 percent of the unit

Landform(s): hills on moraines

Slope gradient: 12 to 18 percent

Parent material: fine-loamy till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 4e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 8 in	loam	moderate	1.57 to 1.73 in	5.6 to 7.3
Bw -- 8 to 22 in	loam	moderate	2.41 to 2.69 in	5.6 to 7.3
C -- 22 to 60 in	loam	moderate	6.43 to 7.18 in	7.4 to 8.4

Map Unit Description (MN)

Martin County, Minnesota

960D2--Storden-Clarion loams, 12 to 18 percent slopes, eroded

Delft

Extent: 15 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Terril

Extent: 15 percent of the unit

Landform(s): moraines

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

960E--Storden-Clarion loams, 18 to 25 percent slopes

Storden

Extent: 45 percent of the unit

Landform(s): hills on moraines

Slope gradient: 18 to 25 percent

Parent material: fine-loamy till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .32

Land capability, nonirrigated 6e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 9 in	loam	moderate	1.81 to 1.99 in	7.4 to 8.4
C -- 9 to 60 in	loam	moderate	7.62 to 9.65 in	7.4 to 8.4

Clarion

Extent: 30 percent of the unit

Landform(s): hills on moraines

Slope gradient: 18 to 25 percent

Parent material: fine-loamy till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 6e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 15 in	loam	moderate	2.99 to 3.29 in	5.6 to 7.3
Bw -- 15 to 30 in	loam	moderate	2.54 to 2.84 in	5.6 to 7.3
C -- 30 to 60 in	loam	moderate	5.09 to 5.69 in	7.4 to 8.4

Map Unit Description (MN)

Martin County, Minnesota

960E--Storden-Clarion loams, 18 to 25 percent slopes

Terril

Extent: 15 percent of the unit

Landform(s): moraines

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Delft

Extent: 10 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

1029--Pits, gravel

Pits, gravel

Extent: 100 percent of the unit

Landform(s): outwash plains

Slope gradient:

Parent material: sandy and gravelly outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: unranked

Hydrologic group:

Potential for frost action:

Representative soil profile:

Texture

Permeability

*Available water
capacity*

pH

Map Unit Description (MN)

Martin County, Minnesota

1052--Palms and Okoboji soils, ponded

Palms, ponded

Extent: 50 percent of the unit

Landform(s): depressions on moraines

Slope gradient: 0 to 1 percent

Parent material: muck herbaceous organic material over fine-loamy till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: frequent

Drainage class: very poorly drained

Soil loss tolerance (T factor): 1

Wind erodibility group (WEG): 8

Wind erodibility index (WEI): 0

Kw factor (surface layer) .02

Land capability, nonirrigated 8w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Oa -- 0 to 26 in	muck	moderately rapid	9.09 to 11.69 in	
A -- 26 to 60 in	clay loam	moderate	4.74 to 7.45 in	

Okoboji, ponded

Extent: 50 percent of the unit

Landform(s): depressions on moraines

Slope gradient: 0 to 1 percent

Parent material: fine-silty alluvium

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: frequent

Drainage class: very poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 8

Wind erodibility index (WEI): 0

Kw factor (surface layer) .28

Land capability, nonirrigated 8w

Hydric soil: yes

Hydrologic group: C/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 35 in	silty clay loam	moderately slow	6.31 to 7.01 in	6.6 to 7.8
Bg -- 35 to 46 in	silty clay loam	moderately slow	1.98 to 2.20 in	6.6 to 7.8
Cg -- 46 to 60 in	clay loam	moderately slow	2.48 to 2.76 in	6.6 to 7.8

Map Unit Description (MN)

Martin County, Minnesota

1090--Blue Earth silt loam

Blue Earth

Extent: 85 percent of the unit

Landform(s): flats on moraines

Slope gradient: 0 to 2 percent

Parent material: coprogenic material

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .32

Land capability, nonirrigated 2w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 9 in	silt loam	moderate	1.63 to 1.99 in	7.4 to 8.4
C -- 9 to 60 in	silt loam	moderate	7.11 to 10.16 in	7.4 to 8.4

Glencoe

Extent: 10 percent of the unit

Landform(s): depressions

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

1090--Blue Earth silt loam

Harps

Extent: 5 percent of the unit

Landform(s): rims

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

1356--Water, miscellaneous

Water, miscellaneous

Extent: 100 percent of the unit

Landform(s):

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil:

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

1833--Coland loam, occasionally flooded

Coland, occasionally flooded

Extent: 90 percent of the unit

Landform(s): flood plains

Slope gradient: 0 to 2 percent

Parent material: fine-loamy alluvium

Restrictive feature(s): greater than 60 inches

Flooding: occasional

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 2w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 8 in	loam	moderate	1.57 to 1.73 in	6.1 to 7.3
A -- 8 to 60 in	clay loam	moderate	10.39 to 11.43 in	6.1 to 7.3

Millington

Extent: 4 percent of the unit

Landform(s): flood plains

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>

Map Unit Description (MN)

Martin County, Minnesota

1833--Coland loam, occasionally flooded

Zook

Extent: 3 percent of the unit

Landform(s): flood plains

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

Representative soil profile:

Texture

Permeability

*Available water
capacity*

pH

Linder

Extent: 3 percent of the unit

Landform(s): outwash plains

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

Representative soil profile:

Texture

Permeability

*Available water
capacity*

pH

Map Unit Description (MN)

Martin County, Minnesota

1834--Coland loam, frequently flooded

Coland, frequently flooded

Extent: 90 percent of the unit

Landform(s): flood plains

Slope gradient: 0 to 2 percent

Parent material: fine-loamy alluvium

Restrictive feature(s): greater than 60 inches

Flooding: frequent

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 5w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 16 in	loam	moderate	3.23 to 3.55 in	6.1 to 7.3
AC -- 16 to 44 in	clay loam	moderate	5.59 to 6.15 in	6.1 to 7.3
Cg -- 44 to 60 in	clay loam	moderately rapid	2.05 to 2.68 in	6.1 to 7.8

Millington

Extent: 5 percent of the unit

Landform(s): flood plains

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

1834--Coland loam, frequently flooded

Zook

Extent: 5 percent of the unit

Landform(s): flood plains

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

Representative soil profile:

Texture

Permeability

*Available water
capacity*

pH

Map Unit Description (MN)

Martin County, Minnesota

1852F--Terril-Swanlake loams, 18 to 40 percent slopes

Terril

Extent: 45 percent of the unit

Landform(s): hills on moraines

Slope gradient: 18 to 40 percent

Parent material: fine-loamy colluvium

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 7e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 15 in	loam	moderate	2.99 to 3.29 in	6.1 to 7.3
AB -- 15 to 40 in	loam	moderate	4.28 to 4.79 in	6.1 to 7.3
Bw -- 40 to 60 in	loam	moderate	3.15 to 3.54 in	6.1 to 7.8

Swanlake

Extent: 25 percent of the unit

Landform(s): hills on moraines

Slope gradient: 18 to 40 percent

Parent material: fine-loamy till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .28

Land capability, nonirrigated 6e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 14 in	loam	moderate	2.83 to 3.40 in	7.4 to 8.4
Bw -- 14 to 30 in	loam	moderate	2.68 to 2.99 in	7.4 to 8.4
C -- 30 to 60 in	loam	moderate	5.09 to 5.69 in	7.4 to 8.4

Map Unit Description (MN)

Martin County, Minnesota

1852F--Terril-Swanlake loams, 18 to 40 percent slopes

Delft

Extent: 15 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Clarion

Extent: 15 percent of the unit

Landform(s): moraines

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

1877--Fostoria clay loam

Fostoria

Extent: 90 percent of the unit

Landform(s): moraines

Slope gradient: 0 to 2 percent

Parent material: loamy till over silty glaciolacustrine deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .24

Land capability, nonirrigated 1

Hydric soil: no

Hydrologic group: B/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 19 in	clay loam	moderate	3.78 to 4.16 in	6.1 to 7.3
Bw -- 19 to 25 in	clay loam	moderate	1.26 to 1.39 in	6.1 to 7.3
2C -- 25 to 60 in	silt loam	moderate	6.93 to 7.62 in	7.4 to 8.4

Waldorf

Extent: 4 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

1877--Fostoria clay loam

Ocheyedan

Extent: 3 percent of the unit

Landform(s): lake plains

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Kingston

Extent: 3 percent of the unit

Landform(s): lake plains

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

1907--Lakefield silt loam

Lakefield

Extent: 90 percent of the unit

Landform(s): lake plains

Slope gradient: 0 to 3 percent

Parent material: fine-silty lacustrine deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .37

Land capability, nonirrigated 1

Hydric soil: no

Hydrologic group: C

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 20 in	silt loam	moderate	3.61 to 4.82 in	7.4 to 8.4
Bw -- 20 to 60 in	silt loam	moderate	6.36 to 7.95 in	7.4 to 8.4

Madelia

Extent: 5 percent of the unit

Landform(s): drainageways

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>

Map Unit Description (MN)

Martin County, Minnesota

1907--Lakefield silt loam

Truman

Extent: 5 percent of the unit

Landform(s): lake plains

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: no

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

L13A--Klossner muck, depressional, 0 to 1 percent slopes

Klossner, drained

Extent: 65 to 85 percent of the unit

Landform(s): depressions on moraines

Slope gradient: 0 to 1 percent

Parent material: herbaceous organic material over loamy glaciofluvial deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: frequent

Drainage class: very poorly drained

Soil loss tolerance (T factor): 1

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .02

Land capability, nonirrigated 3w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Op -- 0 to 10 in	muck	moderately rapid	3.44 to 4.72 in	
Oa -- 10 to 26 in	muck	moderately rapid	5.65 to 7.75 in	
2A1 -- 26 to 36 in	mucky silty clay loam	moderate	2.17 to 2.56 in	
2A2 -- 36 to 48 in	silty clay loam	moderate	2.20 to 2.69 in	
2Cg -- 48 to 80 in	loam	moderate	4.78 to 6.06 in	

Map Unit Description (MN)

Martin County, Minnesota

L83A--Webster clay loam, 0 to 2 percent slopes

Webster

Extent: 50 to 85 percent of the unit

Landform(s): flats on moraines, swales on moraines

Slope gradient: 0 to 2 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .24

Land capability, nonirrigated 2w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 19 in	clay loam	moderate	3.59 to 3.97 in	6.6 to 7.3
Bg -- 19 to 26 in	clay loam	moderate	1.13 to 1.28 in	6.6 to 7.8
BCg,Cg -- 26 to 60 in	loam	moderate	5.08 to 6.43 in	7.4 to 8.4

L84A--Glencoe clay loam, depressional, 0 to 1 percent slopes

Glencoe, depressional

Extent: 75 to 100 percent of the unit

Landform(s): depressions on moraines

Slope gradient: 0 to 1 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: frequent

Drainage class: very poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .24

Land capability, nonirrigated 3w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 24 in	clay loam	moderate	4.32 to 5.28 in	6.1 to 7.8
ABg -- 24 to 35 in	clay loam	moderate	1.98 to 2.43 in	6.1 to 7.8
Bg -- 35 to 48 in	loam	moderate	1.95 to 2.47 in	6.6 to 7.8
Cg -- 48 to 60 in	loam	moderate	1.77 to 2.24 in	7.4 to 8.4

Map Unit Description (MN)

Martin County, Minnesota

L85A--Nicollet clay loam, 1 to 3 percent slopes

Nicollet

Extent: 70 to 95 percent of the unit

Landform(s): flats on moraines, rises on moraines

Slope gradient: 1 to 3 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .24

Land capability, nonirrigated 1

Hydric soil: no

Hydrologic group: B/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 17 in	clay loam	moderate	2.88 to 3.72 in	5.6 to 7.3
Bw,Bg -- 17 to 33 in	clay loam	moderate	2.42 to 3.07 in	5.6 to 7.3
Bg -- 33 to 36 in	clay loam	moderate	0.41 to 0.52 in	7.4 to 8.4
Cg -- 36 to 60 in	loam	moderate	3.60 to 4.56 in	7.4 to 8.4

Map Unit Description (MN)

Martin County, Minnesota

L107A--Canisteo-Glencoe, depressional complex, 0 to 2 percent slopes

Canisteo

Extent: 30 to 70 percent of the unit

Landform(s): rims on moraines

Slope gradient: 0 to 2 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .24

Land capability, nonirrigated 2w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 18 in	clay loam	moderate	3.26 to 3.98 in	7.4 to 8.4
Bkg -- 18 to 39 in	loam	moderate	2.50 to 3.76 in	7.4 to 8.4
Cg -- 39 to 80 in	loam	moderate	6.14 to 7.78 in	7.4 to 8.4

Glencoe, depressional

Extent: 15 to 55 percent of the unit

Landform(s): depressions on moraines

Slope gradient: 0 to 1 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: frequent

Drainage class: very poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .24

Land capability, nonirrigated 3w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 10 in	clay loam	moderate	1.77 to 2.17 in	6.1 to 7.8
A,ABg -- 10 to 35 in	clay loam	moderate	4.54 to 5.54 in	6.1 to 7.8
Bg -- 35 to 48 in	loam	moderate	1.95 to 2.47 in	6.6 to 7.8
Cg -- 48 to 60 in	loam	moderate	1.77 to 2.24 in	7.4 to 8.4

Map Unit Description (MN)

Martin County, Minnesota

L163A--Okoboji silty clay loam, depressional, 0 to 1 percent slopes

Okoboji, depressional

Extent: 70 to 95 percent of the unit

Landform(s): depressions on lake plains, depressions on moraines

Slope gradient: 0 to 1 percent

Parent material: lacustrine sediments over till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: frequent

Drainage class: very poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .28

Land capability, nonirrigated 3w

Hydric soil: yes

Hydrologic group: C/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 26 in	silty clay loam	moderately slow	5.46 to 5.98 in	6.1 to 7.8
Bg -- 26 to 42 in	silty clay	moderately slow	2.91 to 3.23 in	6.6 to 7.8
Cg -- 42 to 60 in	silty clay loam	moderately slow	3.19 to 3.54 in	6.6 to 8.4

W--Water

Water

Extent: 100 percent of the unit

Landform(s):

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: unranked

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Martin County, Minnesota

This report provides a semitabular listing of some soil and site properties and interpretations that are valuable in communicating the concept of a map unit. The report also provides easy access to the commonly used conservation planning information in one place. The major soil components in each map unit are displayed. Minor components may be displayed if they are included in the database and are selected at the time the report is generated.