

Map Unit Description (MN)

Redwood County, Minnesota

[Data apply to the entire extent of the map unit within the survey area. Map unit and soil properties for a specific parcel of land may vary somewhat and should be determined by onsite investigation]

27A--Dickinson fine sandy loam, 0 to 2 percent slopes

Dickinson

Extent: 90 percent of the unit

Landform(s): outwash plains, terraces

Slope gradient: 0 to 2 percent

Parent material: outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .20

Land capability, nonirrigated 3s

Hydric soil: no

Hydrologic group: A

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|-----------------|---------------------|---------------------------------|------------|
| Ap,A -- 0 to 17 in | fine sandy loam | moderately rapid | 2.71 to 3.05 in | 6.1 to 7.3 |
| Bw -- 17 to 36 in | fine sandy loam | moderately rapid | 2.83 to 3.21 in | 6.1 to 7.3 |
| C -- 36 to 60 in | sand | rapid | 1.20 to 1.68 in | 6.1 to 7.3 |

27B--Dickinson fine sandy loam, 2 to 6 percent slopes

Dickinson

Extent: 90 percent of the unit

Landform(s): outwash plains, terraces

Slope gradient: 2 to 6 percent

Parent material: outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .20

Land capability, nonirrigated 3e

Hydric soil: no

Hydrologic group: A

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|-----------------|---------------------|---------------------------------|------------|
| Ap,A -- 0 to 17 in | fine sandy loam | moderately rapid | 2.71 to 3.05 in | 6.1 to 7.3 |
| Bw -- 17 to 36 in | fine sandy loam | moderately rapid | 2.83 to 3.21 in | 6.1 to 7.3 |
| C -- 36 to 60 in | sand | rapid | 1.20 to 1.68 in | 6.1 to 7.3 |

Map Unit Description (MN)

Redwood County, Minnesota

31E--Storden loam, 18 to 25 percent slopes

Storden

Extent: 85 percent of the unit

Landform(s): moraines

Slope gradient: 18 to 25 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .32

Land capability, nonirrigated 6e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| A -- 0 to 10 in | loam | moderate | 1.97 to 2.17 in | 7.4 to 8.4 |
| C1 -- 10 to 37 in | loam | moderate | 4.62 to 5.16 in | 7.4 to 8.4 |
| C2 -- 37 to 60 in | loam | moderate | 3.88 to 4.34 in | 7.4 to 8.4 |

31F--Storden loam, 25 to 40 percent slopes

Storden

Extent: 86 percent of the unit

Landform(s): moraines

Slope gradient: 25 to 40 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .32

Land capability, nonirrigated 7e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| A -- 0 to 8 in | loam | moderate | 1.57 to 1.73 in | 7.4 to 8.4 |
| C1 -- 8 to 37 in | loam | moderate | 4.95 to 5.54 in | 7.4 to 8.4 |
| C2 -- 37 to 60 in | loam | moderate | 3.88 to 4.34 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

39A--Wadena loam, 0 to 2 percent slopes

Wadena

Extent: 90 percent of the unit

Landform(s): outwash plains, terraces

Slope gradient: 0 to 2 percent

Parent material: outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .20

Land capability, nonirrigated 2s

Hydric soil: no

Hydrologic group: B

Potential for frost action: low

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|-----------------------------------------|---------------------|---------------------------------|------------|
| Ap,A -- 0 to 13 in | loam | moderate | 2.60 to 2.86 in | 6.1 to 7.3 |
| Bw -- 13 to 29 in | loam | moderate | 2.74 to 3.07 in | 6.1 to 7.3 |
| 2C -- 29 to 60 in | stratified gravelly coarse sand to sand | rapid | 0.61 to 1.23 in | 6.6 to 8.4 |

39B--Wadena loam, 2 to 6 percent slopes

Wadena

Extent: 90 percent of the unit

Landform(s): outwash plains, terraces

Slope gradient: 2 to 6 percent

Parent material: outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .20

Land capability, nonirrigated 2e

Hydric soil: no

Hydrologic group: B

Potential for frost action: low

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|-----------------------------------------|---------------------|---------------------------------|------------|
| Ap,A -- 0 to 13 in | loam | moderate | 2.60 to 2.86 in | 6.1 to 7.3 |
| Bw -- 13 to 29 in | loam | moderate | 2.74 to 3.07 in | 6.1 to 7.3 |
| 2C -- 29 to 60 in | stratified gravelly coarse sand to sand | rapid | 0.61 to 1.23 in | 6.6 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

41A--Estherville sandy loam, 0 to 2 percent slopes

Estherville

Extent: 90 percent of the unit
Landform(s): outwash plains, terraces
Slope gradient: 0 to 2 percent
Parent material: outwash
Restrictive feature(s): greater than 60 inches
Flooding: none
Ponding: none
Drainage class: somewhat excessively drained

Soil loss tolerance (T factor): 2
Wind erodibility group (WEG): 3
Wind erodibility index (WEI): 86
Kw factor (surface layer) .20
Land capability, nonirrigated 3s
Hydric soil: no
Hydrologic group: A
Potential for frost action: low

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------------|---------------------|---------------------------------|------------|
| Ap,A -- 0 to 14 in | sandy loam | moderately rapid | 1.79 to 2.07 in | 6.1 to 7.3 |
| Bw -- 14 to 19 in | sandy loam | moderately rapid | 0.61 to 0.72 in | 6.1 to 7.3 |
| 2C -- 19 to 60 in | gravelly coarse sand | rapid | 0.82 to 1.64 in | 7.4 to 8.4 |

41B--Estherville sandy loam, 2 to 6 percent slopes

Estherville

Extent: 90 percent of the unit
Landform(s): outwash plains, terraces
Slope gradient: 2 to 6 percent
Parent material: outwash
Restrictive feature(s): greater than 60 inches
Flooding: none
Ponding: none
Drainage class: somewhat excessively drained

Soil loss tolerance (T factor): 2
Wind erodibility group (WEG): 3
Wind erodibility index (WEI): 86
Kw factor (surface layer) .20
Land capability, nonirrigated 3s
Hydric soil: no
Hydrologic group: A
Potential for frost action: low

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------------|---------------------|---------------------------------|------------|
| Ap,A -- 0 to 14 in | sandy loam | moderately rapid | 1.79 to 2.07 in | 6.1 to 7.3 |
| Bw -- 14 to 19 in | sandy loam | moderately rapid | 0.61 to 0.72 in | 6.1 to 7.3 |
| 2C -- 19 to 60 in | gravelly coarse sand | rapid | 0.82 to 1.64 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

42C--Salida gravelly sandy loam, 2 to 12 percent slopes

Salida

Extent: 90 percent of the unit

Landform(s): outwash plains, terraces

Slope gradient: 2 to 12 percent

Parent material: outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: excessively drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .15

Land capability, nonirrigated 4s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|---------------------|---------------------|---------------------------------|------------|
| A -- 0 to 8 in | gravelly sandy loam | moderately rapid | 1.02 to 1.18 in | 6.1 to 7.3 |
| C -- 8 to 60 in | very gravelly sand | very rapid | 2.60 to 3.64 in | 7.4 to 8.4 |

42E--Salida gravelly sandy loam, 12 to 35 percent slopes

Salida

Extent: 90 percent of the unit

Landform(s): outwash plains, terraces

Slope gradient: 12 to 35 percent

Parent material: outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: excessively drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .15

Land capability, nonirrigated 7s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|---------------------|---------------------|---------------------------------|------------|
| A -- 0 to 8 in | gravelly sandy loam | moderately rapid | 1.02 to 1.18 in | 6.1 to 7.3 |
| C -- 8 to 60 in | very gravelly sand | very rapid | 2.60 to 3.64 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

86--Canisteo clay loam

Canisteo

Extent: 85 percent of the unit

Landform(s): flats

Slope gradient: 0 to 2 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .24

Land capability, nonirrigated 2w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap,A -- 0 to 18 in | clay loam | moderate | 3.08 to 3.44 in | 7.4 to 8.4 |
| AB,Bg -- 18 to 29 in | clay loam | moderate | 1.65 to 2.09 in | 7.4 to 8.4 |
| C -- 29 to 60 in | loam | moderate | 5.22 to 5.83 in | 7.4 to 8.4 |

94B--Terril loam, 2 to 6 percent slopes

Terril

Extent: 85 percent of the unit

Landform(s): moraines

Slope gradient: 2 to 6 percent

Parent material: colluvium

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 2e

Hydric soil: no

Hydrologic group: C

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap,A1 -- 0 to 24 in | loam | moderate | 4.80 to 5.28 in | 6.1 to 7.3 |
| A2 -- 24 to 34 in | loam | moderate | 1.97 to 2.17 in | 6.1 to 7.3 |
| Bw -- 34 to 60 in | loam | moderate | 4.42 to 4.94 in | 6.1 to 7.3 |

Map Unit Description (MN)

Redwood County, Minnesota

94C--Terril loam, 6 to 12 percent slopes

Terril

Extent: 85 percent of the unit

Landform(s): moraines

Slope gradient: 6 to 12 percent

Parent material: colluvium

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 3e

Hydric soil: no

Hydrologic group: C

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap,A1 -- 0 to 24 in | loam | moderate | 4.80 to 5.28 in | 6.1 to 7.3 |
| A2 -- 24 to 34 in | loam | moderate | 1.97 to 2.17 in | 6.1 to 7.3 |
| Bw -- 34 to 60 in | loam | moderate | 4.42 to 4.94 in | 6.1 to 7.3 |

128A--Grogan loam, 0 to 2 percent slopes

Grogan

Extent: 85 percent of the unit

Landform(s): moraines

Slope gradient: 0 to 2 percent

Parent material: lacustrine deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .37

Land capability, nonirrigated 1

Hydric soil: no

Hydrologic group: A

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------------------------------------|---------------------|---------------------------------|------------|
| Ap,AB -- 0 to 15 in | loam | moderately rapid | 2.99 to 3.29 in | 6.1 to 7.3 |
| Bw,BC -- 15 to 34 in | very fine sandy loam | moderately rapid | 3.21 to 3.59 in | 6.1 to 7.3 |
| C -- 34 to 60 in | stratified loamy very fine sand to silt loam | moderately rapid | 4.42 to 4.94 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

128B--Grogan loam, 2 to 6 percent slopes

Grogan

Extent: 85 percent of the unit

Landform(s): moraines

Slope gradient: 2 to 6 percent

Parent material: lacustrine deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .37

Land capability, nonirrigated 2e

Hydric soil: no

Hydrologic group: A

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------------------------------------|---------------------|---------------------------------|------------|
| Ap,AB -- 0 to 15 in | loam | moderately rapid | 2.99 to 3.29 in | 6.1 to 7.3 |
| Bw,Bc -- 15 to 34 in | very fine sandy loam | moderately rapid | 3.21 to 3.59 in | 6.1 to 7.3 |
| C -- 34 to 60 in | stratified loamy very fine sand to silt loam | moderately rapid | 4.42 to 4.94 in | 7.4 to 8.4 |

227--Lemond loam

Lemond

Extent: 85 percent of the unit

Landform(s): flats

Slope gradient: 0 to 2 percent

Parent material: outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .32

Land capability, nonirrigated 2w

Hydric soil: yes

Hydrologic group: A/D

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap -- 0 to 12 in | loam | moderately rapid | 2.36 to 2.60 in | 7.4 to 8.4 |
| Bg -- 12 to 32 in | sandy loam | moderately rapid | 2.41 to 2.81 in | 7.4 to 8.4 |
| 2Cg -- 32 to 60 in | loamy sand | rapid | 2.24 to 2.80 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

247--Linder loam

Linder

Extent: 85 percent of the unit

Landform(s): moraines

Slope gradient: 0 to 2 percent

Parent material: outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .32

Land capability, nonirrigated 2s

Hydric soil: no

Hydrologic group: B/D

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------------------|---------------------|---------------------------------|------------|
| Ap,A -- 0 to 18 in | loam | moderate | 3.62 to 3.98 in | 6.1 to 7.3 |
| Bw,Bg -- 18 to 28 in | sandy loam | moderately rapid | 1.18 to 1.38 in | 6.1 to 7.3 |
| 2Cg -- 28 to 60 in | gravelly loamy coarse sand | very rapid | 0.64 to 1.28 in | 7.4 to 8.4 |

255--Mayer loam

Mayer

Extent: 85 percent of the unit

Landform(s): flats

Slope gradient: 0 to 2 percent

Parent material: outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .28

Land capability, nonirrigated 2w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------------|---------------------|---------------------------------|------------|
| Ap,A -- 0 to 22 in | loam | moderate | 4.41 to 4.85 in | 7.4 to 8.4 |
| Bg -- 22 to 32 in | silt loam | moderate | 1.97 to 2.17 in | 7.4 to 8.4 |
| 2Cg -- 32 to 60 in | gravelly coarse sand | rapid | 0.56 to 1.12 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

282--Hanska fine sandy loam

Hanska

Extent: 85 percent of the unit

Landform(s): flats

Slope gradient: 0 to 2 percent

Parent material: outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .20

Land capability, nonirrigated 2w

Hydric soil: yes

Hydrologic group: A/D

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|-----------------|---------------------|---------------------------------|------------|
| Ap,A -- 0 to 18 in | fine sandy loam | moderately rapid | 2.90 to 3.26 in | 6.1 to 7.3 |
| Bg -- 18 to 38 in | sandy loam | moderately rapid | 2.36 to 2.76 in | 6.1 to 7.3 |
| 2Cg -- 38 to 60 in | sand | rapid | 1.10 to 1.54 in | 6.6 to 7.8 |

313--Spillville loam, occasionally flooded

Spillville, occasionally flooded

Extent: 85 percent of the unit

Landform(s): flood plains

Slope gradient: 0 to 1 percent

Parent material: alluvium

Restrictive feature(s): greater than 60 inches

Flooding: occasional

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 2w

Hydric soil: no

Hydrologic group: C

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap,A -- 0 to 48 in | loam | moderate | 9.61 to 10.57 in | 6.1 to 7.3 |
| C -- 48 to 60 in | loam | moderately rapid | 2.01 to 2.24 in | 6.1 to 7.3 |

Map Unit Description (MN)

Redwood County, Minnesota

317--Oshawa silty clay loam

Oshawa, frequently flooded

Extent: 85 percent of the unit

Landform(s): flood plains

Slope gradient: 0 to 1 percent

Parent material: alluvium

Restrictive feature(s): greater than 60 inches

Flooding: frequent

Ponding: none

Drainage class: very poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .32

Land capability, nonirrigated 6w

Hydric soil: yes

Hydrologic group: C/D

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|-----------------|---------------------|---------------------------------|------------|
| A -- 0 to 39 in | silty clay loam | moderately slow | 7.02 to 8.57 in | 7.4 to 7.8 |
| Cg -- 39 to 60 in | silty clay loam | moderately slow | 3.34 to 3.96 in | 7.4 to 7.8 |

321--Tilfer clay loam

Tilfer, occasionally flooded

Extent: 86 percent of the unit

Landform(s): flats on benches

Slope gradient: 0 to 2 percent

Parent material: alluvium

Restrictive feature(s): lithic bedrock at 20 to 40 inches

Flooding: occasional

Ponding: none

Drainage class: very poorly drained

Soil loss tolerance (T factor): 2

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .24

Land capability, nonirrigated 3w

Hydric soil: yes

Hydrologic group: C/D

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|---------------------|---------------------|---------------------------------|------------|
| Ap,A -- 0 to 19 in | clay loam | moderate | 3.21 to 3.59 in | 7.4 to 8.4 |
| Bg -- 19 to 32 in | loam | moderate | 2.21 to 2.47 in | 7.4 to 8.4 |
| R -- 32 to 36 in | unweathered bedrock | very slow | | |

Map Unit Description (MN)

Redwood County, Minnesota

327A--Dickman sandy loam, 0 to 2 percent slopes

Dickman

Extent: 90 percent of the unit
Landform(s): outwash plains, terraces
Slope gradient: 0 to 2 percent
Parent material: outwash
Restrictive feature(s): greater than 60 inches
Flooding: none
Ponding: none
Drainage class: somewhat excessively drained

Soil loss tolerance (T factor): 3
Wind erodibility group (WEG): 3
Wind erodibility index (WEI): 86
Kw factor (surface layer) .20
Land capability, nonirrigated 3s
Hydric soil: no
Hydrologic group: A
Potential for frost action: low

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap,A -- 0 to 12 in | sandy loam | moderately rapid | 1.54 to 1.77 in | 6.1 to 6.5 |
| Bw -- 12 to 33 in | sandy loam | moderately rapid | 2.55 to 2.98 in | 6.1 to 7.3 |
| C -- 33 to 60 in | fine sand | rapid | 1.34 to 1.87 in | 6.1 to 7.8 |

327B--Dickman sandy loam, 2 to 6 percent slopes

Dickman

Extent: 90 percent of the unit
Landform(s): outwash plains, terraces
Slope gradient: 2 to 6 percent
Parent material: outwash
Restrictive feature(s): greater than 60 inches
Flooding: none
Ponding: none
Drainage class: somewhat excessively drained

Soil loss tolerance (T factor): 3
Wind erodibility group (WEG): 3
Wind erodibility index (WEI): 86
Kw factor (surface layer) .20
Land capability, nonirrigated 3e
Hydric soil: no
Hydrologic group: A
Potential for frost action: low

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap,A -- 0 to 12 in | sandy loam | moderately rapid | 1.54 to 1.77 in | 6.1 to 7.3 |
| Bw -- 12 to 33 in | sandy loam | moderately rapid | 2.55 to 2.98 in | 6.1 to 7.3 |
| C -- 33 to 60 in | fine sand | rapid | 1.34 to 1.87 in | 6.1 to 7.8 |

Map Unit Description (MN)

Redwood County, Minnesota

392--Biscay loam

Biscay

Extent: 85 percent of the unit

Landform(s): flats

Slope gradient: 0 to 2 percent

Parent material: outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 2w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|-----------------------------------------------|---------------------|---------------------------------|------------|
| Ap,A1,A2 -- 0 to 22 in | loam | moderate | 4.41 to 4.85 in | 6.1 to 7.3 |
| Bg -- 22 to 30 in | loam | moderate | 1.34 to 1.50 in | 6.6 to 7.3 |
| BCg -- 30 to 33 in | sandy loam | moderately rapid | 0.35 to 0.41 in | 7.4 to 8.4 |
| 2Cg -- 33 to 60 in | stratified gravelly coarse sand to loamy sand | rapid | 0.54 to 1.07 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

399--Biscay loam, depressional

Biscay, depressional

Extent: 86 percent of the unit

Landform(s): depressions

Slope gradient: 0 to 1 percent

Parent material: outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: frequent

Drainage class: very poorly drained

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 3w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|-----------------------------------------------|---------------------|---------------------------------|------------|
| Ap,A -- 0 to 31 in | loam | moderate | 6.22 to 6.84 in | 6.1 to 7.3 |
| Bg -- 31 to 35 in | sandy loam | moderately rapid | 0.47 to 0.55 in | 6.6 to 7.3 |
| 2Cg -- 35 to 60 in | stratified gravelly coarse sand to loamy sand | rapid | 0.50 to 0.99 in | 7.4 to 8.4 |

421B--Ves loam, 1 to 4 percent slopes

Ves

Extent: 85 percent of the unit

Landform(s): moraines

Slope gradient: 1 to 4 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .24

Land capability, nonirrigated 2e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap -- 0 to 10 in | loam | moderate | 1.97 to 2.17 in | 6.1 to 7.3 |
| Bw -- 10 to 25 in | loam | moderate | 2.61 to 2.92 in | 6.1 to 7.3 |
| Bk -- 25 to 39 in | loam | moderate | 2.34 to 2.62 in | 7.4 to 8.4 |
| C -- 39 to 60 in | loam | moderate | 3.55 to 3.96 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

421B2--Ves loam, 3 to 6 percent slopes, eroded

Ves, eroded

Extent: 85 percent of the unit

Landform(s): moraines

Slope gradient: 3 to 6 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 2e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap -- 0 to 10 in | loam | moderate | 1.97 to 2.17 in | 6.1 to 7.8 |
| Bw -- 10 to 25 in | loam | moderate | 2.61 to 2.92 in | 6.6 to 7.8 |
| Bk -- 25 to 39 in | loam | moderate | 2.34 to 2.62 in | 7.4 to 8.4 |
| C -- 39 to 60 in | loam | moderate | 3.55 to 3.96 in | 7.4 to 8.4 |

423--Seaforth loam

Seaforth

Extent: 85 percent of the unit

Landform(s): moraines

Slope gradient: 1 to 3 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .24

Land capability, nonirrigated 2s

Hydric soil: no

Hydrologic group: C

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap,A -- 0 to 15 in | loam | moderate | 2.99 to 3.29 in | 7.4 to 8.4 |
| Bk -- 15 to 21 in | clay loam | moderate | 0.89 to 1.12 in | 7.4 to 8.4 |
| C -- 21 to 60 in | loam | moderate | 6.63 to 7.41 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

562--Knoke silty clay loam

Knoke

Extent: 86 percent of the unit

Landform(s): depressions

Slope gradient: 0 to 1 percent

Parent material: lacustrine deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: frequent

Drainage class: very poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .32

Land capability, nonirrigated 3w

Hydric soil: yes

Hydrologic group: C/D

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|-----------------|---------------------|---------------------------------|------------|
| Ap,A1 -- 0 to 15 in | silty clay loam | moderately slow | 2.69 to 3.29 in | 7.4 to 8.4 |
| A2,Bg -- 15 to 60 in | clay loam | moderately slow | 7.63 to 8.53 in | 7.4 to 8.4 |

574--Du Page loam

Du Page, occasionally flooded

Extent: 85 percent of the unit

Landform(s): flood plains

Slope gradient: 0 to 2 percent

Parent material: alluvium

Restrictive feature(s): greater than 60 inches

Flooding: occasional

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .28

Land capability, nonirrigated 2w

Hydric soil: no

Hydrologic group: C

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap,A -- 0 to 32 in | loam | moderate | 6.38 to 7.02 in | 7.4 to 8.4 |
| C -- 32 to 60 in | loam | moderate | 4.75 to 5.31 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

575--Nishna clay loam

Nishna, occasionally flooded

Extent: 85 percent of the unit

Landform(s): flood plains

Slope gradient: 0 to 1 percent

Parent material: alluvium

Restrictive feature(s): greater than 60 inches

Flooding: occasional

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .32

Land capability, nonirrigated 3w

Hydric soil: yes

Hydrologic group: C/D

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|-----------------|---------------------|---------------------------------|------------|
| Ap,A -- 0 to 45 in | silty clay loam | very slow | 8.08 to 9.87 in | 7.4 to 8.4 |
| Cg -- 45 to 60 in | silty clay | very slow | 1.65 to 2.24 in | 7.4 to 8.4 |

654--Revere clay loam

Revere

Extent: 85 percent of the unit

Landform(s): flats

Slope gradient: 0 to 2 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .24

Land capability, nonirrigated 2w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap,Ay -- 0 to 15 in | clay loam | moderate | 2.54 to 2.84 in | 7.4 to 8.4 |
| Byg -- 15 to 35 in | clay loam | moderate | 3.01 to 3.81 in | 7.4 to 8.4 |
| Cyg,Cg -- 35 to 60 in | loam | moderate | 4.22 to 4.71 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

818--Lemond-Linder-Estherville complex

Lemond

Extent: 45 percent of the unit

Landform(s): beaches

Slope gradient: 0 to 2 percent

Parent material: outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .10

Land capability, nonirrigated 3w

Hydric soil: yes

Hydrologic group: A/D

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|---------------------|---------------------|---------------------------------|------------|
| Ap,A -- 0 to 17 in | gravelly sandy loam | moderately rapid | 2.20 to 2.54 in | 7.4 to 7.8 |
| Bg1 -- 17 to 28 in | sandy loam | moderately rapid | 1.32 to 1.54 in | 7.4 to 7.8 |
| Bg2 -- 28 to 44 in | loamy sand | rapid | 1.45 to 1.78 in | 7.4 to 8.4 |
| Cg -- 44 to 60 in | sandy loam | moderate | 1.73 to 2.05 in | 7.4 to 8.4 |

Linder

Extent: 25 percent of the unit

Landform(s): beaches

Slope gradient: 0 to 2 percent

Parent material: outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .20

Land capability, nonirrigated 2s

Hydric soil: no

Hydrologic group: B/D

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------------------|---------------------|---------------------------------|------------|
| Ap,A -- 0 to 18 in | sandy loam | moderate | 2.35 to 2.72 in | 6.1 to 7.3 |
| Bw,Bg -- 18 to 28 in | sandy loam | moderately rapid | 1.18 to 1.38 in | 6.1 to 7.3 |
| 2Cg -- 28 to 60 in | gravelly loamy coarse sand | very rapid | 2.55 to 3.19 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

818--Lemond-Linder-Estherville complex

Estherville

Extent: 25 percent of the unit

Landform(s): beaches

Slope gradient: 0 to 2 percent

Parent material: outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat excessively drained

Soil loss tolerance (T factor): 2

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .20

Land capability, nonirrigated 3s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------------|---------------------|---------------------------------|------------|
| Ap,A -- 0 to 14 in | sandy loam | moderately rapid | 1.79 to 2.07 in | 6.1 to 7.3 |
| Bw -- 14 to 19 in | sandy loam | moderately rapid | 0.61 to 0.72 in | 6.1 to 7.3 |
| 2C -- 19 to 60 in | gravelly coarse sand | rapid | 0.82 to 1.64 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

884--Delft-Webster complex

Delft

Extent: 50 percent of the unit

Landform(s): swales

Slope gradient: 1 to 2 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .24

Land capability, nonirrigated 2w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap -- 0 to 10 in | loam | moderate | 1.97 to 2.17 in | 6.1 to 7.3 |
| A1,AB -- 10 to 28 in | clay loam | moderate | 3.62 to 3.98 in | 6.1 to 7.3 |
| Bg -- 28 to 47 in | loam | moderate | 3.21 to 3.59 in | 6.6 to 7.8 |
| Cg -- 47 to 60 in | loam | moderate | 2.21 to 2.47 in | 7.4 to 8.4 |

Webster

Extent: 45 percent of the unit

Landform(s): swales

Slope gradient: 0 to 2 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .24

Land capability, nonirrigated 2w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap,A -- 0 to 20 in | clay loam | moderate | 3.41 to 3.81 in | 6.6 to 7.3 |
| Bg -- 20 to 49 in | clay loam | moderate | 4.31 to 5.46 in | 6.6 to 7.3 |
| Cg -- 49 to 60 in | loam | moderate | 1.87 to 2.09 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

954B2--Ves-Storden loams, 3 to 6 percent slopes, eroded

Ves, eroded

Extent: 50 percent of the unit

Landform(s): moraines

Slope gradient: 3 to 6 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 2e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap -- 0 to 10 in | loam | moderate | 1.97 to 2.17 in | 6.1 to 7.8 |
| Bw -- 10 to 24 in | loam | moderate | 2.41 to 2.69 in | 6.6 to 7.8 |
| Bk -- 24 to 39 in | loam | moderate | 2.54 to 2.84 in | 7.4 to 8.4 |
| C -- 39 to 60 in | loam | moderate | 3.55 to 3.96 in | 7.4 to 8.4 |

Storden, eroded

Extent: 35 percent of the unit

Landform(s): moraines

Slope gradient: 4 to 6 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .32

Land capability, nonirrigated 2e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap -- 0 to 8 in | loam | moderate | 1.57 to 1.73 in | 7.4 to 8.4 |
| C1 -- 8 to 37 in | loam | moderate | 4.95 to 5.54 in | 7.4 to 8.4 |
| C2 -- 37 to 60 in | loam | moderate | 3.88 to 4.34 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

954C2--Storden-Ves loams, 6 to 12 percent slopes, eroded

Storden, eroded

Extent: 50 percent of the unit

Landform(s): moraines

Slope gradient: 6 to 12 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .32

Land capability, nonirrigated 3e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap -- 0 to 8 in | loam | moderate | 1.57 to 1.73 in | 7.4 to 8.4 |
| C1 -- 8 to 37 in | loam | moderate | 4.95 to 5.54 in | 7.4 to 8.4 |
| C2 -- 37 to 60 in | loam | moderate | 3.88 to 4.34 in | 7.4 to 8.4 |

Ves, eroded

Extent: 25 percent of the unit

Landform(s): moraines

Slope gradient: 6 to 12 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 3e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap -- 0 to 10 in | loam | moderate | 1.97 to 2.17 in | 6.1 to 7.8 |
| Bw -- 10 to 24 in | loam | moderate | 2.41 to 2.69 in | 6.6 to 7.8 |
| Bk -- 24 to 39 in | loam | moderate | 2.54 to 2.84 in | 7.4 to 8.4 |
| C -- 39 to 60 in | loam | moderate | 3.55 to 3.96 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

954D2--Storden-Ves loams, 12 to 18 percent slopes, eroded

Storden, eroded

Extent: 60 percent of the unit

Landform(s): moraines

Slope gradient: 12 to 18 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .32

Land capability, nonirrigated 4e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap -- 0 to 8 in | loam | moderate | 1.57 to 1.73 in | 7.4 to 8.4 |
| C1 -- 8 to 37 in | loam | moderate | 4.95 to 5.54 in | 7.4 to 8.4 |
| C2 -- 37 to 60 in | loam | moderate | 3.88 to 4.34 in | 7.4 to 8.4 |

Ves, eroded

Extent: 20 percent of the unit

Landform(s): moraines

Slope gradient: 12 to 18 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 4e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap -- 0 to 10 in | loam | moderate | 1.97 to 2.17 in | 6.1 to 7.8 |
| Bw -- 10 to 24 in | loam | moderate | 2.41 to 2.69 in | 6.6 to 7.8 |
| Bk -- 24 to 39 in | loam | moderate | 2.54 to 2.84 in | 7.4 to 8.4 |
| C -- 39 to 60 in | loam | moderate | 3.55 to 3.96 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

992E--Rock outcrop-Copaston complex, 2 to 40 percent slopes

Rock outcrop

Extent: 50 percent of the unit

Landform(s): terraces

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil:

Hydrologic group:

Potential for frost action:

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|-----------|
|-------------------------------------|----------------|---------------------|---------------------------------|-----------|

Copaston

Extent: 35 percent of the unit

Landform(s): terraces

Slope gradient: 2 to 40 percent

Parent material: alluvium

Restrictive feature(s): lithic bedrock at 12 to 20 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 1

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .28

Land capability, nonirrigated 6e

Hydric soil: no

Hydrologic group: D

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|---------------------|---------------------|---------------------------------|------------|
| A,AB -- 0 to 14 in | sandy loam | moderate | 1.84 to 2.13 in | 6.1 to 7.3 |
| Bw -- 14 to 18 in | sandy loam | moderately rapid | 0.47 to 0.55 in | 6.1 to 7.8 |
| R -- 18 to 22 in | unweathered bedrock | very slow | | |

Map Unit Description (MN)

Redwood County, Minnesota

999B2--Ves-Estherville-Storden complex, 3 to 6 percent slopes, eroded

Ves, eroded

Extent: 30 percent of the unit

Landform(s): moraines

Slope gradient: 3 to 6 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 2e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap -- 0 to 10 in | loam | moderate | 1.97 to 2.17 in | 6.1 to 7.8 |
| Bw -- 10 to 22 in | loam | moderate | 2.07 to 2.32 in | 6.6 to 7.8 |
| Bk -- 22 to 39 in | loam | moderate | 2.88 to 3.22 in | 7.4 to 8.4 |
| C -- 39 to 60 in | loam | moderate | 3.55 to 3.96 in | 7.4 to 8.4 |

Estherville, eroded

Extent: 30 percent of the unit

Landform(s): moraines

Slope gradient: 3 to 6 percent

Parent material: outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat excessively drained

Soil loss tolerance (T factor): 2

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .20

Land capability, nonirrigated 3s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------------|---------------------|---------------------------------|------------|
| Ap,A -- 0 to 14 in | sandy loam | moderately rapid | 1.79 to 2.07 in | 6.1 to 7.3 |
| Bw -- 14 to 19 in | sandy loam | moderately rapid | 0.61 to 0.72 in | 6.1 to 7.3 |
| 2C -- 19 to 60 in | gravelly coarse sand | rapid | 0.82 to 1.64 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

999B2--Ves-Estherville-Storden complex, 3 to 6 percent slopes, eroded

Storden, eroded

Extent: 20 percent of the unit

Landform(s): moraines

Slope gradient: 4 to 6 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .32

Land capability, nonirrigated 2e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap -- 0 to 11 in | loam | moderate | 2.20 to 2.43 in | 7.4 to 8.4 |
| C1 -- 11 to 37 in | loam | moderate | 4.42 to 4.94 in | 7.4 to 8.4 |
| C2 -- 37 to 60 in | loam | moderate | 3.88 to 4.34 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

999C2--Storden-Estherville-Ves loams, 6 to 12 percent slopes, eroded

Storden, eroded

Extent: 30 percent of the unit

Landform(s): moraines

Slope gradient: 6 to 12 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .32

Land capability, nonirrigated 3e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap -- 0 to 11 in | loam | moderate | 2.20 to 2.43 in | 7.4 to 8.4 |
| C1 -- 11 to 37 in | loam | moderate | 4.42 to 4.94 in | 7.4 to 8.4 |
| C2 -- 37 to 60 in | loam | moderate | 3.88 to 4.34 in | 7.4 to 8.4 |

Estherville, eroded

Extent: 30 percent of the unit

Landform(s): moraines

Slope gradient: 6 to 12 percent

Parent material: outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat excessively drained

Soil loss tolerance (T factor): 2

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .32

Land capability, nonirrigated 4s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------------|---------------------|---------------------------------|------------|
| Ap -- 0 to 10 in | loam | moderately rapid | 1.97 to 2.17 in | 6.1 to 7.3 |
| Bw -- 10 to 19 in | sandy loam | moderately rapid | 1.09 to 1.27 in | 6.1 to 7.3 |
| 2C -- 19 to 60 in | gravelly coarse sand | rapid | 0.82 to 1.64 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

999C2--Storden-Estherville-Ves loams, 6 to 12 percent slopes, eroded

Ves, eroded

Extent: 20 percent of the unit

Landform(s): moraines

Slope gradient: 6 to 12 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 3e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap -- 0 to 10 in | loam | moderate | 1.97 to 2.17 in | 6.1 to 7.8 |
| Bw -- 10 to 22 in | loam | moderate | 2.07 to 2.32 in | 6.6 to 7.8 |
| Bk -- 22 to 39 in | loam | moderate | 2.88 to 3.22 in | 7.4 to 8.4 |
| C -- 39 to 60 in | loam | moderate | 3.55 to 3.96 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

999D2--Storden-Estherville-Ves complex, 12 to 18 percent slopes, eroded

Storden, eroded

Extent: 30 percent of the unit

Landform(s): moraines

Slope gradient: 12 to 18 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .32

Land capability, nonirrigated 4e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap -- 0 to 11 in | loam | moderate | 2.20 to 2.43 in | 7.4 to 8.4 |
| C1 -- 11 to 37 in | loam | moderate | 4.42 to 4.94 in | 7.4 to 8.4 |
| C2 -- 37 to 60 in | loam | moderate | 3.88 to 4.34 in | 7.4 to 8.4 |

Estherville, eroded

Extent: 30 percent of the unit

Landform(s): moraines

Slope gradient: 12 to 18 percent

Parent material: outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat excessively drained

Soil loss tolerance (T factor): 2

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .20

Land capability, nonirrigated 6e

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------------|---------------------|---------------------------------|------------|
| Ap -- 0 to 10 in | sandy loam | moderately rapid | 1.28 to 1.48 in | 6.1 to 7.3 |
| Bw -- 10 to 19 in | sandy loam | moderately rapid | 1.09 to 1.27 in | 6.1 to 7.3 |
| 2C -- 19 to 60 in | gravelly coarse sand | rapid | 0.82 to 1.64 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

999D2--Storden-Estherville-Ves complex, 12 to 18 percent slopes, eroded

Ves, eroded

Extent: 15 percent of the unit

Landform(s): moraines

Slope gradient: 12 to 18 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 4e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap -- 0 to 10 in | loam | moderate | 1.97 to 2.17 in | 6.1 to 7.8 |
| Bw -- 10 to 22 in | loam | moderate | 2.07 to 2.32 in | 6.6 to 7.8 |
| Bk -- 22 to 39 in | loam | moderate | 2.88 to 3.22 in | 7.4 to 8.4 |
| C -- 39 to 60 in | loam | moderate | 3.55 to 3.96 in | 7.4 to 8.4 |

1003B--Udorthents, (cut, and Fill land), 0 to 6 percent slopes

Udorthents, loamy (cut and fill land)

Extent: 100 percent of the unit

Landform(s): moraines

Slope gradient: 0 to 6 percent

Parent material: variable loamy material

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil:

Hydrologic group:

Potential for frost action:

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|-----------|
|-------------------------------------|----------------|---------------------|---------------------------------|-----------|

Map Unit Description (MN)

Redwood County, Minnesota

1016--Udorthents, loamy

Udorthents, loamy

Extent: 100 percent of the unit

Landform(s): moraines

Slope gradient: 0 to 2 percent

Parent material: variable soil material

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class: well drained

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil:

Hydrologic group:

Potential for frost action:

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|-----------|
| | | | | |

1024A--Havelock clay loam, 0 to 2 percent slopes. occasionally flooded

Havelock, occasionally flooded

Extent: 75 to 85 percent of the unit

Landform(s): flats on flood plains

Slope gradient: 0 to 2 percent

Parent material: alluvium

Restrictive feature(s): greater than 60 inches

Flooding: occasional

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .24

Land capability, nonirrigated 2w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap,A1,A2 -- 0 to 32 in | clay loam | moderate | 5.42 to 7.33 in | 7.4 to 8.4 |
| Cg -- 32 to 60 in | clay loam | moderate | 4.75 to 5.59 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

1029--Pits, gravel

Pits, gravel

Extent: 100 percent of the unit

Landform(s): outwash plains, terraces

Slope gradient:

Parent material: sandy and gravelly outwash

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil:

Hydrologic group:

Potential for frost action:

| | | | | |
|-------------------------------------|----------------|---------------------|---------------------------------|-----------|
| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|-----------|

1053--Aquolls, ponded

Aquolls, ponded

Extent: 100 percent of the unit

Landform(s): depressions

Slope gradient: 0 to 1 percent

Parent material: alluvium

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: frequent

Drainage class: very poorly drained

Soil loss tolerance (T factor):

Wind erodibility group (WEG): 8

Wind erodibility index (WEI): 0

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil: yes

Hydrologic group:

Potential for frost action:

| | | | | |
|-------------------------------------|----------------|---------------------|---------------------------------|-----------|
| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|-----------|

Map Unit Description (MN)

Redwood County, Minnesota

1056A--Spillville loam, 0 to 2 percent slopes. frequently flooded

Spillville, frequently flooded

Extent: 80 to 95 percent of the unit

Landform(s): flood plains

Slope gradient: 0 to 2 percent

Parent material: alluvium

Restrictive feature(s): greater than 60 inches

Flooding: frequent

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 5w

Hydric soil: no

Hydrologic group: C

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|-----------------|---------------------|---------------------------------|------------|
| A1,A2 -- 0 to 30 in | loam | moderate | 5.98 to 6.58 in | 6.1 to 7.3 |
| AC -- 30 to 50 in | fine sandy loam | moderately rapid | 2.21 to 2.61 in | 6.1 to 7.3 |
| C -- 50 to 60 in | fine sandy loam | moderately rapid | 1.08 to 1.28 in | 6.1 to 7.8 |

1833--Coland clay loam, occasionally flooded

Coland, occasionally flooded

Extent: 85 percent of the unit

Landform(s): flood plains

Slope gradient: 0 to 2 percent

Parent material: alluvium

Restrictive feature(s): greater than 60 inches

Flooding: occasional

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .24

Land capability, nonirrigated 2w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap, -- 0 to 10 in | clay loam | moderate | 1.67 to 1.87 in | 6.1 to 7.3 |
| A -- 10 to 39 in | clay loam | moderate | 4.95 to 5.54 in | 6.1 to 7.3 |
| Cg -- 39 to 60 in | loam | moderately rapid | 3.55 to 3.96 in | 7.4 to 7.8 |

Map Unit Description (MN)

Redwood County, Minnesota

1834--Coland clay loam, frequently flooded

Coland, frequently flooded

Extent: 85 percent of the unit

Landform(s): flood plains

Slope gradient: 0 to 2 percent

Parent material: alluvium

Restrictive feature(s): greater than 60 inches

Flooding: frequent

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .24

Land capability, nonirrigated 6w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap -- 0 to 10 in | clay loam | moderate | 1.67 to 1.87 in | 6.1 to 7.3 |
| A -- 10 to 39 in | clay loam | moderate | 4.95 to 5.54 in | 6.1 to 7.3 |
| Cg -- 39 to 60 in | loam | moderately rapid | 3.55 to 3.96 in | 7.4 to 7.8 |

1850--Oshawa variant stony clay loam

Oshawa, variant

Extent: 85 percent of the unit

Landform(s): flood plains

Slope gradient: 1 to 3 percent

Parent material: alluvium

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: frequent

Drainage class: very poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .15

Land capability, nonirrigated 6s

Hydric soil: yes

Hydrologic group: C/D

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|-----------------|---------------------|---------------------------------|------------|
| A1 -- 0 to 13 in | stony clay loam | moderately slow | 2.21 to 2.47 in | 7.4 to 8.4 |
| A2,A3 -- 13 to 42 in | clay loam | moderately slow | 4.95 to 5.54 in | 7.4 to 8.4 |
| Cg -- 42 to 60 in | silty clay loam | moderately slow | 2.83 to 3.37 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

1851B--Blue Earth mucky clay loam, sloping

Blue Earth, sloping

Extent: 85 percent of the unit

Landform(s): swales

Slope gradient: 1 to 3 percent

Parent material: lacustrine deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: very poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .32

Land capability, nonirrigated 7w

Hydric soil: yes

Hydrologic group: C/D

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|-----------------|---------------------|---------------------------------|------------|
| A -- 0 to 10 in | mucky clay loam | moderately slow | 1.77 to 2.36 in | 7.4 to 8.4 |
| Cg -- 10 to 52 in | mucky clay loam | moderately slow | 7.58 to 10.11 in | 7.4 to 8.4 |
| 2Cg -- 52 to 60 in | clay loam | moderately slow | 1.10 to 1.26 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

1852F--Terril-Swanlake loams, 25 to 70 percent slopes

Terril

Extent: 50 percent of the unit

Landform(s): moraines

Slope gradient: 25 to 60 percent

Parent material: colluvium

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 7e

Hydric soil: no

Hydrologic group: C

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| A,A1 -- 0 to 24 in | loam | moderate | 4.80 to 5.28 in | 6.1 to 7.3 |
| A2 -- 24 to 36 in | loam | moderate | 2.36 to 2.60 in | 6.1 to 7.3 |
| Bw -- 36 to 60 in | loam | moderate | 4.08 to 4.56 in | 6.1 to 7.3 |

Swanlake

Extent: 30 percent of the unit

Landform(s): moraines

Slope gradient: 25 to 70 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .32

Land capability, nonirrigated 7e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| A -- 0 to 9 in | loam | moderate | 1.81 to 1.99 in | 7.4 to 8.4 |
| Bk1 -- 9 to 17 in | loam | moderate | 1.34 to 1.50 in | 7.4 to 8.4 |
| Bk2,C -- 17 to 60 in | loam | moderate | 7.30 to 8.15 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

1853A--Wadena variant loam, 0 to 2 percent slopes

Wadena, variant

Extent: 90 percent of the unit

Landform(s): terraces

Slope gradient: 0 to 2 percent

Parent material: outwash

Restrictive feature(s): lithic bedrock at 20 to 40 inches

Flooding: none

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 2

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 2s

Hydric soil: no

Hydrologic group: C

Potential for frost action: low

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|-------------------|---------------------|---------------------------------|------------|
| Ap -- 0 to 11 in | loam | moderately rapid | 2.20 to 2.43 in | 6.6 to 7.3 |
| Bw -- 11 to 18 in | loam | moderate | 1.20 to 1.35 in | 6.6 to 7.3 |
| C -- 18 to 32 in | loam | moderately rapid | 2.34 to 2.62 in | 7.4 to 7.8 |
| R -- 32 to 36 in | weathered bedrock | very slow | | |

Map Unit Description (MN)

Redwood County, Minnesota

1853B--Wadena variant loam, 2 to 6 percent slopes

Wadena, variant

Extent: 90 percent of the unit

Landform(s): terraces

Slope gradient: 2 to 6 percent

Parent material: outwash

Restrictive feature(s): lithic bedrock at 20 to 40 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 2

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 2e

Hydric soil: no

Hydrologic group: C

Potential for frost action: low

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|-------------------|---------------------|---------------------------------|------------|
| Ap -- 0 to 11 in | loam | moderately rapid | 2.20 to 2.43 in | 6.6 to 7.3 |
| Bw -- 11 to 18 in | loam | moderate | 1.20 to 1.35 in | 6.6 to 7.3 |
| C -- 18 to 32 in | loam | moderately rapid | 2.34 to 2.62 in | 7.4 to 7.8 |
| R -- 32 to 36 in | weathered bedrock | very slow | | |

Map Unit Description (MN)

Redwood County, Minnesota

1897--Seaforth-Wilmonton clay loams

Seaforth

Extent: 50 percent of the unit

Landform(s): moraines

Slope gradient: 1 to 3 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .24

Land capability, nonirrigated 2s

Hydric soil: no

Hydrologic group: C

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap,A -- 0 to 12 in | loam | moderate | 2.36 to 2.60 in | 7.4 to 8.4 |
| Bk -- 12 to 30 in | loam | moderate | 3.08 to 3.44 in | 7.4 to 8.4 |
| C -- 30 to 60 in | loam | moderate | 5.09 to 5.69 in | 7.4 to 8.4 |

Wilmonton

Extent: 30 percent of the unit

Landform(s): moraines

Slope gradient: 1 to 3 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .24

Land capability, nonirrigated 1

Hydric soil: no

Hydrologic group: C

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap,A -- 0 to 14 in | clay loam | moderate | 2.41 to 2.69 in | 6.1 to 7.3 |
| Bw -- 14 to 37 in | clay loam | moderately slow | 3.43 to 4.34 in | 6.1 to 7.3 |
| 2Bk,2C -- 37 to 60 in | clay loam | moderately slow | 3.20 to 3.65 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

1899B--Wilmonton variant loam, 2 to 12 percent slopes

Wilmonton, variant

Extent: 85 percent of the unit

Landform(s): moraines

Slope gradient: 2 to 12 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .37

Land capability, nonirrigated 3e

Hydric soil: no

Hydrologic group: D

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap,A -- 0 to 19 in | loam | moderate | 3.78 to 4.16 in | 6.6 to 7.3 |
| 2Bt -- 19 to 41 in | clay | very slow | 2.20 to 3.09 in | 6.6 to 7.3 |
| 2Cg -- 41 to 60 in | clay | very slow | 1.70 to 2.46 in | 7.4 to 8.4 |

1899E--Wilmonton variant sandy clay loam, 12 to 40 percent slopes

Wilmonton, variant

Extent: 85 percent of the unit

Landform(s): moraines

Slope gradient: 12 to 40 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .24

Land capability, nonirrigated 7e

Hydric soil: no

Hydrologic group: D

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|-----------------|---------------------|---------------------------------|------------|
| A -- 0 to 9 in | sandy clay loam | moderate | 1.63 to 1.81 in | 6.6 to 7.3 |
| 2Bt -- 9 to 29 in | clay | very slow | 2.01 to 2.81 in | 6.6 to 7.3 |
| 2Cg -- 29 to 60 in | clay | very slow | 2.76 to 3.99 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

GP--Pits, gravel-Udipsamments complex

Pits, gravel

Extent: 50 to 100 percent of the unit
Landform(s): outwash plains, stream terraces
Slope gradient:
Parent material: outwash
Restrictive feature(s): greater than 60 inches
Flooding:
Ponding:
Drainage class:

Soil loss tolerance (T factor):
Wind erodibility group (WEG):
Wind erodibility index (WEI):
Kw factor (surface layer)
Land capability, nonirrigated
Hydric soil: unranked
Hydrologic group:
Potential for frost action:

| | | | | |
|-------------------------------------|----------------|---------------------|---------------------------------|-----------|
| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|-----------|

Udipsamments

Extent: 15 to 30 percent of the unit
Landform(s): outwash plains, stream terraces
Slope gradient:
Parent material: outwash
Restrictive feature(s): greater than 60 inches
Flooding:
Ponding:
Drainage class:

Soil loss tolerance (T factor):
Wind erodibility group (WEG):
Wind erodibility index (WEI):
Kw factor (surface layer)
Land capability, nonirrigated
Hydric soil: unranked
Hydrologic group:
Potential for frost action:

| | | | | |
|-------------------------------------|----------------|---------------------|---------------------------------|-----------|
| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|-----------|

Map Unit Description (MN)

Redwood County, Minnesota

L83A--Webster clay loam, 0 to 2 percent slopes

Webster

Extent: 50 to 85 percent of the unit

Landform(s): flats on moraines, swales on moraines

Slope gradient: 0 to 2 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .24

Land capability, nonirrigated 2w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap,A -- 0 to 19 in | clay loam | moderate | 3.59 to 3.97 in | 6.6 to 7.3 |
| Bg -- 19 to 26 in | clay loam | moderate | 1.13 to 1.28 in | 6.6 to 7.8 |
| BCg,Cg -- 26 to 60 in | loam | moderate | 5.08 to 6.43 in | 7.4 to 8.4 |

L84A--Glencoe clay loam, depressional, 0 to 1 percent slopes

Glencoe, depressional

Extent: 75 to 100 percent of the unit

Landform(s): depressions on moraines

Slope gradient: 0 to 1 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: frequent

Drainage class: very poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .24

Land capability, nonirrigated 3w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap,A -- 0 to 24 in | clay loam | moderate | 4.32 to 5.28 in | 6.1 to 7.8 |
| ABg -- 24 to 35 in | clay loam | moderate | 1.98 to 2.43 in | 6.1 to 7.8 |
| Bg -- 35 to 48 in | loam | moderate | 1.95 to 2.47 in | 6.6 to 7.8 |
| Cg -- 48 to 60 in | loam | moderate | 1.77 to 2.24 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

L141A--Spillville loam, 0 to 2 percent slopes, occasionally flooded

Spillville, occasionally flooded

Extent: 80 to 90 percent of the unit

Landform(s): flats on flood plains, rises on flood plains

Slope gradient: 0 to 2 percent

Parent material: alluvium

Restrictive feature(s): greater than 60 inches

Flooding: occasional

Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 2w

Hydric soil: no

Hydrologic group: B/D

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap,A -- 0 to 51 in | loam | moderate | 9.72 to 10.75 in | 5.6 to 7.3 |
| C -- 51 to 60 in | loam | moderately rapid | 1.30 to 1.56 in | 5.6 to 7.3 |

L142A--Jeffers clay loam, 0 to 2 percent slopes

Jeffers

Extent: 80 to 95 percent of the unit

Landform(s): rims on depressions on moraines, flats on moraines

Slope gradient: 0 to 2 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .20

Land capability, nonirrigated 2w

Hydric soil: yes

Hydrologic group: C/D

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap,A -- 0 to 18 in | clay loam | moderate | 3.08 to 3.44 in | 7.4 to 8.4 |
| Bg -- 18 to 22 in | clay loam | moderate | 0.59 to 0.75 in | 7.9 to 8.4 |
| Bkg,Bk -- 22 to 35 in | clay loam | moderate | 1.95 to 2.47 in | 7.9 to 8.4 |
| BC1,BC2 -- 35 to 60 in | clay loam | moderately slow | 3.47 to 3.97 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

L149A--Romnell clay loam, 0 to 3 percent slopes

Romnell

Extent: 80 to 95 percent of the unit

Landform(s): drainageways on moraines, swales on moraines

Slope gradient: 0 to 3 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .20

Land capability, nonirrigated 2w

Hydric soil: yes

Hydrologic group: C/D

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap,A -- 0 to 18 in | clay loam | moderate | 3.08 to 4.35 in | 6.6 to 7.8 |
| Bg -- 18 to 27 in | clay loam | moderate | 1.54 to 2.17 in | 6.6 to 7.8 |
| Btyg -- 27 to 33 in | clay loam | moderately slow | 0.89 to 1.12 in | 6.6 to 7.8 |
| Bkg -- 33 to 43 in | clay loam | moderately slow | 1.38 to 1.57 in | 7.4 to 8.4 |
| BCkg -- 43 to 80 in | clay loam | moderately slow | 5.18 to 5.92 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

L163A--Okoboji silty clay loam, depressional, 0 to 1 percent slopes

Okoboji, depressional

Extent: 70 to 95 percent of the unit

Landform(s): depressions on lake plains, depressions on moraines

Slope gradient: 0 to 1 percent

Parent material: lacustrine sediments over till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: frequent

Drainage class: very poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .28

Land capability, nonirrigated 3w

Hydric soil: yes

Hydrologic group: C/D

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|-----------------|---------------------|---------------------------------|------------|
| Ap,A -- 0 to 26 in | silty clay loam | moderately slow | 5.46 to 5.98 in | 6.1 to 7.8 |
| Bg -- 26 to 42 in | silty clay | moderately slow | 2.91 to 3.23 in | 6.6 to 7.8 |
| Cg -- 42 to 60 in | silty clay loam | moderately slow | 3.19 to 3.54 in | 6.6 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

L172D2--Storden, firm till-Annton complex, 12 to 18 percent slopes, moderately eroded

Storden, moderately eroded, firm till

Extent: 40 to 60 percent of the unit

Landform(s): hills on moraines

Slope gradient: 12 to 18 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .32

Land capability, nonirrigated 4e

Hydric soil: no

Hydrologic group: C

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap -- 0 to 8 in | loam | moderate | 1.57 to 1.73 in | 7.4 to 8.4 |
| Bk -- 8 to 20 in | loam | moderate | 2.07 to 2.32 in | 7.9 to 8.4 |
| BC -- 20 to 80 in | clay loam | moderately slow | 8.38 to 9.57 in | 7.4 to 8.4 |

Annton, moderately, eroded

Extent: 20 to 45 percent of the unit

Landform(s): hills on moraines

Slope gradient: 12 to 18 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .24

Land capability, nonirrigated 3e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap -- 0 to 8 in | clay loam | moderate | 1.42 to 1.73 in | 6.1 to 7.3 |
| Bw -- 8 to 26 in | clay loam | moderate | 3.08 to 3.44 in | 6.1 to 7.3 |
| Bk -- 26 to 60 in | loam | moderate | 5.76 to 6.43 in | 7.9 to 8.4 |
| BC -- 60 to 80 in | clay loam | moderately slow | 2.81 to 3.21 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

L173A--Moines clay loam, 1 to 3 percent slopes

Moines

Extent: 80 to 95 percent of the unit

Landform(s): flats on moraines, rises on moraines

Slope gradient: 1 to 3 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .24

Land capability, nonirrigated 2s

Hydric soil: no

Hydrologic group: B/D

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap,ABy -- 0 to 14 in | clay loam | moderate | 2.41 to 2.69 in | 7.4 to 8.4 |
| Byg1,Byg2,Bk - 14 to 40 in | loam | moderate | 4.42 to 4.94 in | 7.9 to 8.4 |
| - | | | | |
| BCg -- 40 to 60 in | clay loam | moderately slow | 2.76 to 3.15 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

L198A--North Twin-Walnut grove complex, 0 to 2 percent slopes

North Twin

Extent: 60 to 80 percent of the unit

Landform(s): hills on moraines

Slope gradient: 1 to 2 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 1

Hydric soil: no

Hydrologic group: C

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap -- 0 to 10 in | clay loam | moderately slow | 1.67 to 1.87 in | 6.6 to 7.3 |
| Bw -- 10 to 18 in | clay loam | moderately slow | 1.16 to 1.49 in | 6.1 to 7.3 |
| Bk -- 18 to 31 in | clay loam | moderately slow | 1.82 to 2.08 in | 7.4 to 8.4 |
| BCg,BC1,BC2 -- 31 to 80 in | clay loam | moderately slow | 6.83 to 7.81 in | 7.4 to 8.4 |

Walnut Grove

Extent: 15 to 25 percent of the unit

Landform(s): flats on moraines, rises on moraines

Slope gradient: 0 to 2 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 1

Hydric soil: no

Hydrologic group: C/D

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap,A -- 0 to 14 in | clay loam | moderately slow | 2.41 to 2.69 in | 6.6 to 7.3 |
| Bw -- 14 to 28 in | clay loam | moderate | 2.48 to 3.31 in | 6.1 to 7.3 |
| Bk -- 28 to 39 in | clay loam | moderately slow | 1.54 to 1.76 in | 7.4 to 8.4 |
| BC1,BC2 -- 39 to 80 in | clay loam | moderately slow | 5.73 to 6.55 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

L198B--North Twin-Walnut grove complex, 1 to 4 percent slopes

North Twin

Extent: 50 to 70 percent of the unit

Landform(s): hills on moraines

Slope gradient: 1 to 4 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 2e

Hydric soil: no

Hydrologic group: C

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap -- 0 to 10 in | clay loam | moderately slow | 1.67 to 1.87 in | 6.6 to 7.3 |
| Bw -- 10 to 18 in | clay loam | moderately slow | 1.16 to 1.49 in | 6.1 to 7.3 |
| Bk -- 18 to 31 in | clay loam | moderately slow | 1.82 to 2.08 in | 7.4 to 8.4 |
| BCg,BC1,BC2 -- 31 to 80 in | clay loam | moderately slow | 6.83 to 7.81 in | 7.4 to 8.4 |

Walnut Grove

Extent: 15 to 35 percent of the unit

Landform(s): flats on moraines, rises on moraines

Slope gradient: 1 to 3 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 1

Hydric soil: no

Hydrologic group: C/D

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap,A -- 0 to 14 in | clay loam | moderately slow | 2.41 to 2.69 in | 6.6 to 7.3 |
| Bw -- 14 to 28 in | clay loam | moderate | 2.48 to 3.31 in | 6.1 to 7.3 |
| Bk -- 28 to 39 in | clay loam | moderately slow | 1.54 to 1.76 in | 7.4 to 8.4 |
| BC1,BC2 -- 39 to 80 in | clay loam | moderately slow | 5.73 to 6.55 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

L201A--Normania loam, 0 to 3 percent slopes

Normania

Extent: 75 to 90 percent of the unit

Landform(s): flats on moraines, rises on moraines

Slope gradient: 0 to 3 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 1

Hydric soil: no

Hydrologic group: B/D

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap,A,AB -- 0 to 17 in | loam | moderate | 3.39 to 3.72 in | 6.1 to 7.3 |
| Bw -- 17 to 26 in | loam | moderate | 1.36 to 1.72 in | 6.6 to 7.3 |
| Bk -- 26 to 50 in | loam | moderate | 3.60 to 4.56 in | 7.4 to 8.4 |
| Cg -- 50 to 60 in | loam | moderate | 1.48 to 1.87 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

L202A--Pell Creek-Romnell complex, 0 to 2 percent slopes

Pell Creek

| | |
|-------------------------------------------------------------------------------------|------------------------------------------|
| <i>Extent:</i> 35 to 70 percent of the unit | <i>Soil loss tolerance (T factor):</i> 5 |
| <i>Landform(s):</i> drainageways on moraines, flats on moraines, swales on moraines | <i>Wind erodibility group (WEG):</i> 6 |
| <i>Slope gradient:</i> 0 to 2 percent | <i>Wind erodibility index (WEI):</i> 48 |
| <i>Parent material:</i> till | <i>Kw factor (surface layer)</i> .24 |
| <i>Restrictive feature(s):</i> greater than 60 inches | <i>Land capability, nonirrigated</i> 2w |
| <i>Flooding:</i> none | <i>Hydric soil:</i> yes |
| <i>Ponding:</i> none | <i>Hydrologic group:</i> C/D |
| <i>Drainage class:</i> poorly drained | <i>Potential for frost action:</i> high |

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap,A,AB -- 0 to 20 in | clay loam | moderate | 3.41 to 3.81 in | 6.1 to 7.3 |
| Bg -- 20 to 34 in | clay loam | moderate | 2.07 to 2.62 in | 7.4 to 8.4 |
| BCg,BC1,BC2 -- 34 to 80 in | clay loam | moderately slow | 6.45 to 7.37 in | 7.4 to 8.4 |

Romnell

| | |
|------------------------------------------------------------------|------------------------------------------|
| <i>Extent:</i> 15 to 45 percent of the unit | <i>Soil loss tolerance (T factor):</i> 5 |
| <i>Landform(s):</i> drainageways on moraines, swales on moraines | <i>Wind erodibility group (WEG):</i> 4L |
| <i>Slope gradient:</i> 0 to 2 percent | <i>Wind erodibility index (WEI):</i> 86 |
| <i>Parent material:</i> till | <i>Kw factor (surface layer)</i> .20 |
| <i>Restrictive feature(s):</i> greater than 60 inches | <i>Land capability, nonirrigated</i> 2w |
| <i>Flooding:</i> none | <i>Hydric soil:</i> yes |
| <i>Ponding:</i> none | <i>Hydrologic group:</i> C/D |
| <i>Drainage class:</i> poorly drained | <i>Potential for frost action:</i> high |

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap,A -- 0 to 18 in | clay loam | moderate | 3.08 to 4.35 in | 6.6 to 7.8 |
| Bg -- 18 to 27 in | clay loam | moderate | 1.54 to 2.17 in | 6.6 to 7.8 |
| Btyg -- 27 to 33 in | clay loam | moderately slow | 0.89 to 1.12 in | 6.6 to 7.8 |
| Bkg -- 33 to 43 in | clay loam | moderately slow | 1.38 to 1.57 in | 7.4 to 8.4 |
| BCkg -- 43 to 60 in | clay loam | moderately slow | 2.37 to 2.71 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

L207E--Belview-Ridgeton complex, 18 to 25 percent slopes, firm till substratum

Belview, firm till substratum

Extent: 65 to 85 percent of the unit

Landform(s): escarpments on moraines

Slope gradient: 18 to 25 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .28

Land capability, nonirrigated 6e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| A -- 0 to 8 in | loam | moderate | 1.57 to 1.73 in | 7.4 to 8.4 |
| Bk -- 8 to 48 in | loam | moderate | 6.02 to 7.63 in | 7.4 to 8.4 |
| BC1,2 -- 48 to 80 in | clay loam | moderately slow | 4.46 to 5.10 in | 7.4 to 8.4 |

Ridgeton, firm till substratum

Extent: 10 to 20 percent of the unit

Landform(s): escarpments on moraines

Slope gradient: 18 to 25 percent

Parent material: colluvium over till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 6e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| A1,A2,AB -- 0 to 31 in | loam | moderate | 6.22 to 6.84 in | 6.1 to 7.3 |
| BA -- 31 to 49 in | loam | moderate | 3.54 to 3.90 in | 6.1 to 7.3 |
| Bw -- 49 to 64 in | clay loam | moderate | 2.46 to 2.76 in | 6.1 to 7.3 |
| BC -- 64 to 80 in | clay loam | moderately slow | 2.20 to 2.52 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

L207F--Belview-Ridgeton complex, 18 to 40 percent slopes, firm till substratum

Belview, firm till substratum

Extent: 65 to 85 percent of the unit

Landform(s): escarpments on moraines

Slope gradient: 18 to 40 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .28

Land capability, nonirrigated 6e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| A -- 0 to 8 in | loam | moderate | 1.57 to 1.73 in | 7.4 to 8.4 |
| Bk -- 8 to 48 in | loam | moderate | 6.02 to 7.63 in | 7.4 to 8.4 |
| BC1,BC2 -- 48 to 80 in | clay loam | moderately slow | 4.46 to 5.10 in | 7.4 to 8.4 |

Ridgeton, firm till substratum

Extent: 10 to 20 percent of the unit

Landform(s): escarpments on moraines

Slope gradient: 18 to 35 percent

Parent material: colluvium over till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 6e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| A1,A2,AB -- 0 to 31 in | loam | moderate | 6.22 to 6.84 in | 6.1 to 7.3 |
| BA -- 31 to 49 in | loam | moderate | 3.54 to 3.90 in | 6.1 to 7.3 |
| Bw -- 49 to 64 in | clay loam | moderate | 2.46 to 2.76 in | 6.1 to 7.3 |
| BC -- 64 to 80 in | clay loam | moderately slow | 2.20 to 2.52 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

L214A--Calco-Du Page complex, 0 to 2 percent slopes, frequently flooded

Calco, frequently flooded

Extent: 40 to 70 percent of the unit

Landform(s): flats on flood plains

Slope gradient: 0 to 2 percent

Parent material: alluvium

Restrictive feature(s): greater than 60 inches

Flooding: frequent

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .32

Land capability, nonirrigated 5w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|-----------------|---------------------|---------------------------------|------------|
| Ap,A1 -- 0 to 14 in | silty clay loam | moderate | 2.98 to 3.26 in | 7.4 to 8.4 |
| A2 -- 14 to 40 in | silty clay loam | moderate | 5.46 to 5.98 in | 7.4 to 8.4 |
| Cg -- 40 to 60 in | silty clay loam | moderate | 3.54 to 3.94 in | 7.4 to 8.4 |

Du Page, frequently flooded

Extent: 40 to 60 percent of the unit

Landform(s): flats on flood plains, rises on flood plains

Slope gradient: 0 to 2 percent

Parent material: alluvium

Restrictive feature(s): greater than 60 inches

Flooding: frequent

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .28

Land capability, nonirrigated 5w

Hydric soil: yes

Hydrologic group: C

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| A1,A2 -- 0 to 36 in | loam | moderate | 7.17 to 7.88 in | 6.6 to 8.4 |
| C1,C2 -- 36 to 60 in | loam | moderate | 4.08 to 4.56 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

L219A--Coland clay loam, 0 to 2 percent slopes, occasionally flooded

Coland, occasionally flooded

Extent: 65 to 90 percent of the unit

Landform(s): flats on flood plains

Slope gradient: 0 to 2 percent

Parent material: alluvium

Restrictive feature(s): greater than 60 inches

Flooding: occasional

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .24

Land capability, nonirrigated 2w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap,A1,A2,A3 -- 0 to 42 in | clay loam | moderate | 8.43 to 9.27 in | 6.1 to 7.3 |
| AB -- 42 to 55 in | clay loam | moderate | 2.60 to 2.86 in | 6.1 to 7.3 |
| Cg -- 55 to 60 in | clay loam | moderate | 0.94 to 1.04 in | 6.1 to 7.3 |

L221A--Du Page loam, 0 to 2 percent slopes, occasionally flooded

Du Page, occasionally flooded

Extent: 75 to 90 percent of the unit

Landform(s): flats on flood plains, rises on flood plains

Slope gradient: 0 to 2 percent

Parent material: alluvium

Restrictive feature(s): greater than 60 inches

Flooding: occasional

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .28

Land capability, nonirrigated 2w

Hydric soil: no

Hydrologic group: C

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap,A1,A2 -- 0 to 36 in | loam | moderate | 7.17 to 7.88 in | 6.6 to 8.4 |
| C1,C2 -- 36 to 60 in | loam | moderate | 4.08 to 4.56 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

L224A--Coland clay loam, 0 to 2 percent slopes, frequently flooded

Coland, frequently flooded

Extent: 65 to 90 percent of the unit

Landform(s): flats on flood plains

Slope gradient: 0 to 2 percent

Parent material: alluvium

Restrictive feature(s): greater than 60 inches

Flooding: frequent

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .24

Land capability, nonirrigated 5w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| A -- 0 to 42 in | clay loam | moderate | 8.43 to 9.27 in | 6.1 to 7.3 |
| C -- 42 to 60 in | clay loam | moderate | 3.54 to 3.90 in | 6.1 to 7.3 |

Map Unit Description (MN)

Redwood County, Minnesota

L225B--Annton-North Twin complex, 3 to 6 percent slopes

Annton

Extent: 40 to 55 percent of the unit

Landform(s): hills on moraines

Slope gradient: 3 to 6 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .24

Land capability, nonirrigated 2e

Hydric soil: no

Hydrologic group: C

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap -- 0 to 10 in | clay loam | moderate | 1.77 to 2.17 in | 6.1 to 7.3 |
| Bw -- 10 to 22 in | clay loam | moderate | 2.07 to 2.32 in | 6.1 to 7.3 |
| Bk -- 22 to 60 in | loam | moderately slow | 5.29 to 6.05 in | 7.4 to 8.4 |
| BC -- 60 to 80 in | clay loam | moderately slow | 2.81 to 3.21 in | 7.4 to 8.4 |

North Twin

Extent: 15 to 45 percent of the unit

Landform(s): hills on moraines

Slope gradient: 3 to 4 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 2e

Hydric soil: no

Hydrologic group: C

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap -- 0 to 10 in | clay loam | moderately slow | 1.67 to 1.87 in | 6.6 to 7.3 |
| Bw -- 10 to 18 in | clay loam | moderately slow | 1.16 to 1.49 in | 6.1 to 7.3 |
| Bk -- 18 to 31 in | clay loam | moderately slow | 1.82 to 2.08 in | 7.4 to 8.4 |
| BCg,BC1,BC2 -- 31 to 80 in | clay loam | moderately slow | 6.83 to 7.81 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

L226C2--Annton-Storden, firm till complex, 6 to 12 percent slopes, moderately eroded

Annton, moderately eroded

Extent: 30 to 60 percent of the unit

Landform(s): hills on moraines

Slope gradient: 6 to 12 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .24

Land capability, nonirrigated 3e

Hydric soil: no

Hydrologic group: C

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap -- 0 to 8 in | clay loam | moderate | 1.42 to 1.73 in | 6.1 to 7.3 |
| Bw -- 8 to 21 in | clay loam | moderate | 2.21 to 2.47 in | 6.1 to 7.3 |
| Bk -- 21 to 34 in | clay loam | moderately slow | 1.82 to 2.08 in | 7.4 to 8.4 |
| BC1,BC2 -- 34 to 80 in | clay loam | moderately slow | 6.45 to 7.37 in | 7.4 to 8.4 |

Storden, moderately eroded, firm till

Extent: 25 to 50 percent of the unit

Landform(s): hills on moraines

Slope gradient: 6 to 12 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .37

Land capability, nonirrigated 3e

Hydric soil: no

Hydrologic group: C

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap -- 0 to 10 in | loam | moderate | 1.97 to 2.17 in | 7.4 to 8.4 |
| Bk -- 10 to 31 in | clay loam | moderately slow | 2.98 to 3.40 in | 7.4 to 8.4 |
| BC -- 31 to 80 in | clay loam | moderately slow | 6.83 to 7.81 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

L229A--Romnell silty clay loam, depressional, 0 to 1 percent slopes

Romnell, depressional

Extent: 75 to 95 percent of the unit

Landform(s): depressions on moraines

Slope gradient: 0 to 1 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: frequent

Drainage class: very poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .28

Land capability, nonirrigated 3w

Hydric soil: yes

Hydrologic group: C/D

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|-----------------|---------------------|---------------------------------|------------|
| Ap,A -- 0 to 23 in | silty clay loam | moderately slow | 4.80 to 5.25 in | 6.1 to 7.8 |
| Bg1,Bg2 -- 23 to 43 in | silty clay loam | moderate | 3.41 to 4.82 in | 6.6 to 7.8 |
| BCg1 -- 43 to 51 in | clay loam | moderately slow | 1.24 to 1.57 in | 6.6 to 7.8 |
| BCg2 -- 51 to 80 in | clay loam | moderately slow | 4.02 to 4.60 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

L241B--Dickinson fine sandy loam, firm till substratum, 1 to 6 percent slopes

Dickinson, firm till substratum

Extent: 70 to 90 percent of the unit

Landform(s): hills on moraines

Slope gradient: 1 to 6 percent

Parent material: outwash over till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .20

Land capability, nonirrigated 3e

Hydric soil: no

Hydrologic group: A

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|-----------------|---------------------|---------------------------------|------------|
| Ap,AB -- 0 to 16 in | fine sandy loam | moderately rapid | 1.94 to 2.42 in | 5.6 to 7.3 |
| Bw -- 16 to 30 in | fine sandy loam | moderately rapid | 1.65 to 2.07 in | 5.1 to 6.5 |
| C1 -- 30 to 47 in | loamy sand | rapid | 0.34 to 0.68 in | 5.6 to 7.8 |
| C2 -- 47 to 60 in | sand | rapid | 0.26 to 0.52 in | 5.6 to 7.8 |
| 2BC -- 60 to 80 in | clay loam | moderately slow | 2.81 to 3.21 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

L242B--Terril loam, firm till substratum, 2 to 6 percent slopes

Terril, firm till substratum

Extent: 80 to 95 percent of the unit

Landform(s): hills on moraines

Slope gradient: 2 to 6 percent

Parent material: colluvium over till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 2e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap,A1,A2 -- 0 to 30 in | loam | moderate | 5.98 to 6.58 in | 6.1 to 7.3 |
| A3,AB -- 30 to 40 in | loam | moderate | 1.74 to 1.94 in | 6.1 to 7.3 |
| Bw -- 40 to 61 in | loam | moderate | 3.34 to 3.76 in | 6.1 to 7.3 |
| BC -- 61 to 80 in | clay loam | moderately slow | 2.65 to 3.02 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

L243A--Havelock-Zumbro complex, 0 to 3 percent slopes, frequently flooded

Havelock, frequently flooded

Extent: 50 to 85 percent of the unit

Landform(s): flats on flood plains

Slope gradient: 0 to 2 percent

Parent material: alluvium

Restrictive feature(s): greater than 60 inches

Flooding: frequent

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .28

Land capability, nonirrigated 5w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| A1,A2 -- 0 to 32 in | loam | moderate | 5.42 to 7.33 in | 7.4 to 8.4 |
| Cg -- 32 to 60 in | clay loam | moderate | 4.75 to 5.59 in | 7.4 to 8.4 |

Zumbro, frequently flooded

Extent: 10 to 20 percent of the unit

Landform(s): flood plains

Slope gradient: 0 to 2 percent

Parent material: alluvium

Restrictive feature(s): greater than 60 inches

Flooding: frequent

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .28

Land capability, nonirrigated 3s

Hydric soil: no

Hydrologic group: B

Potential for frost action: low

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|-----------------|---------------------|---------------------------------|------------|
| A -- 0 to 10 in | fine sandy loam | moderately rapid | 1.57 to 1.77 in | 6.1 to 7.8 |
| AB -- 10 to 42 in | loamy fine sand | rapid | 2.58 to 3.23 in | 6.1 to 7.8 |
| C -- 42 to 60 in | fine sand | rapid | 0.35 to 1.24 in | 7.4 to 7.8 |

Map Unit Description (MN)

Redwood County, Minnesota

L244A--Du Page-Zumbro complex, 0 to 3 percent slopes, occasionally flooded

Du Page, occasionally flooded

Extent: 50 to 80 percent of the unit

Landform(s): flats on flood plains, rises on flood plains

Slope gradient: 0 to 2 percent

Parent material: alluvium

Restrictive feature(s): greater than 60 inches

Flooding: occasional

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .28

Land capability, nonirrigated 2w

Hydric soil: no

Hydrologic group: C

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap,A1,A2 -- 0 to 36 in | loam | moderate | 7.17 to 7.88 in | 6.6 to 8.4 |
| C1,C2 -- 36 to 60 in | loam | moderate | 4.08 to 4.56 in | 7.4 to 8.4 |

Havelock, occasionally flooded

Extent: 15 to 25 percent of the unit

Landform(s): flats on flood plains

Slope gradient: 0 to 2 percent

Parent material: alluvium

Restrictive feature(s): greater than 60 inches

Flooding: occasional

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .24

Land capability, nonirrigated 2w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap,A1,A2 -- 0 to 32 in | clay loam | moderate | 5.42 to 7.33 in | 7.4 to 8.4 |
| Cg -- 32 to 60 in | clay loam | moderate | 4.75 to 5.59 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

L247A--Moines-Walnut grove complex, 1 to 3 percent slopes

Moines

Extent: 35 to 60 percent of the unit

Landform(s): flats on moraines, rises on moraines

Slope gradient: 1 to 3 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .24

Land capability, nonirrigated 2s

Hydric soil: no

Hydrologic group: B/D

Potential for frost action: high

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap,ABy -- 0 to 14 in | clay loam | moderate | 2.41 to 2.69 in | 7.4 to 8.4 |
| Byg1,2,Bk -- 14 to 40 in | loam | moderate | 4.42 to 4.94 in | 7.9 to 8.4 |
| BCg -- 40 to 60 in | clay loam | moderately slow | 2.76 to 3.15 in | 7.4 to 8.4 |

Walnut grove

Extent: 20 to 40 percent of the unit

Landform(s): flats on moraines, rises on moraines

Slope gradient: 1 to 3 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .28

Land capability, nonirrigated 1

Hydric soil: no

Hydrologic group: C/D

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap,A -- 0 to 14 in | clay loam | moderately slow | 2.41 to 2.69 in | 6.6 to 7.3 |
| Bw -- 14 to 28 in | clay loam | moderate | 2.48 to 3.31 in | 6.1 to 7.3 |
| Bk -- 28 to 39 in | clay loam | moderately slow | 1.54 to 1.76 in | 7.4 to 8.4 |
| BC1,2 -- 39 to 80 in | clay loam | moderately slow | 5.73 to 6.55 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

L248B--Annton-Swanlake, firm till complex, 3 to 6 percent slopes

Annton

Extent: 35 to 65 percent of the unit

Landform(s): hills on moraines

Slope gradient: 3 to 6 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .24

Land capability, nonirrigated 2e

Hydric soil: no

Hydrologic group: C

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap -- 0 to 10 in | clay loam | moderate | 1.77 to 2.17 in | 6.1 to 7.3 |
| Bw -- 10 to 22 in | clay loam | moderate | 2.07 to 2.32 in | 6.1 to 7.3 |
| Bk -- 22 to 60 in | loam | moderately slow | 5.29 to 6.05 in | 7.4 to 8.4 |
| BC -- 60 to 80 in | clay loam | moderately slow | 2.81 to 3.21 in | 7.4 to 8.4 |

Swanlake, firm till

Extent: 30 to 50 percent of the unit

Landform(s): hills on moraines

Slope gradient: 3 to 6 percent

Parent material: till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .28

Land capability, nonirrigated 2e

Hydric soil: no

Hydrologic group: C

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|------------|
| Ap -- 0 to 8 in | clay loam | moderately slow | 1.34 to 1.50 in | 7.4 to 8.4 |
| ABk -- 8 to 18 in | clay loam | moderately slow | 1.43 to 1.64 in | 7.4 to 8.4 |
| Bk -- 18 to 40 in | clay loam | moderately slow | 3.09 to 3.53 in | 7.4 to 8.4 |
| BC -- 40 to 80 in | clay loam | moderately slow | 5.57 to 6.36 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

L250A--Lowlein loam, firm till, 1 to 3 percent slopes

Lowlein, firm till

Extent: 80 to 90 percent of the unit

Landform(s): flats on moraines, rises on moraines, flats on stream terraces, rises on stream terraces

Slope gradient: 1 to 3 percent

Parent material: outwash over till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .20

Land capability, nonirrigated 1

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

Representative soil profile:

| | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|---------------------|----------------|---------------------|---------------------------------|------------|
| Ap,AB -- 0 to 14 in | loam | moderately rapid | 1.84 to 2.13 in | 6.1 to 7.3 |
| Bw1 -- 14 to 24 in | sandy loam | moderately rapid | 1.18 to 1.38 in | 6.1 to 7.3 |
| 2Bw2 -- 24 to 31 in | loamy sand | rapid | 0.43 to 0.78 in | 6.1 to 7.3 |
| 3BC -- 31 to 80 in | clay loam | moderately slow | 6.83 to 7.81 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

L252A--Mayer clay loam, firm till substratum, 0 to 2 percent slopes

Mayer, firm till substratum

Extent: 85 to 95 percent of the unit

Landform(s): rims on depressions on moraines, flats on moraines

Slope gradient: 0 to 2 percent

Parent material: fine-loamy alluvium over sandy outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 4L

Wind erodibility index (WEI): 86

Kw factor (surface layer) .24

Land capability, nonirrigated 2w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

Representative soil profile:

| | | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|----------|----------------|----------------|---------------------|---------------------------------|------------|
| Ap,AB | -- 0 to 14 in | clay loam | moderate | 2.83 to 3.12 in | 7.4 to 8.4 |
| Bg, BCg1 | -- 14 to 36 in | clay loam | moderate | 4.33 to 4.76 in | 7.4 to 8.4 |
| BCg2 | -- 36 to 39 in | sandy loam | moderately rapid | 0.38 to 0.44 in | 7.4 to 8.4 |
| 2Cg | -- 39 to 61 in | sand | rapid | 0.44 to 0.88 in | 7.4 to 8.4 |
| 3BCg | -- 61 to 80 in | clay loam | moderately slow | 2.65 to 3.02 in | 7.4 to 8.4 |

Map Unit Description (MN)

Redwood County, Minnesota

L253B--Dickinson fine sandy loam, terrace, 1 to 6 percent slopes

Dickinson, terrace

Extent: 80 to 90 percent of the unit

Landform(s): outwash plains, terraces

Slope gradient: 1 to 6 percent

Parent material: outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .20

Land capability, nonirrigated 3e

Hydric soil: no

Hydrologic group: A

Potential for frost action: moderate

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|-----------------|---------------------|---------------------------------|------------|
| Ap,A -- 0 to 17 in | fine sandy loam | moderately rapid | 2.71 to 3.05 in | 6.1 to 7.3 |
| Bw -- 17 to 36 in | fine sandy loam | moderately rapid | 2.83 to 3.21 in | 6.1 to 7.3 |
| C -- 36 to 60 in | sand | rapid | 1.20 to 1.68 in | 6.1 to 7.3 |

M-W--Water, miscellaneous

Water, miscellaneous

Extent: 100 percent of the unit

Landform(s):

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil:

Hydrologic group:

Potential for frost action:

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|-----------|
|-------------------------------------|----------------|---------------------|---------------------------------|-----------|

Map Unit Description (MN)

Redwood County, Minnesota

W--Water

Water

Extent: 100 percent of the unit

Landform(s):

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated

Hydric soil:

Hydrologic group:

Potential for frost action:

| <i>Representative soil profile:</i> | <i>Texture</i> | <i>Permeability</i> | <i>Available water capacity</i> | <i>pH</i> |
|-------------------------------------|----------------|---------------------|---------------------------------|-----------|
|-------------------------------------|----------------|---------------------|---------------------------------|-----------|

This report provides a semitabular listing of some soil and site properties and interpretations that are valuable in communicating the concept of a map unit. The report also provides easy access to the commonly used conservation planning information in one place. The major soil components in each map unit are displayed. Minor components may be displayed if they are included in the database and are selected at the time the report is generated.