

Map Unit Description (MN)

Benton County, Minnesota

[Data apply to the entire extent of the map unit within the survey area. Map unit and soil properties for a specific parcel of land may vary somewhat and should be determined by onsite investigation]

1002A--Udorthents, wet substratum, 0 to 2 percent slopes

Udorthents, wet substratum

Extent: 100 percent of the unit

Landform(s): moraines, outwash plains, stream terraces

Slope gradient: 0 to 2 percent

Parent material: variable soil material

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated:

Hydric soil:

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

1010--Pits, quarry

Pits, quarry

Extent: 100 percent of the unit

Landform(s): stream terraces, outwash plains

Slope gradient: 0 to 50 percent

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated:

Hydric soil:

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Benton County, Minnesota

1011A--Fordum-Winterfield complex, 0 to 2 percent slopes, frequently flooded

Fordum, frequently flooded

Extent: 50 to 80 percent of the unit

Landform(s): flood plains

Slope gradient: 0 to 1 percent

Parent material: alluvium

Restrictive feature(s): greater than 60 inches

Flooding: frequent

Ponding: none

Drainage class: very poorly drained

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .28

Land capability, nonirrigated: 5w

Hydric soil: yes

Hydrologic group: A/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 7 in	fine sandy loam	moderately rapid	0.99 to 1.28 in	5.1 to 7.3
Cg -- 7 to 28 in	sandy loam	moderately rapid	2.09 to 5.01 in	5.1 to 7.3
2Cg -- 28 to 80 in	sand	rapid	1.04 to 5.20 in	5.6 to 7.3

Winterfield, frequently flooded

Extent: 20 to 40 percent of the unit

Landform(s): flood plains

Slope gradient: 0 to 2 percent

Parent material: alluvium

Restrictive feature(s): greater than 60 inches

Flooding: frequent

Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .17

Land capability, nonirrigated: 5w

Hydric soil: no

Hydrologic group: A/D

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 8 in	loamy fine sand	rapid	0.79 to 0.94 in	5.6 to 7.3
C1,C2 -- 8 to 20 in	sand	rapid	0.61 to 1.34 in	5.6 to 7.3
C3,C4,C5 -- 20 to 80 in	sand	rapid	2.39 to 5.98 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

1013A--Seelyeville and Cathro soils, ponded, 0 to 1 percent slopes

Seelyeville, ponded

<p><i>Extent:</i> 0 to 95 percent of the unit</p> <p><i>Landform(s):</i> interdrumlins, moraines</p> <p><i>Slope gradient:</i> 0 to 1 percent</p> <p><i>Parent material:</i> highly decomposed organic material</p> <p><i>Restrictive feature(s):</i> greater than 60 inches</p> <p><i>Flooding:</i> none</p> <p><i>Ponding:</i> frequent</p> <p><i>Drainage class:</i> very poorly drained</p>	<p><i>Soil loss tolerance (T factor):</i> 2</p> <p><i>Wind erodibility group (WEG):</i> 8</p> <p><i>Wind erodibility index (WEI):</i> 0</p> <p><i>Kw factor (surface layer)</i> .02</p> <p><i>Land capability, nonirrigated:</i> 8w</p> <p><i>Hydric soil:</i> yes</p> <p><i>Hydrologic group:</i> A/D</p> <p><i>Potential for frost action:</i> high</p>
-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Oa1 -- 0 to 10 in	muck	moderately rapid	3.44 to 4.43 in	
Oa2,Oa5 -- 10 to 80 in	muck	moderately rapid	24.53 to 31.54 in	

Cathro, ponded, milaca catena

<p><i>Extent:</i> 0 to 95 percent of the unit</p> <p><i>Landform(s):</i> interdrumlins, moraines</p> <p><i>Slope gradient:</i> 0 to 1 percent</p> <p><i>Parent material:</i> highly decomposed organic material over dense loamy till</p> <p><i>Restrictive feature(s):</i> densic material at 40 to 80 inches</p> <p><i>Flooding:</i> none</p> <p><i>Ponding:</i> frequent</p> <p><i>Drainage class:</i> very poorly drained</p>	<p><i>Soil loss tolerance (T factor):</i> 1</p> <p><i>Wind erodibility group (WEG):</i> 8</p> <p><i>Wind erodibility index (WEI):</i> 0</p> <p><i>Kw factor (surface layer)</i> .02</p> <p><i>Land capability, nonirrigated:</i> 8w</p> <p><i>Hydric soil:</i> yes</p> <p><i>Hydrologic group:</i> B/D</p> <p><i>Potential for frost action:</i> high</p>
-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Oa -- 0 to 31 in	muck	moderately rapid	10.89 to 14.00 in	
A -- 31 to 37 in	loam	moderate	1.18 to 1.42 in	5.6 to 6.5
BC -- 37 to 62 in	fine sandy loam	moderately rapid	2.02 to 3.02 in	5.6 to 7.3
BCd -- 62 to 80 in	fine sandy loam	moderately rapid	0.89 to 1.42 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

1020A--Bowstring and Fluvaquents, loamy, 0 to 2 percent slopes, frequently flooded

Bowstring, frequently flooded

Extent: 25 to 75 percent of the unit

Landform(s): flood plains

Slope gradient: 0 to 1 percent

Parent material: highly decomposed organic materials containing strata of alluvium

Restrictive feature(s): greater than 60 inches

Flooding: frequent

Ponding: none

Drainage class: very poorly drained

Soil loss tolerance (T factor): 1

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .02

Land capability, nonirrigated: 7w

Hydric soil: yes

Hydrologic group: A/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Oa -- 0 to 38 in	muck	moderately rapid	13.37 to 17.19 in	
Cg -- 38 to 47 in	stratified fine sand to loamy fine sand	rapid	0.43 to 0.87 in	5.6 to 7.3
Oa' -- 47 to 80 in	muck	moderately rapid	11.57 to 14.88 in	

Fluvaquents, frequently flooded, loamy

Extent: 20 to 60 percent of the unit

Landform(s): flood plains

Slope gradient: 0 to 1 percent

Parent material: alluvium

Restrictive feature(s): greater than 60 inches

Flooding: frequent

Ponding: none

Drainage class: very poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .32

Land capability, nonirrigated: 7w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 6 in	mucky silt loam	moderate	1.18 to 1.42 in	5.6 to 7.3
Cg -- 6 to 80 in	stratified silt loam to gravelly loamy coarse sand	rapid	4.44 to 16.28 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

1023A--Seelyeville and Markey soils, ponded, 0 to 1 percent slopes

Seelyeville, ponded

Extent: 0 to 100 percent of the unit

Landform(s): lake plains, outwash plains, stream terraces

Slope gradient: 0 to 1 percent

Parent material: highly decomposed organic material

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: frequent

Drainage class: very poorly drained

Soil loss tolerance (T factor): 2

Wind erodibility group (WEG): 8

Wind erodibility index (WEI): 0

Kw factor (surface layer) .02

Land capability, nonirrigated: 8w

Hydric soil: yes

Hydrologic group: A/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Oa1 -- 0 to 15 in	muck	moderately rapid	5.24 to 6.73 in	
Oa2,Oa5 -- 15 to 80 in	muck	moderately rapid	22.74 to 29.23 in	

Markey, ponded

Extent: 0 to 100 percent of the unit

Landform(s): lake plains, outwash plains, stream terraces

Slope gradient: 0 to 1 percent

Parent material: highly decomposed organic material over outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: frequent

Drainage class: very poorly drained

Soil loss tolerance (T factor): 1

Wind erodibility group (WEG): 8

Wind erodibility index (WEI): 0

Kw factor (surface layer) .02

Land capability, nonirrigated: 8w

Hydric soil: yes

Hydrologic group: A/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Oa -- 0 to 27 in	muck	moderately rapid	9.51 to 12.22 in	
A -- 27 to 32 in	loamy sand	rapid	0.28 to 0.57 in	5.6 to 7.3
Cg -- 32 to 80 in	sand	rapid	1.92 to 3.36 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

1025A--Fluvaquents and Udifluvents, loamy, 0 to 2 percent slopes, frequently flooded

Fluvaquents, frequently flooded, loamy

Extent: 25 to 80 percent of the unit

Landform(s): flood plains

Slope gradient: 0 to 2 percent

Parent material: alluvium

Restrictive feature(s): greater than 60 inches

Flooding: frequent

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .32

Land capability, nonirrigated: 5w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 6 in	silt loam	moderate	1.18 to 1.42 in	5.6 to 7.3
Cg -- 6 to 80 in	stratified silt loam to gravelly loamy coarse sand	rapid	4.44 to 16.28 in	5.6 to 7.3

Udifluvents, frequently flooded, loamy

Extent: 10 to 50 percent of the unit

Landform(s): flood plains

Slope gradient: 0 to 2 percent

Parent material: alluvium

Restrictive feature(s): greater than 60 inches

Flooding: frequent

Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .32

Land capability, nonirrigated: 4w

Hydric soil: no

Hydrologic group: B/D

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 6 in	silt loam	moderate	1.18 to 1.42 in	5.6 to 7.3
Cg -- 6 to 80 in	stratified silt loam to gravelly loamy coarse sand	rapid	4.44 to 16.28 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

1052B--Udorthents, loamy-Rock outcrop complex, 1 to 6 percent slopes

Udorthents, loamy

Extent: 40 to 90 percent of the unit

Landform(s): drumlins

Slope gradient: 1 to 6 percent

Parent material: loamy drift over bedrock

Restrictive feature(s): lithic bedrock at 20 to 80 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 2

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .15

Land capability, nonirrigated: 6s

Hydric soil: no

Hydrologic group: B

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 4 in	fine sandy loam	moderately rapid	0.55 to 0.71 in	5.1 to 6.5
C -- 4 to 22 in	fine sandy loam	moderately rapid	1.99 to 3.44 in	5.1 to 6.5
2R -- 22 to 80 in	unweathered bedrock	impermeable		

Rock outcrop

Extent: 10 to 60 percent of the unit

Landform(s): drumlins

Slope gradient: 1 to 6 percent

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated:

Hydric soil:

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>

Map Unit Description (MN)

Benton County, Minnesota

C2A--Adolph mucky silty clay loam, depressional, 0 to 1 percent slopes, stony

Adolph, depressional, stony

Extent: 80 to 95 percent of the unit

Landform(s): drumlins, moraines

Slope gradient: 0 to 1 percent

Parent material: loamy mantled dense loamy till

Restrictive feature(s): densic material at 40 to 80 inches

Flooding: none

Ponding: frequent

Drainage class: very poorly drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .32

Land capability, nonirrigated: 6w

Hydric soil: yes

Hydrologic group: C/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A1 -- 0 to 5 in	mucky silty clay loam	moderately slow	1.07 to 1.18 in	5.1 to 6.5
A2 -- 5 to 13 in	silt loam	moderate	1.65 to 1.89 in	5.1 to 6.5
Bg -- 13 to 32 in	silt loam	moderate	3.21 to 4.16 in	5.1 to 7.3
2BC -- 32 to 44 in	fine sandy loam	slow	0.98 to 1.46 in	5.6 to 7.3
2BCd -- 44 to 80 in	fine sandy loam	very slow	1.79 to 2.87 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

C4A--Cebana silt loam, 0 to 1 percent slopes, stony

Cebana, stony

Extent: 85 to 95 percent of the unit

Landform(s): drumlins, moraines

Slope gradient: 0 to 1 percent

Parent material: loamy mantled dense loamy till

Restrictive feature(s): densic material at 40 to 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .37

Land capability, nonirrigated: 4w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 8 in	silt loam	moderate	1.50 to 1.89 in	5.1 to 6.5
Eg,E/B -- 8 to 27 in	silt loam	moderate	3.28 to 4.24 in	5.1 to 6.5
2Bt -- 27 to 49 in	loam	moderate	1.95 to 3.90 in	5.1 to 7.3
2BCd -- 49 to 80 in	fine sandy loam	very slow	1.56 to 2.49 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

C5C--Milaca fine sandy loam, 8 to 15 percent slopes, stony

Milaca, stony

Extent: 70 to 90 percent of the unit

Landform(s): drumlins, moraines

Slope gradient: 8 to 15 percent

Parent material: dense loamy till

Restrictive feature(s): densic material at 40 to 60 inches

Flooding: none

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .15

Land capability, nonirrigated: 3e

Hydric soil: no

Hydrologic group: A/D

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 4 in	fine sandy loam	moderately rapid	0.55 to 0.71 in	5.1 to 6.5
E -- 4 to 15 in	fine sandy loam	moderately rapid	1.21 to 2.09 in	5.1 to 6.5
E/B,Bt1,Bt2 -- 15 to 42 in	fine sandy loam	moderately rapid	2.72 to 4.07 in	5.1 to 6.5
BCd -- 42 to 80 in	fine sandy loam	very slow	1.89 to 3.02 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

C5E--Milaca fine sandy loam, 15 to 30 percent slopes, stony

Milaca, stony

Extent: 70 to 90 percent of the unit

Landform(s): drumlins, moraines

Slope gradient: 15 to 30 percent

Parent material: dense loamy till

Restrictive feature(s): densic material at 40 to 60 inches

Flooding: none

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .15

Land capability, nonirrigated: 6e

Hydric soil: no

Hydrologic group: A/D

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 4 in	fine sandy loam	moderately rapid	0.55 to 0.71 in	5.1 to 6.5
E -- 4 to 15 in	fine sandy loam	moderately rapid	1.21 to 2.09 in	5.1 to 6.5
E/B,Bt1,Bt2 -- 15 to 42 in	fine sandy loam	moderately rapid	2.72 to 4.07 in	5.1 to 6.5
BCd -- 42 to 80 in	fine sandy loam	very slow	1.89 to 3.02 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

C8A--Brennyville, wet-Cebana complex, 0 to 3 percent slopes, stony

Brennyville, wet, stony

Extent: 68 to 98 percent of the unit

Landform(s): drumlins, moraines

Slope gradient: 1 to 3 percent

Parent material: silt mantled dense loamy till

Restrictive feature(s): densic material at 40 to 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .43

Land capability, nonirrigated: 3w

Hydric soil: no

Hydrologic group: C/D

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 8 in	silt loam	moderate	1.50 to 1.89 in	5.1 to 6.5
B/E -- 8 to 11 in	silt loam	moderate	0.63 to 0.69 in	5.1 to 6.5
Bt1 -- 11 to 21 in	silt loam	moderate	1.67 to 2.17 in	5.1 to 6.5
2Bt2,2Bt3 -- 21 to 38 in	fine sandy loam	moderately rapid	1.56 to 3.12 in	5.1 to 7.3
2BC -- 38 to 45 in	fine sandy loam	slow	0.54 to 0.80 in	5.6 to 7.3
2BCd -- 45 to 80 in	fine sandy loam	very slow	1.75 to 2.80 in	5.6 to 7.3

Cebana, stony

Extent: 5 to 30 percent of the unit

Landform(s): drumlins, moraines

Slope gradient: 0 to 2 percent

Parent material: loamy mantled dense loamy till

Restrictive feature(s): densic material at 40 to 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .37

Land capability, nonirrigated: 4w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 8 in	silt loam	moderate	1.50 to 1.89 in	5.1 to 6.5
Eg,E/B -- 8 to 27 in	silt loam	moderate	3.28 to 4.24 in	5.1 to 6.5
2Bt -- 27 to 49 in	loam	moderate	1.95 to 3.90 in	5.1 to 7.3
2BCd -- 49 to 80 in	fine sandy loam	very slow	1.56 to 2.49 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

C10B--Brennyville complex, 1 to 6 percent slopes, stony

Brennyville, stony

Extent: 40 to 85 percent of the unit

Landform(s): drumlins, moraines

Slope gradient: 3 to 6 percent

Parent material: silt mantled dense loamy till

Restrictive feature(s): densic material at 40 to 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .43

Land capability, nonirrigated: 3s

Hydric soil: no

Hydrologic group: C/D

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 8 in	silt loam	moderate	1.50 to 1.89 in	5.1 to 6.5
B/E -- 8 to 11 in	silt loam	moderate	0.63 to 0.69 in	5.1 to 6.5
Bt1 -- 11 to 21 in	silt loam	moderate	1.67 to 2.17 in	5.1 to 6.5
2Bt2,2Bt3 -- 21 to 38 in	fine sandy loam	moderately rapid	1.56 to 3.12 in	5.1 to 7.3
2BC -- 38 to 45 in	fine sandy loam	slow	0.54 to 0.80 in	5.6 to 7.3
2BCd -- 45 to 80 in	fine sandy loam	very slow	1.75 to 2.80 in	5.6 to 7.3

Brennyville, wet, stony

Extent: 5 to 35 percent of the unit

Landform(s): drumlins, moraines

Slope gradient: 1 to 3 percent

Parent material: silt mantled dense loamy till

Restrictive feature(s): densic material at 40 to 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .43

Land capability, nonirrigated: 3w

Hydric soil: no

Hydrologic group: C/D

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 8 in	silt loam	moderate	1.50 to 1.89 in	5.1 to 6.5
B/E -- 8 to 11 in	silt loam	moderate	0.63 to 0.69 in	5.1 to 6.5
Bt1 -- 11 to 21 in	silt loam	moderate	1.67 to 2.17 in	5.1 to 6.5
2Bt2,2Bt3 -- 21 to 38 in	fine sandy loam	moderately rapid	1.56 to 3.12 in	5.1 to 7.3
2BC -- 38 to 45 in	fine sandy loam	slow	0.54 to 0.80 in	5.6 to 7.3
2BCd -- 45 to 80 in	fine sandy loam	very slow	1.75 to 2.80 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

C10B--Brennyville complex, 1 to 6 percent slopes, stony

Map Unit Description (MN)

Benton County, Minnesota

C14A--Cathro, Twig, and Adolph soils, ponded, 0 to 1 percent slopes

Adolph, ponded, stony

Extent: 0 to 100 percent of the unit

Landform(s): interdrumlins, moraines

Slope gradient: 0 to 1 percent

Parent material: loamy mantled dense loamy till

Restrictive feature(s): densic material at 40 to 80 inches

Flooding: none

Ponding: frequent

Drainage class: very poorly drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 8

Wind erodibility index (WEI): 0

Kw factor (surface layer) .32

Land capability, nonirrigated: 8w

Hydric soil: yes

Hydrologic group: C/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A1 -- 0 to 5 in	mucky silty clay loam	moderately slow	1.07 to 1.18 in	5.1 to 6.5
A2 -- 5 to 13 in	silt loam	moderate	1.65 to 1.89 in	5.1 to 6.5
Bg -- 13 to 32 in	silt loam	moderate	3.21 to 4.16 in	5.1 to 7.3
2BC -- 32 to 44 in	fine sandy loam	slow	0.98 to 1.46 in	5.6 to 7.3
2BCd -- 44 to 80 in	fine sandy loam	very slow	1.79 to 2.87 in	5.6 to 7.3

Cathro, ponded, milaca catena

Extent: 0 to 100 percent of the unit

Landform(s): interdrumlins, moraines

Slope gradient: 0 to 1 percent

Parent material: highly decomposed organic material over dense loamy till

Restrictive feature(s): densic material at 40 to 80 inches

Flooding: none

Ponding: frequent

Drainage class: very poorly drained

Soil loss tolerance (T factor): 1

Wind erodibility group (WEG): 8

Wind erodibility index (WEI): 0

Kw factor (surface layer) .02

Land capability, nonirrigated: 8w

Hydric soil: yes

Hydrologic group: C/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Oa -- 0 to 31 in	muck	moderately rapid	10.89 to 14.00 in	
A -- 31 to 37 in	loam	moderate	1.18 to 1.42 in	5.6 to 6.5
BC -- 37 to 62 in	fine sandy loam	slow	2.02 to 3.02 in	5.6 to 7.3
BCd -- 62 to 80 in	fine sandy loam	very slow	0.89 to 1.42 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

C14A--Cathro, Twig, and Adolph soils, ponded, 0 to 1 percent slopes

Twig, ponded

<p><i>Extent:</i> 0 to 100 percent of the unit</p> <p><i>Landform(s):</i> interdrumlins, moraines</p> <p><i>Slope gradient:</i> 0 to 1 percent</p> <p><i>Parent material:</i> thin highly decomposed organic material over dense loamy till</p> <p><i>Restrictive feature(s):</i> densic material at 40 to 80 inches</p> <p><i>Flooding:</i> none</p> <p><i>Ponding:</i> frequent</p> <p><i>Drainage class:</i> very poorly drained</p>	<p><i>Soil loss tolerance (T factor):</i> 1</p> <p><i>Wind erodibility group (WEG):</i> 8</p> <p><i>Wind erodibility index (WEI):</i> 0</p> <p><i>Kw factor (surface layer)</i> .02</p> <p><i>Land capability, nonirrigated:</i> 8w</p> <p><i>Hydric soil:</i> yes</p> <p><i>Hydrologic group:</i> B/D</p> <p><i>Potential for frost action:</i> high</p>
-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Oa -- 0 to 10 in	muck	moderately rapid	3.44 to 4.43 in	3.5 to 6.0
A -- 10 to 16 in	loam	moderate	1.07 to 1.51 in	3.5 to 6.0
Eg -- 16 to 21 in	fine sandy loam	moderately rapid	0.61 to 1.04 in	3.5 to 6.0
2Bg,2Bw -- 21 to 44 in	sandy loam	moderately rapid	2.09 to 4.41 in	3.5 to 6.0
2BC -- 44 to 52 in	fine sandy loam	slow	0.63 to 0.94 in	5.6 to 7.3
2BCd -- 52 to 80 in	fine sandy loam	very slow	1.40 to 2.24 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

C23B--Antigo-Chetek complex, 2 to 8 percent slopes

Antigo

Extent: 45 to 80 percent of the unit

Landform(s): outwash plains, stream terraces

Slope gradient: 2 to 6 percent

Parent material: outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .49

Land capability, nonirrigated: 3s

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 4 in	silt loam	moderate	0.75 to 0.87 in	5.1 to 6.5
E -- 4 to 12 in	silt loam	moderate	1.50 to 1.73 in	5.1 to 6.5
B/E -- 12 to 18 in	silt loam	moderate	1.20 to 1.39 in	5.1 to 6.5
2Bt -- 18 to 30 in	fine sandy loam	moderate	1.06 to 2.13 in	5.1 to 6.5
3C -- 30 to 80 in	gravelly coarse sand	very rapid	0.50 to 3.00 in	5.1 to 6.5

Chetek

Extent: 15 to 40 percent of the unit

Landform(s): outwash plains, stream terraces

Slope gradient: 2 to 8 percent

Parent material: outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat excessively drained

Soil loss tolerance (T factor): 2

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .28

Land capability, nonirrigated: 6s

Hydric soil: no

Hydrologic group: A

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 5 in	fine sandy loam	moderately rapid	0.72 to 0.92 in	5.1 to 6.0
E -- 5 to 12 in	fine sandy loam	moderately rapid	0.67 to 1.14 in	5.1 to 6.0
Bt1 -- 12 to 18 in	sandy loam	moderately rapid	0.57 to 1.20 in	5.1 to 6.0
2Bt2 -- 18 to 25 in	gravelly loamy coarse sand	rapid	0.07 to 0.64 in	5.1 to 6.5
2BC,2C -- 25 to 80 in	gravelly coarse sand	very rapid	0.55 to 3.28 in	5.1 to 6.5

Map Unit Description (MN)

Benton County, Minnesota

C26A--Foglake silt loam, 0 to 2 percent slopes

Foglake

Extent: 75 to 90 percent of the unit

Landform(s): lake plains

Slope gradient: 0 to 2 percent

Parent material: silty glaciolacustrine

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 6

Wind erodibility index (WEI): 48

Kw factor (surface layer) .37

Land capability, nonirrigated: 3w

Hydric soil: yes

Hydrologic group: C/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 8 in	silt loam	moderate	1.73 to 1.89 in	5.1 to 7.3
Eg -- 8 to 16 in	silt loam	moderate	1.41 to 1.82 in	5.1 to 7.3
Btg,BCg -- 16 to 47 in	silty clay loam	slow	2.46 to 6.14 in	5.1 to 7.3
Cg -- 47 to 80 in	silty clay loam	moderately slow	5.95 to 7.28 in	7.4 to 8.4

Map Unit Description (MN)

Benton County, Minnesota

C28A--Cathro and Twig soils, depressional, 0 to 1 percent slopes

Cathro, depressional, milaca catena

<i>Extent:</i> 0 to 95 percent of the unit	<i>Soil loss tolerance (T factor):</i> 1
<i>Landform(s):</i> interdrumlins, moraines	<i>Wind erodibility group (WEG):</i> 8
<i>Slope gradient:</i> 0 to 1 percent	<i>Wind erodibility index (WEI):</i> 0
<i>Parent material:</i> highly decomposed organic material over dense loamy till	<i>Kw factor (surface layer)</i> .02
<i>Restrictive feature(s):</i> densic material at 40 to 80 inches	<i>Land capability, nonirrigated:</i> 7w
<i>Flooding:</i> none	<i>Hydric soil:</i> yes
<i>Ponding:</i> frequent	<i>Hydrologic group:</i> C/D
<i>Drainage class:</i> very poorly drained	<i>Potential for frost action:</i> high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Oa -- 0 to 31 in	muck	moderately rapid	10.89 to 14.00 in	
A -- 31 to 37 in	loam	moderate	1.18 to 1.42 in	5.6 to 6.5
BC -- 37 to 62 in	fine sandy loam	slow	2.02 to 3.02 in	5.6 to 7.3
BCd -- 62 to 80 in	fine sandy loam	very slow	0.89 to 1.42 in	5.6 to 7.3

Twig, depressional

<i>Extent:</i> 0 to 95 percent of the unit	<i>Soil loss tolerance (T factor):</i> 1
<i>Landform(s):</i> interdrumlins, moraines	<i>Wind erodibility group (WEG):</i> 2
<i>Slope gradient:</i> 0 to 1 percent	<i>Wind erodibility index (WEI):</i> 134
<i>Parent material:</i> thin highly decomposed organic material over dense loamy till	<i>Kw factor (surface layer)</i> .02
<i>Restrictive feature(s):</i> densic material at 40 to 80 inches	<i>Land capability, nonirrigated:</i> 6w
<i>Flooding:</i> none	<i>Hydric soil:</i> yes
<i>Ponding:</i> frequent	<i>Hydrologic group:</i> B/D
<i>Drainage class:</i> very poorly drained	<i>Potential for frost action:</i> high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Oa -- 0 to 10 in	muck	moderately rapid	3.44 to 4.43 in	3.5 to 6.0
A -- 10 to 16 in	loam	moderate	1.07 to 1.51 in	3.5 to 6.0
Eg -- 16 to 21 in	fine sandy loam	moderately rapid	0.61 to 1.04 in	3.5 to 6.0
2Bg,2Bw -- 21 to 44 in	sandy loam	moderately rapid	2.09 to 4.41 in	3.5 to 6.0
2BC -- 44 to 52 in	fine sandy loam	slow	0.63 to 0.94 in	5.6 to 7.3
2BCd -- 52 to 80 in	fine sandy loam	very slow	1.40 to 2.24 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

C28A--Cathro and Twig soils, depressional, 0 to 1 percent slopes

C36A--Nokasippi loamy fine sand, depressional, 0 to 1 percent slopes

Nokasippi, depressional

Extent: 75 to 95 percent of the unit

Landform(s): drainageways on interdrumlins, depressions on interdrumlins

Slope gradient: 0 to 1 percent

Parent material: sandy outwash over dense loamy till

Restrictive feature(s): densic material at 40 to 60 inches

Flooding: none

Ponding: frequent

Drainage class: very poorly drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .15

Land capability, nonirrigated: 6w

Hydric soil: yes

Hydrologic group: C/D

Potential for frost action: high

<i>Representative soil profile:</i>		<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A --	0 to 10 in	loamy fine sand	rapid	0.98 to 1.18 in	4.5 to 6.5
Bg1 --	10 to 21 in	loamy sand	rapid	0.66 to 1.32 in	4.5 to 7.3
Bg2 --	21 to 32 in	loamy sand	moderately rapid	1.21 to 1.98 in	4.5 to 7.3
2Bg3 --	32 to 41 in	sandy loam	slow	0.00 to 0.72 in	4.5 to 7.3
2BCd --	41 to 80 in	fine sandy loam	very slow	1.95 to 3.12 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

C37C--Braham loamy fine sand, 6 to 12 percent slopes

Braham

Extent: 75 to 95 percent of the unit

Landform(s): drumlins, moraines

Slope gradient: 6 to 12 percent

Parent material: sandy outwash over till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .28

Land capability, nonirrigated: 4e

Hydric soil: no

Hydrologic group: B

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 9 in	loamy fine sand	rapid	0.91 to 1.09 in	5.6 to 7.3
E -- 9 to 28 in	loamy fine sand	rapid	1.51 to 1.89 in	5.6 to 7.3
2Bt -- 28 to 39 in	sandy clay loam	moderate	1.65 to 1.98 in	5.1 to 7.3
2C -- 39 to 60 in	loam	moderate	3.13 to 3.76 in	7.4 to 8.4

Map Unit Description (MN)

Benton County, Minnesota

C38B--Grasston fine sandy loam, terrace, 2 to 6 percent slopes

Grasston, terrace

Extent: 55 to 85 percent of the unit

Landform(s): lake plains, terraces

Slope gradient: 2 to 6 percent

Parent material: silty and clayey glaciolacustrine deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .24

Land capability, nonirrigated: 2s

Hydric soil: no

Hydrologic group: D

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 9 in	fine sandy loam	moderately rapid	1.27 to 1.63 in	5.1 to 6.5
B/E -- 9 to 14 in	silty clay loam	moderately slow	0.56 to 1.13 in	5.1 to 7.3
Bt1,Bt2 -- 14 to 42 in	silty clay	slow	2.24 to 5.59 in	5.1 to 7.3
Bk -- 42 to 66 in	silt loam	moderate	4.32 to 5.28 in	7.4 to 8.4
C -- 66 to 80 in	silt loam	moderate	2.34 to 3.03 in	7.4 to 8.4

Map Unit Description (MN)

Benton County, Minnesota

C39A--Foglake fine sandy loam, terrace, 0 to 2 percent slopes

Foglake, terrace

Extent: 50 to 75 percent of the unit

Landform(s): lake plains, terraces

Slope gradient: 0 to 2 percent

Parent material: silty and clayey glaciolacustrine deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .24

Land capability, nonirrigated: 3w

Hydric soil: yes

Hydrologic group: C/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 8 in	fine sandy loam	moderately rapid	1.10 to 1.42 in	5.1 to 6.5
Eg -- 8 to 16 in	loam	moderate	1.41 to 1.82 in	5.1 to 7.3
Btg,BCg -- 16 to 47 in	silty clay loam	slow	2.46 to 6.14 in	5.1 to 7.3
Cg -- 47 to 80 in	silty clay loam	moderately slow	5.95 to 7.28 in	7.4 to 8.4

Map Unit Description (MN)

Benton County, Minnesota

C42B--Sartell, till substratum-Bushville complex, 1 to 6 percent slopes

Sartell, till substratum

Extent: 65 to 95 percent of the unit

Landform(s): drumlins

Slope gradient: 1 to 6 percent

Parent material: eolian sands over till

Restrictive feature(s): densic material at 60 to 80 inches

Flooding: none

Ponding: none

Drainage class: excessively drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 1

Wind erodibility index (WEI): 250

Kw factor (surface layer) .10

Land capability, nonirrigated: 4s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 4 in	fine sand	rapid	0.35 to 0.43 in	5.1 to 6.0
Bw -- 4 to 33 in	fine sand	rapid	1.75 to 2.91 in	5.1 to 6.0
C -- 33 to 66 in	fine sand	rapid	1.65 to 2.98 in	5.6 to 7.3
2BCd -- 66 to 80 in	fine sandy loam	very slow	0.69 to 1.10 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

C46A--Foglake loam, terrace, 0 to 2 percent slopes

Foglake, terrace

Extent: 50 to 75 percent of the unit

Landform(s): lake plains, terraces

Slope gradient: 0 to 2 percent

Parent material: silty and clayey glaciolacustrine deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .37

Land capability, nonirrigated: 3w

Hydric soil: yes

Hydrologic group: C/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 8 in	loam	moderate	1.42 to 1.81 in	5.1 to 6.5
Eg -- 8 to 16 in	loam	moderate	1.41 to 1.82 in	5.1 to 7.3
Btg,BCg -- 16 to 47 in	silty clay loam	slow	2.46 to 6.14 in	5.1 to 7.3
Cg -- 47 to 80 in	silty clay loam	moderately slow	5.95 to 7.28 in	7.4 to 8.4

Map Unit Description (MN)

Benton County, Minnesota

C48A--Ronneby loam, 0 to 2 percent slopes, stony

Ronneby, stony

Extent: 75 to 95 percent of the unit

Landform(s): interdrumlins

Slope gradient: 0 to 2 percent

Parent material: dense till

Restrictive feature(s): densic material at 40 to 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .32

Land capability, nonirrigated: 2w

Hydric soil: no

Hydrologic group: C/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 4 in	loam	moderate	0.71 to 0.91 in	5.1 to 6.5
E -- 4 to 12 in	fine sandy loam	moderately rapid	0.94 to 1.50 in	5.1 to 6.5
Bt1,Bt2 -- 12 to 33 in	fine sandy loam	moderate	2.55 to 4.04 in	5.6 to 6.5
Bt3 -- 33 to 45 in	fine sandy loam	moderately slow	0.94 to 1.42 in	5.6 to 7.3
BCd -- 45 to 80 in	fine sandy loam	very slow	1.75 to 2.80 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

C49A--Antigo silt loam, 0 to 2 percent slopes

Antigo

Extent: 55 to 93 percent of the unit

Landform(s): outwash plains, stream terraces

Slope gradient: 0 to 2 percent

Parent material: outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .49

Land capability, nonirrigated: 3s

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 4 in	silt loam	moderate	0.75 to 0.87 in	5.1 to 6.5
E -- 4 to 12 in	silt loam	moderate	1.50 to 1.73 in	5.1 to 6.5
B/E -- 12 to 18 in	silt loam	moderate	1.20 to 1.39 in	5.1 to 6.5
2Bt -- 18 to 30 in	fine sandy loam	moderate	1.06 to 2.13 in	5.1 to 6.5
3C -- 30 to 80 in	gravelly coarse sand	very rapid	0.50 to 3.00 in	5.1 to 6.5

Map Unit Description (MN)

Benton County, Minnesota

C51D--Emmert-St. Francis complex, 6 to 25 percent slopes

Emmert

Extent: 50 to 80 percent of the unit

Landform(s): outwash plains, terraces

Slope gradient: 6 to 25 percent

Parent material: gravelly outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: excessively drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .17

Land capability, nonirrigated: 6s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 2 in	fine sandy loam	rapid	0.16 to 0.24 in	5.1 to 6.5
E -- 2 to 12 in	gravelly loamy sand	rapid	0.49 to 1.48 in	5.1 to 6.5
Bt -- 12 to 37 in	gravelly loamy coarse sand	rapid	0.76 to 2.27 in	5.1 to 6.5
C -- 37 to 80 in	very gravelly coarse sand	very rapid	0.43 to 2.57 in	5.1 to 6.5

St. francis

Extent: 20 to 50 percent of the unit

Landform(s): terraces, outwash plains

Slope gradient: 3 to 8 percent

Parent material: loamy glaciofluvial deposits over sandy and gravelly outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat excessively drained

Soil loss tolerance (T factor): 2

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .32

Land capability, nonirrigated: 3e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 6 in	fine sandy loam	moderately rapid	0.83 to 1.06 in	5.1 to 6.5
E -- 6 to 11 in	fine sandy loam	moderately rapid	0.51 to 0.87 in	5.1 to 6.5
Bt1 -- 11 to 16 in	loam	moderate	0.46 to 0.97 in	5.1 to 6.5
2Bt2 -- 16 to 20 in	gravelly loamy coarse sand	rapid	0.04 to 0.35 in	5.1 to 6.5
2C -- 20 to 80 in	gravelly coarse sand	very rapid	0.60 to 3.59 in	5.1 to 6.5

Map Unit Description (MN)

Benton County, Minnesota

C52B--Waukon fine sandy loam, dense substratum, 2 to 6 percent slopes

Waukon, dense substratum

Extent: 65 to 90 percent of the unit

Landform(s): moraines

Slope gradient: 2 to 6 percent

Parent material: loamy till over dense loamy till

Restrictive feature(s): densic material at 60 to 80 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .24

Land capability, nonirrigated: 2e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 8 in	fine sandy loam	moderately rapid	1.02 to 1.18 in	6.1 to 7.3
E -- 8 to 12 in	fine sandy loam	moderately rapid	0.43 to 0.75 in	5.6 to 7.3
BE,Bt -- 12 to 43 in	loam	moderate	4.67 to 5.91 in	6.1 to 8.4
Bk -- 43 to 64 in	loam	moderate	3.19 to 4.04 in	7.4 to 8.4
2BCd -- 64 to 80 in	fine sandy loam	very slow	0.79 to 1.26 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

C53C--Pomroy loamy fine sand, 6 to 12 percent slopes

Pomroy

Extent: 75 to 95 percent of the unit

Landform(s): drumlins, moraines

Slope gradient: 6 to 12 percent

Parent material: sandy mantled dense loamy till

Restrictive feature(s): densic material at 40 to 60 inches

Flooding: none

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .28

Land capability, nonirrigated: 3e

Hydric soil: no

Hydrologic group: C/D

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 6 in	loamy fine sand	rapid	0.47 to 0.71 in	5.1 to 6.5
E -- 6 to 22 in	loamy fine sand	rapid	0.81 to 1.78 in	5.1 to 6.5
2Bt -- 22 to 31 in	sandy loam	moderately rapid	0.91 to 1.45 in	5.1 to 6.5
2BC -- 31 to 41 in	sandy loam	slow	0.79 to 1.18 in	5.6 to 7.3
2BCd -- 41 to 60 in	fine sandy loam	very slow	0.94 to 1.51 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

C54B--Chetek-Mahtomedi complex, 2 to 6 percent slopes

Chetek

<i>Extent:</i> 45 to 70 percent of the unit	<i>Soil loss tolerance (T factor):</i> 2
<i>Landform(s):</i> outwash plains, stream terraces, moraines	<i>Wind erodibility group (WEG):</i> 3
<i>Slope gradient:</i> 2 to 6 percent	<i>Wind erodibility index (WEI):</i> 86
<i>Parent material:</i> loamy glaciofluvial deposits and/or sandy and gravelly outwash	<i>Kw factor (surface layer)</i> .28
<i>Restrictive feature(s):</i> greater than 60 inches	<i>Land capability, nonirrigated:</i> 4s
<i>Flooding:</i> none	<i>Hydric soil:</i> no
<i>Ponding:</i> none	<i>Hydrologic group:</i> A
<i>Drainage class:</i> somewhat excessively drained	<i>Potential for frost action:</i> moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 5 in	fine sandy loam	moderately rapid	0.72 to 0.92 in	5.1 to 6.0
E -- 5 to 12 in	fine sandy loam	moderately rapid	0.67 to 1.14 in	5.1 to 6.0
Bt1 -- 12 to 18 in	sandy loam	moderately rapid	0.57 to 1.20 in	5.1 to 6.0
2Bt2 -- 18 to 25 in	gravelly loamy coarse sand	rapid	0.07 to 0.64 in	5.1 to 6.5
2BC,2C -- 25 to 80 in	gravelly coarse sand	very rapid	0.55 to 3.28 in	5.1 to 6.5

Mahtomedi

<i>Extent:</i> 10 to 40 percent of the unit	<i>Soil loss tolerance (T factor):</i> 5
<i>Landform(s):</i> outwash plains, stream terraces, moraines	<i>Wind erodibility group (WEG):</i> 3
<i>Slope gradient:</i> 2 to 6 percent	<i>Wind erodibility index (WEI):</i> 86
<i>Parent material:</i> sandy outwash	<i>Kw factor (surface layer)</i> .24
<i>Restrictive feature(s):</i> greater than 60 inches	<i>Land capability, nonirrigated:</i> 4s
<i>Flooding:</i> none	<i>Hydric soil:</i> no
<i>Ponding:</i> none	<i>Hydrologic group:</i> A
<i>Drainage class:</i> excessively drained	<i>Potential for frost action:</i> low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 4 in	fine sandy loam	moderately rapid	0.55 to 0.71 in	5.1 to 6.5
E -- 4 to 9 in	gravelly coarse sandy loam	moderately rapid	0.20 to 0.77 in	5.1 to 6.5
Bw -- 9 to 20 in	gravelly coarse sand	very rapid	0.11 to 0.66 in	5.1 to 6.5
BC,C -- 20 to 80 in	gravelly coarse sand	very rapid	0.60 to 3.59 in	5.1 to 6.5

Map Unit Description (MN)

Benton County, Minnesota

C54C--Chetek-Mahtomedi complex, 6 to 12 percent slopes

Chetek

<i>Extent:</i> 45 to 75 percent of the unit	<i>Soil loss tolerance (T factor):</i> 2
<i>Landform(s):</i> outwash plains, stream terraces, moraines	<i>Wind erodibility group (WEG):</i> 3
<i>Slope gradient:</i> 6 to 12 percent	<i>Wind erodibility index (WEI):</i> 86
<i>Parent material:</i> loamy glaciofluvial deposits and/or sandy and gravelly outwash	<i>Kw factor (surface layer)</i> .28
<i>Restrictive feature(s):</i> greater than 60 inches	<i>Land capability, nonirrigated:</i> 6s
<i>Flooding:</i> none	<i>Hydric soil:</i> no
<i>Ponding:</i> none	<i>Hydrologic group:</i> A
<i>Drainage class:</i> somewhat excessively drained	<i>Potential for frost action:</i> moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 5 in	fine sandy loam	moderately rapid	0.72 to 0.92 in	5.1 to 6.0
E -- 5 to 12 in	fine sandy loam	moderately rapid	0.67 to 1.14 in	5.1 to 6.0
Bt1 -- 12 to 18 in	sandy loam	moderately rapid	0.57 to 1.20 in	5.1 to 6.0
2Bt2 -- 18 to 25 in	gravelly loamy coarse sand	rapid	0.07 to 0.64 in	5.1 to 6.5
2BC,2C -- 25 to 80 in	gravelly coarse sand	very rapid	0.55 to 3.28 in	5.1 to 6.5

Mahtomedi

<i>Extent:</i> 10 to 40 percent of the unit	<i>Soil loss tolerance (T factor):</i> 5
<i>Landform(s):</i> outwash plains, stream terraces, moraines	<i>Wind erodibility group (WEG):</i> 3
<i>Slope gradient:</i> 6 to 12 percent	<i>Wind erodibility index (WEI):</i> 86
<i>Parent material:</i> sandy outwash	<i>Kw factor (surface layer)</i> .24
<i>Restrictive feature(s):</i> greater than 60 inches	<i>Land capability, nonirrigated:</i> 6s
<i>Flooding:</i> none	<i>Hydric soil:</i> no
<i>Ponding:</i> none	<i>Hydrologic group:</i> A
<i>Drainage class:</i> excessively drained	<i>Potential for frost action:</i> low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 4 in	fine sandy loam	moderately rapid	0.55 to 0.71 in	5.1 to 6.5
E -- 4 to 9 in	gravelly coarse sandy loam	moderately rapid	0.20 to 0.77 in	5.1 to 6.5
Bw -- 9 to 20 in	gravelly coarse sand	very rapid	0.11 to 0.66 in	5.1 to 6.5
BC,C -- 20 to 80 in	gravelly coarse sand	very rapid	0.60 to 3.59 in	5.1 to 6.5

Map Unit Description (MN)

Benton County, Minnesota

C55A--Watab loamy fine sand, 0 to 2 percent slopes

Watab

Extent: 75 to 95 percent of the unit

Landform(s): flats on interdrumlins

Slope gradient: 0 to 2 percent

Parent material: sandy outwash over dense loamy till

Restrictive feature(s): densic material at 40 to 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .20

Land capability, nonirrigated: 3w

Hydric soil: yes

Hydrologic group: C/D

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 8 in	loamy fine sand	rapid	0.79 to 0.94 in	5.1 to 6.0
E,Bw -- 8 to 23 in	loamy fine sand	rapid	0.90 to 1.35 in	5.1 to 6.5
2Bt -- 23 to 33 in	fine sandy loam	moderately rapid	0.82 to 1.23 in	5.1 to 6.5
2BC -- 33 to 45 in	fine sandy loam	slow	0.94 to 1.42 in	5.6 to 7.3
2BCd -- 45 to 80 in	fine sandy loam	very slow	1.75 to 2.80 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

C56A--Langola loamy fine sand, 0 to 2 percent slopes

Langola, wet

Extent: 80 to 95 percent of the unit

Landform(s): drumlins

Slope gradient: 0 to 2 percent

Parent material: sandy outwash over dense loamy till

Restrictive feature(s): densic material at 40 to 60 inches

Flooding: none

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .20

Land capability, nonirrigated: 3s

Hydric soil: no

Hydrologic group: C/D

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 15 in	loamy fine sand	rapid	1.50 to 1.80 in	5.1 to 6.5
Bw -- 15 to 31 in	loamy sand	rapid	1.29 to 1.61 in	5.1 to 6.5
2Bt -- 31 to 39 in	sandy loam	moderate	0.79 to 1.18 in	5.1 to 6.5
2BC -- 39 to 43 in	fine sandy loam	slow	0.31 to 0.47 in	5.6 to 7.3
2BCd -- 43 to 80 in	fine sandy loam	very slow	1.85 to 2.96 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

C57A--Warman loam, depressional, 0 to 1 percent slopes

Warman, depressional

Extent: 70 to 95 percent of the unit

Landform(s): outwash plains, stream terraces

Slope gradient: 0 to 1 percent

Parent material: outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: frequent

Drainage class: very poorly drained

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .37

Land capability, nonirrigated: 6w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 16 in	loam	moderate	2.74 to 3.55 in	5.1 to 6.5
Bg -- 16 to 34 in	loam	moderate	1.77 to 3.37 in	5.1 to 6.5
2Cg -- 34 to 60 in	gravelly coarse sand	very rapid	0.26 to 1.56 in	5.1 to 6.5

C58A--Ogilvie loam, 0 to 2 percent slopes

Ogilvie

Extent: 65 to 90 percent of the unit

Landform(s): outwash plains, stream terraces

Slope gradient: 0 to 2 percent

Parent material: outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .43

Land capability, nonirrigated: 2w

Hydric soil: no

Hydrologic group: B/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 8 in	loam	moderate	1.18 to 1.50 in	5.1 to 6.5
E -- 8 to 18 in	loam	moderate	1.54 to 2.25 in	5.1 to 6.5
Bt -- 18 to 31 in	loam	moderate	1.95 to 2.86 in	5.1 to 6.5
2C -- 31 to 60 in	gravelly coarse sand	very rapid	0.29 to 1.72 in	5.1 to 6.5

Map Unit Description (MN)

Benton County, Minnesota

C59A--Pierz sandy loam, 0 to 2 percent slopes

Pierz

Extent: 75 to 95 percent of the unit

Landform(s): stream terraces, outwash plains

Slope gradient: 0 to 2 percent

Parent material: loamy glaciofluvial deposits over sandy outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .24

Land capability, nonirrigated: 2s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

Representative soil profile:

		<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap --	0 to 10 in	sandy loam	moderately rapid	1.28 to 1.67 in	5.1 to 6.5
Bt --	10 to 23 in	sandy loam	moderately rapid	2.08 to 2.60 in	5.1 to 6.5
2Bt --	23 to 27 in	gravelly sandy loam	moderately rapid	0.69 to 0.87 in	5.1 to 6.5
2C --	27 to 60 in	gravelly coarse sand	very rapid	0.65 to 1.31 in	5.1 to 6.5

Map Unit Description (MN)

Benton County, Minnesota

C60A--Bushville fine sand, 0 to 2 percent slopes

Bushville

Extent: 75 to 90 percent of the unit

Landform(s): drumlins

Slope gradient: 0 to 2 percent

Parent material: sandy outwash over dense loamy till

Restrictive feature(s): densic material at 40 to 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 1

Wind erodibility index (WEI): 250

Kw factor (surface layer) .10

Land capability, nonirrigated: 3s

Hydric soil: no

Hydrologic group: C/D

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 10 in	fine sand	rapid	0.69 to 0.98 in	5.1 to 6.5
E -- 10 to 24 in	loamy fine sand	rapid	0.85 to 1.28 in	5.1 to 6.5
2Bt1 -- 24 to 30 in	fine sandy loam	moderate	0.59 to 0.89 in	5.1 to 6.5
2Bt2 -- 30 to 42 in	sandy loam	moderately slow	0.98 to 1.46 in	5.1 to 7.3
2BCd -- 42 to 80 in	fine sandy loam	very slow	1.89 to 3.02 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

C62C--Sartell fine sand, till substratum, 6 to 12 percent slopes

Sartell, till substratum

Extent: 65 to 95 percent of the unit

Landform(s): drumlins

Slope gradient: 6 to 12 percent

Parent material: eolian sands over dense till

Restrictive feature(s): densic material at 60 to 80 inches

Flooding: none

Ponding: none

Drainage class: excessively drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 1

Wind erodibility index (WEI): 250

Kw factor (surface layer) .10

Land capability, nonirrigated: 6s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 4 in	fine sand	rapid	0.35 to 0.43 in	5.1 to 6.0
Bw -- 4 to 33 in	fine sand	rapid	1.75 to 2.91 in	5.1 to 6.0
C -- 33 to 66 in	fine sand	rapid	1.65 to 2.98 in	5.6 to 7.3
2BCd -- 66 to 80 in	fine sandy loam	very slow	0.69 to 1.10 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

C63B--Kost, till substratum-Langola complex, 1 to 6 percent slopes

Kost, till substratum

Extent: 45 to 70 percent of the unit

Landform(s): drumlins

Slope gradient: 1 to 6 percent

Parent material: sandy outwash over dense loamy till

Restrictive feature(s): densic material at 60 to 80 inches

Flooding: none

Ponding: none

Drainage class: excessively drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .17

Land capability, nonirrigated: 4s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 14 in	loamy fine sand	rapid	1.28 to 1.70 in	5.1 to 7.3
Bw -- 14 to 33 in	fine sand	rapid	1.13 to 1.51 in	5.1 to 7.3
BC -- 33 to 66 in	fine sand	rapid	1.65 to 2.31 in	5.6 to 7.3
2BCd -- 66 to 80 in	fine sandy loam	very slow	0.69 to 1.10 in	5.6 to 7.3

Langola

Extent: 30 to 60 percent of the unit

Landform(s): drumlins

Slope gradient: 1 to 6 percent

Parent material: sandy outwash over dense loamy till

Restrictive feature(s): densic material at 40 to 60 inches

Flooding: none

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .20

Land capability, nonirrigated: 3s

Hydric soil: no

Hydrologic group: C/D

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 15 in	loamy fine sand	rapid	1.50 to 1.80 in	5.1 to 6.5
Bw -- 15 to 31 in	loamy sand	rapid	1.29 to 1.61 in	5.1 to 6.5
2Bt -- 31 to 39 in	sandy loam	moderate	0.79 to 1.18 in	5.1 to 6.5
2BC -- 39 to 43 in	fine sandy loam	slow	0.31 to 0.47 in	5.6 to 7.3
2BCd -- 43 to 80 in	fine sandy loam	very slow	1.85 to 2.96 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

C64A--Mora fine sandy loam, 1 to 3 percent slopes, stony

Mora, stony

Extent: 65 to 90 percent of the unit

Landform(s): drumlins

Slope gradient: 1 to 3 percent

Parent material: dense loamy till

Restrictive feature(s): densic material at 40 to 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .15

Land capability, nonirrigated: 2s

Hydric soil: no

Hydrologic group: C/D

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 8 in	fine sandy loam	moderately rapid	1.10 to 1.42 in	5.1 to 6.5
E -- 8 to 12 in	fine sandy loam	moderately rapid	0.43 to 0.75 in	5.1 to 6.5
B/E,Bt -- 12 to 36 in	fine sandy loam	moderately rapid	2.40 to 4.56 in	5.1 to 6.5
BC -- 36 to 46 in	fine sandy loam	slow	0.82 to 1.23 in	5.6 to 7.3
BCd -- 46 to 80 in	fine sandy loam	very slow	1.69 to 2.71 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

C64B--Mora fine sandy loam, 3 to 5 percent slopes, stony

Mora, stony

Extent: 70 to 90 percent of the unit

Landform(s): drumlins

Slope gradient: 3 to 5 percent

Parent material: dense loamy till

Restrictive feature(s): densic material at 40 to 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .15

Land capability, nonirrigated: 3s

Hydric soil: no

Hydrologic group: C/D

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 8 in	fine sandy loam	moderately rapid	1.10 to 1.42 in	5.1 to 6.5
E -- 8 to 12 in	fine sandy loam	moderately rapid	0.43 to 0.75 in	5.1 to 6.5
B/E,Bt -- 12 to 36 in	fine sandy loam	moderately rapid	2.40 to 4.56 in	5.1 to 6.5
BC -- 36 to 46 in	fine sandy loam	slow	0.82 to 1.23 in	5.6 to 7.3
BCd -- 46 to 80 in	fine sandy loam	very slow	1.69 to 2.71 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

C65A--Parent loam, 0 to 2 percent slopes, stony

Parent, stony

Extent: 65 to 95 percent of the unit

Landform(s): flats on interdrumlins

Slope gradient: 0 to 2 percent

Parent material: dense loamy till

Restrictive feature(s): densic material at 40 to 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .28

Land capability, nonirrigated: 4w

Hydric soil: yes

Hydrologic group: C/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 7 in	loam	moderate	1.42 to 1.56 in	5.6 to 7.3
Bg -- 7 to 28 in	fine sandy loam	moderate	2.50 to 3.55 in	5.6 to 7.3
BC -- 28 to 40 in	fine sandy loam	slow	0.98 to 1.46 in	5.6 to 7.3
BCd -- 40 to 80 in	fine sandy loam	very slow	1.99 to 3.18 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

C66A--St. Francis fine sandy loam, 0 to 2 percent slopes

St. francis

Extent: 75 to 95 percent of the unit

Landform(s): stream terraces, outwash plains, moraines

Slope gradient: 0 to 2 percent

Parent material: loamy glaciofluvial deposits and/or sandy and gravelly outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat excessively drained

Soil loss tolerance (T factor): 2

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .32

Land capability, nonirrigated: 3s

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

Representative soil profile:

		<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap --	0 to 6 in	fine sandy loam	moderately rapid	0.83 to 1.06 in	5.1 to 6.5
E --	6 to 11 in	fine sandy loam	moderately rapid	0.51 to 0.87 in	5.1 to 6.5
Bt1 --	11 to 16 in	loam	moderate	0.46 to 0.97 in	5.1 to 6.5
2Bt2 --	16 to 20 in	gravelly loamy coarse sand	rapid	0.04 to 0.35 in	5.1 to 6.5
2C --	20 to 80 in	gravelly coarse sand	very rapid	0.60 to 3.59 in	5.1 to 6.5

Map Unit Description (MN)

Benton County, Minnesota

C67B--Bushville complex, 1 to 6 percent slopes

Bushville

Extent: 55 to 70 percent of the unit

Landform(s): drumlins

Slope gradient: 1 to 6 percent

Parent material: sandy outwash over dense loamy till

Restrictive feature(s): densic material at 40 to 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 1

Wind erodibility index (WEI): 250

Kw factor (surface layer) .10

Land capability, nonirrigated: 3s

Hydric soil: no

Hydrologic group: C/D

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 10 in	fine sand	rapid	0.69 to 0.98 in	5.1 to 6.5
E -- 10 to 24 in	loamy fine sand	rapid	0.85 to 1.28 in	5.1 to 6.5
2Bt1 -- 24 to 30 in	fine sandy loam	moderate	0.59 to 0.89 in	5.1 to 6.5
2Bt2 -- 30 to 42 in	sandy loam	moderately slow	0.98 to 1.46 in	5.1 to 7.3
2BCd -- 42 to 80 in	fine sandy loam	very slow	1.89 to 3.02 in	5.6 to 7.3

Bushville, wet

Extent: 25 to 40 percent of the unit

Landform(s): drumlins

Slope gradient: 0 to 3 percent

Parent material: sandy outwash over dense loamy till

Restrictive feature(s): densic material at 40 to 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 1

Wind erodibility index (WEI): 250

Kw factor (surface layer) .10

Land capability, nonirrigated: 3s

Hydric soil: no

Hydrologic group: C/D

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 10 in	fine sand	rapid	0.69 to 0.98 in	5.1 to 6.5
E -- 10 to 24 in	loamy fine sand	rapid	0.85 to 1.28 in	5.1 to 6.5
2Bt1 -- 24 to 30 in	fine sandy loam	moderate	0.59 to 0.89 in	5.1 to 6.5
2Bt2 -- 30 to 42 in	sandy loam	moderately slow	0.98 to 1.46 in	5.1 to 7.3
2BCd -- 42 to 80 in	fine sandy loam	very slow	1.89 to 3.02 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

C68B--Milaca fine sandy loam, 3 to 6 percent slopes, stony

Milaca, stony

Extent: 75 to 95 percent of the unit

Landform(s): drumlins

Slope gradient: 3 to 6 percent

Parent material: dense loamy till

Restrictive feature(s): densic material at 40 to 60 inches

Flooding: none

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .15

Land capability, nonirrigated: 3s

Hydric soil: no

Hydrologic group: A/D

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 9 in	fine sandy loam	moderately rapid	1.18 to 1.63 in	5.1 to 6.5
E -- 9 to 13 in	fine sandy loam	moderately rapid	0.43 to 0.75 in	5.1 to 6.5
B/E,Bt -- 13 to 43 in	fine sandy loam	moderately rapid	2.99 to 4.49 in	5.1 to 6.5
BCd -- 43 to 80 in	fine sandy loam	very slow	1.85 to 2.96 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

C68C--Milaca fine sandy loam, 6 to 12 percent slopes, stony

Milaca, stony

Extent: 70 to 90 percent of the unit

Landform(s): drumlins, moraines

Slope gradient: 6 to 12 percent

Parent material: dense loamy till

Restrictive feature(s): densic material at 40 to 60 inches

Flooding: none

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .15

Land capability, nonirrigated: 3e

Hydric soil: no

Hydrologic group: A/D

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 4 in	fine sandy loam	moderately rapid	0.55 to 0.71 in	5.1 to 6.5
E -- 4 to 15 in	fine sandy loam	moderately rapid	1.21 to 2.09 in	5.1 to 6.5
E/B,Bt1,Bt2 -- 15 to 42 in	fine sandy loam	moderately rapid	2.72 to 4.07 in	5.1 to 6.5
BCd -- 42 to 80 in	fine sandy loam	very slow	1.89 to 3.02 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

C68E--Milaca fine sandy loam, 12 to 25 percent slopes, stony

Milaca, stony

Extent: 70 to 90 percent of the unit

Landform(s): drumlins, moraines

Slope gradient: 12 to 25 percent

Parent material: dense loamy till

Restrictive feature(s): densic material at 40 to 60 inches

Flooding: none

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .15

Land capability, nonirrigated: 6e

Hydric soil: no

Hydrologic group: A/D

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 4 in	fine sandy loam	moderately rapid	0.55 to 0.71 in	5.1 to 6.5
E -- 4 to 15 in	fine sandy loam	moderately rapid	1.21 to 2.09 in	5.1 to 6.5
E/B,Bt1,Bt2 -- 15 to 42 in	fine sandy loam	moderately rapid	2.72 to 4.07 in	5.1 to 6.5
BCd -- 42 to 80 in	fine sandy loam	very slow	1.89 to 3.02 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

C69B--Milaca, stony-St. Francis complex, 3 to 8 percent slopes

Milaca, stony

<i>Extent:</i> 40 to 75 percent of the unit	<i>Soil loss tolerance (T factor):</i> 4
<i>Landform(s):</i> end moraines, moraines	<i>Wind erodibility group (WEG):</i> 3
<i>Slope gradient:</i> 3 to 8 percent	<i>Wind erodibility index (WEI):</i> 86
<i>Parent material:</i> dense loamy till	<i>Kw factor (surface layer)</i> .15
<i>Restrictive feature(s):</i> densic material at 40 to 60 inches	<i>Land capability, nonirrigated:</i> 3s
<i>Flooding:</i> none	<i>Hydric soil:</i> no
<i>Ponding:</i> none	<i>Hydrologic group:</i> A/D
<i>Drainage class:</i> moderately well drained	<i>Potential for frost action:</i> moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 9 in	fine sandy loam	moderately rapid	1.27 to 1.63 in	5.1 to 6.5
E -- 9 to 13 in	fine sandy loam	moderately rapid	0.43 to 0.75 in	5.1 to 6.5
B/E,Bt -- 13 to 43 in	fine sandy loam	moderately rapid	2.99 to 4.49 in	5.1 to 6.5
BCd -- 43 to 80 in	fine sandy loam	very slow	1.85 to 2.96 in	5.6 to 7.3

St. francis

<i>Extent:</i> 20 to 50 percent of the unit	<i>Soil loss tolerance (T factor):</i> 2
<i>Landform(s):</i> end moraines, moraines	<i>Wind erodibility group (WEG):</i> 3
<i>Slope gradient:</i> 3 to 8 percent	<i>Wind erodibility index (WEI):</i> 86
<i>Parent material:</i> loamy glaciofluvial deposits and/or sandy and gravelly outwash	<i>Kw factor (surface layer)</i> .32
<i>Restrictive feature(s):</i> greater than 60 inches	<i>Land capability, nonirrigated:</i> 4e
<i>Flooding:</i> none	<i>Hydric soil:</i> no
<i>Ponding:</i> none	<i>Hydrologic group:</i> B
<i>Drainage class:</i> somewhat excessively drained	<i>Potential for frost action:</i> moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 6 in	fine sandy loam	moderately rapid	0.83 to 1.06 in	5.1 to 6.5
E -- 6 to 11 in	fine sandy loam	moderately rapid	0.51 to 0.87 in	5.1 to 6.5
Bt1 -- 11 to 16 in	loam	moderate	0.46 to 0.97 in	5.1 to 6.5
2Bt2 -- 16 to 20 in	gravelly loamy coarse sand	rapid	0.04 to 0.35 in	5.1 to 6.5
2C -- 20 to 80 in	gravelly coarse sand	very rapid	0.60 to 3.59 in	5.1 to 6.5

Map Unit Description (MN)

Benton County, Minnesota

C69C--Milaca, stony-St. Francis complex, 8 to 15 percent slopes

Milaca, stony

Extent: 25 to 75 percent of the unit

Landform(s): end moraines, moraines

Slope gradient: 8 to 15 percent

Parent material: dense loamy till

Restrictive feature(s): densic material at 40 to 60 inches

Flooding: none

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .15

Land capability, nonirrigated: 3e

Hydric soil: no

Hydrologic group: A/D

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 9 in	fine sandy loam	moderately rapid	1.27 to 1.63 in	5.1 to 6.5
E -- 9 to 13 in	fine sandy loam	moderately rapid	0.43 to 0.75 in	5.1 to 6.5
B/E,Bt -- 13 to 43 in	fine sandy loam	moderately rapid	2.99 to 4.49 in	5.1 to 6.5
BCd -- 43 to 80 in	fine sandy loam	very slow	1.85 to 2.96 in	5.6 to 7.3

St. francis

Extent: 15 to 45 percent of the unit

Landform(s): end moraines, moraines

Slope gradient: 8 to 15 percent

Parent material: loamy glaciofluvial deposits and/or sandy and gravelly outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat excessively drained

Soil loss tolerance (T factor): 2

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .32

Land capability, nonirrigated: 4e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 6 in	fine sandy loam	moderately rapid	0.83 to 1.06 in	5.1 to 6.5
E -- 6 to 11 in	fine sandy loam	moderately rapid	0.51 to 0.87 in	5.1 to 6.5
Bt1 -- 11 to 16 in	loam	moderate	0.46 to 0.97 in	5.1 to 6.5
2Bt2 -- 16 to 20 in	gravelly loamy coarse sand	rapid	0.04 to 0.35 in	5.1 to 6.5
2C -- 20 to 80 in	gravelly coarse sand	very rapid	0.60 to 3.59 in	5.1 to 6.5

Map Unit Description (MN)

Benton County, Minnesota

C70B--St. Francis-Mahtomedi complex, 2 to 6 percent slopes

St. francis

<i>Extent:</i> 60 to 90 percent of the unit	<i>Soil loss tolerance (T factor):</i> 2
<i>Landform(s):</i> outwash plains, stream terraces, moraines	<i>Wind erodibility group (WEG):</i> 3
<i>Slope gradient:</i> 2 to 6 percent	<i>Wind erodibility index (WEI):</i> 86
<i>Parent material:</i> loamy glaciofluvial deposits and/or sandy and gravelly outwash	<i>Kw factor (surface layer)</i> .32
<i>Restrictive feature(s):</i> greater than 60 inches	<i>Land capability, nonirrigated:</i> 3e
<i>Flooding:</i> none	<i>Hydric soil:</i> no
<i>Ponding:</i> none	<i>Hydrologic group:</i> B
<i>Drainage class:</i> somewhat excessively drained	<i>Potential for frost action:</i> moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 6 in	fine sandy loam	moderately rapid	0.83 to 1.06 in	5.1 to 6.5
E -- 6 to 11 in	fine sandy loam	moderately rapid	0.51 to 0.87 in	5.1 to 6.5
Bt1 -- 11 to 16 in	loam	moderate	0.46 to 0.97 in	5.1 to 6.5
2Bt2 -- 16 to 20 in	gravelly loamy coarse sand	rapid	0.04 to 0.35 in	5.1 to 6.5
2C -- 20 to 80 in	gravelly coarse sand	very rapid	0.60 to 3.59 in	5.1 to 6.5

Mahtomedi

<i>Extent:</i> 10 to 40 percent of the unit	<i>Soil loss tolerance (T factor):</i> 5
<i>Landform(s):</i> outwash plains, stream terraces, moraines	<i>Wind erodibility group (WEG):</i> 3
<i>Slope gradient:</i> 2 to 6 percent	<i>Wind erodibility index (WEI):</i> 86
<i>Parent material:</i> sandy outwash	<i>Kw factor (surface layer)</i> .24
<i>Restrictive feature(s):</i> greater than 60 inches	<i>Land capability, nonirrigated:</i> 4e
<i>Flooding:</i> none	<i>Hydric soil:</i> no
<i>Ponding:</i> none	<i>Hydrologic group:</i> A
<i>Drainage class:</i> excessively drained	<i>Potential for frost action:</i> low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 4 in	fine sandy loam	moderately rapid	0.55 to 0.71 in	5.1 to 6.5
E -- 4 to 9 in	gravelly coarse sandy loam	moderately rapid	0.20 to 0.77 in	5.1 to 6.5
Bw -- 9 to 20 in	gravelly coarse sand	very rapid	0.11 to 0.66 in	5.1 to 6.5
BC,C -- 20 to 80 in	gravelly coarse sand	very rapid	0.60 to 3.59 in	5.1 to 6.5

Map Unit Description (MN)

Benton County, Minnesota

C70C--St. Francis-Mahtomedi complex, 6 to 12 percent slopes

St. francis

<i>Extent:</i> 50 to 85 percent of the unit	<i>Soil loss tolerance (T factor):</i> 2
<i>Landform(s):</i> outwash plains, stream terraces, moraines	<i>Wind erodibility group (WEG):</i> 3
<i>Slope gradient:</i> 6 to 12 percent	<i>Wind erodibility index (WEI):</i> 86
<i>Parent material:</i> loamy glaciofluvial deposits and/or sandy and gravelly outwash	<i>Kw factor (surface layer)</i> .32
<i>Restrictive feature(s):</i> greater than 60 inches	<i>Land capability, nonirrigated:</i> 4e
<i>Flooding:</i> none	<i>Hydric soil:</i> no
<i>Ponding:</i> none	<i>Hydrologic group:</i> B
<i>Drainage class:</i> somewhat excessively drained	<i>Potential for frost action:</i> moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 6 in	fine sandy loam	moderately rapid	0.83 to 1.06 in	5.1 to 6.5
E -- 6 to 11 in	fine sandy loam	moderately rapid	0.51 to 0.87 in	5.1 to 6.5
Bt1 -- 11 to 16 in	loam	moderate	0.46 to 0.97 in	5.1 to 6.5
2Bt2 -- 16 to 20 in	gravelly loamy coarse sand	rapid	0.04 to 0.35 in	5.1 to 6.5
2C -- 20 to 80 in	gravelly coarse sand	very rapid	0.60 to 3.59 in	5.1 to 6.5

Mahtomedi

<i>Extent:</i> 15 to 50 percent of the unit	<i>Soil loss tolerance (T factor):</i> 5
<i>Landform(s):</i> outwash plains, stream terraces, moraines	<i>Wind erodibility group (WEG):</i> 3
<i>Slope gradient:</i> 6 to 12 percent	<i>Wind erodibility index (WEI):</i> 86
<i>Parent material:</i> sandy outwash	<i>Kw factor (surface layer)</i> .24
<i>Restrictive feature(s):</i> greater than 60 inches	<i>Land capability, nonirrigated:</i> 6s
<i>Flooding:</i> none	<i>Hydric soil:</i> no
<i>Ponding:</i> none	<i>Hydrologic group:</i> A
<i>Drainage class:</i> excessively drained	<i>Potential for frost action:</i> low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 4 in	fine sandy loam	moderately rapid	0.55 to 0.71 in	5.1 to 6.5
E -- 4 to 9 in	gravelly coarse sandy loam	moderately rapid	0.20 to 0.77 in	5.1 to 6.5
Bw -- 9 to 20 in	gravelly coarse sand	very rapid	0.11 to 0.66 in	5.1 to 6.5
BC,C -- 20 to 80 in	gravelly coarse sand	very rapid	0.60 to 3.59 in	5.1 to 6.5

Map Unit Description (MN)

Benton County, Minnesota

C71A--Blomford loamy fine sand, 0 to 2 percent slopes

Blomford

Extent: 80 to 90 percent of the unit

Landform(s): flats on moraines

Slope gradient: 0 to 2 percent

Parent material: sandy outwash over till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .28

Land capability, nonirrigated: 3w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 9 in	loamy fine sand	rapid	0.72 to 1.09 in	5.1 to 7.3
E,E&Bt -- 9 to 33 in	fine sand	rapid	1.20 to 1.92 in	5.1 to 7.3
2Btg -- 33 to 42 in	fine sandy loam	moderate	1.18 to 1.54 in	5.1 to 7.3
2Cg -- 42 to 60 in	fine sandy loam	moderate	1.77 to 2.66 in	6.1 to 8.4

Map Unit Description (MN)

Benton County, Minnesota

C72B--Langola complex, 1 to 6 percent slopes

Langola

Extent: 50 to 85 percent of the unit

Landform(s): drumlins

Slope gradient: 1 to 6 percent

Parent material: sandy outwash over dense loamy till

Restrictive feature(s): densic material at 40 to 60 inches

Flooding: none

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .20

Land capability, nonirrigated: 3s

Hydric soil: no

Hydrologic group: C/D

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 15 in	loamy fine sand	rapid	1.50 to 1.80 in	5.1 to 6.5
Bw -- 15 to 31 in	loamy sand	rapid	1.29 to 1.61 in	5.1 to 6.5
2Bt -- 31 to 39 in	sandy loam	moderate	0.79 to 1.18 in	5.1 to 6.5
2BC -- 39 to 43 in	fine sandy loam	slow	0.31 to 0.47 in	5.6 to 7.3
2BCd -- 43 to 80 in	fine sandy loam	very slow	1.85 to 2.96 in	5.6 to 7.3

Langola, wet

Extent: 15 to 35 percent of the unit

Landform(s): drumlins

Slope gradient: 0 to 3 percent

Parent material: sandy outwash over dense loamy till

Restrictive feature(s): densic material at 40 to 60 inches

Flooding: none

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .20

Land capability, nonirrigated: 3s

Hydric soil: no

Hydrologic group: C/D

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 15 in	loamy fine sand	rapid	1.50 to 1.80 in	5.1 to 6.5
Bw -- 15 to 31 in	loamy sand	rapid	1.29 to 1.61 in	5.1 to 6.5
2Bt -- 31 to 39 in	sandy loam	moderate	0.79 to 1.18 in	5.1 to 6.5
2BC -- 39 to 43 in	fine sandy loam	slow	0.31 to 0.47 in	5.6 to 7.3
2BCd -- 43 to 80 in	fine sandy loam	very slow	1.85 to 2.96 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

C73A--Mora loam, 1 to 3 percent slopes, stony

Mora, stony

Extent: 65 to 90 percent of the unit

Landform(s): drumlins

Slope gradient: 1 to 3 percent

Parent material: dense loamy till

Restrictive feature(s): densic material at 40 to 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .32

Land capability, nonirrigated: 2s

Hydric soil: no

Hydrologic group: C/D

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 8 in	loam	moderate	1.42 to 1.81 in	5.1 to 6.5
E -- 8 to 12 in	fine sandy loam	moderately rapid	0.43 to 0.75 in	5.1 to 6.5
B/E,Bt -- 12 to 36 in	fine sandy loam	moderately rapid	2.40 to 4.56 in	5.1 to 6.5
BC -- 36 to 46 in	fine sandy loam	slow	0.82 to 1.23 in	5.6 to 7.3
BCd -- 46 to 80 in	fine sandy loam	very slow	1.69 to 2.71 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

C74A--Parent loam, depressional, 0 to 1 percent slopes, stony

Parent, depressional, stony

Extent: 70 to 95 percent of the unit

Landform(s): depressions on interdrumlins

Slope gradient: 0 to 1 percent

Parent material: dense loamy till

Restrictive feature(s): densic material at 40 to 60 inches

Flooding: none

Ponding: frequent

Drainage class: very poorly drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .28

Land capability, nonirrigated: 6w

Hydric soil: yes

Hydrologic group: C/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 7 in	loam	moderate	1.42 to 1.56 in	5.6 to 7.3
Bg -- 7 to 28 in	fine sandy loam	moderate	2.50 to 3.55 in	5.6 to 7.3
BC -- 28 to 40 in	fine sandy loam	slow	0.98 to 1.46 in	5.6 to 7.3
BCd -- 40 to 80 in	fine sandy loam	very slow	1.99 to 3.18 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

C75A--Seelyeville and Cathro soils, milaca catena, depressional, 0 to 1 percent slopes

Seelyeville, depressional

Extent: 0 to 95 percent of the unit

Landform(s): depressions on interdrumlins

Slope gradient: 0 to 1 percent

Parent material: highly decomposed organic material

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: frequent

Drainage class: very poorly drained

Soil loss tolerance (T factor): 2

Wind erodibility group (WEG): 8

Wind erodibility index (WEI): 0

Kw factor (surface layer) .02

Land capability, nonirrigated: 7w

Hydric soil: yes

Hydrologic group: A/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Oa1 -- 0 to 10 in	muck	moderately rapid	3.44 to 4.43 in	
Oa2,Oa5 -- 10 to 80 in	muck	moderately rapid	24.53 to 31.54 in	

Cathro, depressional, milaca catena

Extent: 0 to 95 percent of the unit

Landform(s): depressions on interdrumlins

Slope gradient: 0 to 1 percent

Parent material: highly decomposed organic material over dense loamy till

Restrictive feature(s): densic material at 40 to 80 inches

Flooding: none

Ponding: frequent

Drainage class: very poorly drained

Soil loss tolerance (T factor): 1

Wind erodibility group (WEG): 8

Wind erodibility index (WEI): 0

Kw factor (surface layer) .02

Land capability, nonirrigated: 7w

Hydric soil: yes

Hydrologic group: C/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Oa -- 0 to 31 in	muck	moderately rapid	10.89 to 14.00 in	
A -- 31 to 37 in	loam	moderate	1.18 to 1.42 in	5.6 to 6.5
BC -- 37 to 62 in	fine sandy loam	slow	2.02 to 3.02 in	5.6 to 7.3
BCd -- 62 to 80 in	fine sandy loam	very slow	0.89 to 1.42 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

C77A--Novak loam, 0 to 2 percent slopes

Novak

Extent: 65 to 80 percent of the unit

Landform(s): outwash plains, stream terraces

Slope gradient: 0 to 2 percent

Parent material: outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .43

Land capability, nonirrigated: 2s

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 8 in	loam	moderate	1.18 to 1.50 in	5.1 to 6.5
B/E -- 8 to 15 in	loam	moderate	1.06 to 1.56 in	5.1 to 6.5
Bt1 -- 15 to 26 in	loam	moderate	1.65 to 2.43 in	5.1 to 6.5
2Bt2 -- 26 to 30 in	sandy loam	moderate	0.35 to 0.67 in	5.1 to 6.5
3C -- 30 to 60 in	gravelly coarse sand	very rapid	0.30 to 1.80 in	5.1 to 6.5

Map Unit Description (MN)

Benton County, Minnesota

C77B--Novak loam, 2 to 6 percent slopes

Novak

Extent: 70 to 90 percent of the unit

Landform(s): outwash plains, stream terraces

Slope gradient: 2 to 6 percent

Parent material: outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .43

Land capability, nonirrigated: 2e

Hydric soil: no

Hydrologic group: B

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 7 in	loam	moderate	1.06 to 1.35 in	5.1 to 6.5
B/E -- 7 to 17 in	loam	moderate	1.48 to 2.17 in	5.1 to 6.5
Bt1 -- 17 to 23 in	loam	moderate	0.89 to 1.30 in	5.1 to 6.5
2Bt2 -- 23 to 26 in	sandy loam	moderate	0.28 to 0.54 in	5.1 to 6.5
3C -- 26 to 60 in	gravelly coarse sand	very rapid	0.34 to 2.03 in	5.1 to 6.5

Map Unit Description (MN)

Benton County, Minnesota

C78A--Warman loam, 0 to 2 percent slopes

Warman

Extent: 70 to 90 percent of the unit

Landform(s): outwash plains, stream terraces

Slope gradient: 0 to 2 percent

Parent material: outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .37

Land capability, nonirrigated: 4w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 9 in	loam	moderate	1.36 to 1.99 in	5.1 to 6.5
AB -- 9 to 12 in	loam	moderate	0.28 to 0.52 in	5.1 to 6.5
Bg -- 12 to 33 in	loam	moderate	1.91 to 4.04 in	5.1 to 6.5
2C -- 33 to 60 in	gravelly coarse sand	very rapid	0.27 to 1.61 in	5.1 to 6.5

Map Unit Description (MN)

Benton County, Minnesota

C79A--Mora loam, 1 to 3 percent slopes, very stony

Mora, very stony

Extent: 70 to 90 percent of the unit

Landform(s): drumlins

Slope gradient: 1 to 3 percent

Parent material: dense loamy till

Restrictive feature(s): densic material at 40 to 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .32

Land capability, nonirrigated: 6s

Hydric soil: no

Hydrologic group: C/D

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 8 in	loam	moderate	1.42 to 1.81 in	5.1 to 6.5
E -- 8 to 12 in	fine sandy loam	moderately rapid	0.43 to 0.75 in	5.1 to 6.5
B/E,Bt -- 12 to 36 in	fine sandy loam	moderately rapid	2.40 to 4.56 in	5.1 to 6.5
BC -- 36 to 46 in	fine sandy loam	slow	0.82 to 1.23 in	5.6 to 7.3
BCd -- 46 to 80 in	fine sandy loam	very slow	1.69 to 2.71 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

C80A--Parent loam, depressional, 0 to 1 percent slopes, very stony

Parent, depressional, very stony

Extent: 70 to 95 percent of the unit

Landform(s): depressions on interdrumlins

Slope gradient: 0 to 1 percent

Parent material: dense loamy till

Restrictive feature(s): densic material at 40 to 60 inches

Flooding: none

Ponding: frequent

Drainage class: very poorly drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .28

Land capability, nonirrigated: 6s

Hydric soil: yes

Hydrologic group: C/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 7 in	loam	moderate	1.42 to 1.56 in	5.6 to 7.3
Bg -- 7 to 28 in	fine sandy loam	moderate	2.50 to 3.55 in	5.6 to 7.3
BC -- 28 to 40 in	fine sandy loam	slow	0.98 to 1.46 in	5.6 to 7.3
BCd -- 40 to 80 in	fine sandy loam	very slow	1.99 to 3.18 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

C82A--Ronneby loam, 0 to 2 percent slopes, very stony

Ronneby, very stony

Extent: 75 to 95 percent of the unit

Landform(s): interdrumlins

Slope gradient: 0 to 2 percent

Parent material: dense loamy till

Restrictive feature(s): densic material at 40 to 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .32

Land capability, nonirrigated: 6s

Hydric soil: no

Hydrologic group: C/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 4 in	loam	moderate	0.71 to 0.91 in	5.1 to 6.5
E -- 4 to 12 in	fine sandy loam	moderately rapid	0.94 to 1.50 in	5.1 to 6.5
Bt1,Bt2 -- 12 to 33 in	fine sandy loam	moderate	2.55 to 4.04 in	5.6 to 6.5
Bt3 -- 33 to 45 in	fine sandy loam	moderately slow	0.94 to 1.42 in	5.6 to 7.3
BCd -- 45 to 80 in	fine sandy loam	very slow	1.75 to 2.80 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

C83A--Parent loam, 0 to 2 percent slopes, very stony

Parent, very stony

Extent: 70 to 95 percent of the unit

Landform(s): flats on interdrumlins

Slope gradient: 0 to 2 percent

Parent material: dense loamy till

Restrictive feature(s): densic material at 40 to 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .28

Land capability, nonirrigated: 6s

Hydric soil: yes

Hydrologic group: C/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 7 in	loam	moderate	1.42 to 1.56 in	5.6 to 7.3
Bg -- 7 to 28 in	fine sandy loam	moderate	2.50 to 3.55 in	5.6 to 7.3
BC -- 28 to 40 in	fine sandy loam	slow	0.98 to 1.46 in	5.6 to 7.3
BCd -- 40 to 80 in	fine sandy loam	very slow	1.99 to 3.18 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

C84B--Brennyville complex, 1 to 6 percent slopes, very stony

Brennyville, very stony

Extent: 40 to 85 percent of the unit

Landform(s): drumlins, moraines

Slope gradient: 3 to 6 percent

Parent material: silt mantled dense loamy till

Restrictive feature(s): densic material at 40 to 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .43

Land capability, nonirrigated: 6s

Hydric soil: no

Hydrologic group: C/D

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 8 in	silt loam	moderate	1.50 to 1.89 in	5.1 to 6.5
B/E -- 8 to 11 in	silt loam	moderate	0.63 to 0.69 in	5.1 to 6.5
Bt1 -- 11 to 21 in	silt loam	moderate	1.67 to 2.17 in	5.1 to 6.5
2Bt2,2Bt3 -- 21 to 38 in	fine sandy loam	moderately rapid	1.56 to 3.12 in	5.1 to 7.3
2BC -- 38 to 45 in	fine sandy loam	slow	0.54 to 0.80 in	5.6 to 7.3
2BCd -- 45 to 80 in	fine sandy loam	very slow	1.75 to 2.80 in	5.6 to 7.3

Brennyville, wet, very stony

Extent: 5 to 35 percent of the unit

Landform(s): drumlins, moraines

Slope gradient: 1 to 3 percent

Parent material: silt mantled dense loamy till

Restrictive feature(s): densic material at 40 to 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .43

Land capability, nonirrigated: 6s

Hydric soil: no

Hydrologic group: C/D

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 8 in	silt loam	moderate	1.50 to 1.89 in	5.1 to 6.5
B/E -- 8 to 11 in	silt loam	moderate	0.63 to 0.69 in	5.1 to 6.5
Bt1 -- 11 to 21 in	silt loam	moderate	1.67 to 2.17 in	5.1 to 6.5
2Bt2,2Bt3 -- 21 to 38 in	fine sandy loam	moderately rapid	1.56 to 3.12 in	5.1 to 7.3
2BC -- 38 to 45 in	fine sandy loam	slow	0.54 to 0.80 in	5.6 to 7.3
2BCd -- 45 to 80 in	fine sandy loam	very slow	1.75 to 2.80 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

C84B--Brennyville complex, 1 to 6 percent slopes, very stony

Map Unit Description (MN)

Benton County, Minnesota

C89A--Cathro and Seelyeville soils, grasston catena, depressional, 0 to 1 percent slopes

Cathro, depressional, grasston catena

Extent: 0 to 95 percent of the unit

Landform(s): lake plains

Slope gradient: 0 to 1 percent

Parent material: highly decomposed organic material over clayey glaciolacustrine deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: frequent

Drainage class: very poorly drained

Soil loss tolerance (T factor): 1

Wind erodibility group (WEG): 8

Wind erodibility index (WEI): 0

Kw factor (surface layer) .02

Land capability, nonirrigated: 7w

Hydric soil: yes

Hydrologic group: C/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Oa -- 0 to 31 in	muck	moderately rapid	10.89 to 14.00 in	
A -- 31 to 37 in	loam	moderate	1.18 to 1.30 in	5.6 to 6.5
Cg1 -- 37 to 62 in	silty clay	slow	2.02 to 5.04 in	7.4 to 8.4
Cg2 -- 62 to 80 in	silty clay	slow	1.42 to 3.54 in	7.4 to 8.4

Seelyeville, depressional

Extent: 0 to 95 percent of the unit

Landform(s): lake plains

Slope gradient: 0 to 1 percent

Parent material: highly decomposed organic material

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: frequent

Drainage class: very poorly drained

Soil loss tolerance (T factor): 2

Wind erodibility group (WEG): 8

Wind erodibility index (WEI): 0

Kw factor (surface layer) .02

Land capability, nonirrigated: 7w

Hydric soil: yes

Hydrologic group: A/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Oa1 -- 0 to 10 in	muck	moderately rapid	3.44 to 4.43 in	
Oa2,Oa5 -- 10 to 80 in	muck	moderately rapid	24.53 to 31.54 in	

Map Unit Description (MN)

Benton County, Minnesota

C90A--Brennyville-Cebana complex, 0 to 3 percent slopes, very stony

Brennyville, wet, very stony

Extent: 68 to 98 percent of the unit

Landform(s): drumlins, moraines

Slope gradient: 1 to 3 percent

Parent material: silt mantled dense loamy till

Restrictive feature(s): densic material at 40 to 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .43

Land capability, nonirrigated: 6s

Hydric soil: no

Hydrologic group: C/D

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 8 in	silt loam	moderate	1.50 to 1.89 in	5.1 to 6.5
B/E -- 8 to 11 in	silt loam	moderate	0.63 to 0.69 in	5.1 to 6.5
Bt1 -- 11 to 21 in	silt loam	moderate	1.67 to 2.17 in	5.1 to 6.5
2Bt2,2Bt3 -- 21 to 38 in	fine sandy loam	moderately rapid	1.56 to 3.12 in	5.1 to 7.3
2BC -- 38 to 45 in	fine sandy loam	slow	0.54 to 0.80 in	5.6 to 7.3
2BCd -- 45 to 80 in	fine sandy loam	very slow	1.75 to 2.80 in	5.6 to 7.3

Cebana, very stony

Extent: 5 to 30 percent of the unit

Landform(s): drumlins, moraines

Slope gradient: 0 to 1 percent

Parent material: loamy mantled dense loamy till

Restrictive feature(s): densic material at 40 to 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .37

Land capability, nonirrigated: 6s

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 8 in	silt loam	moderate	1.50 to 1.89 in	5.1 to 6.5
Eg,E/B -- 8 to 27 in	silt loam	moderate	3.28 to 4.24 in	5.1 to 6.5
2Bt -- 27 to 49 in	loam	moderate	1.95 to 3.90 in	5.1 to 7.3
2BCd -- 49 to 80 in	fine sandy loam	very slow	1.56 to 2.49 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

C91B--Milaca, stony-Chetek complex, 3 to 8 percent slopes

Milaca, stony

Extent: 30 to 70 percent of the unit

Landform(s): drumlins, moraines

Slope gradient: 3 to 8 percent

Parent material: dense loamy till

Restrictive feature(s): densic material at 40 to 60 inches

Flooding: none

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .15

Land capability, nonirrigated: 3s

Hydric soil: no

Hydrologic group: A/D

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 9 in	fine sandy loam	moderately rapid	1.27 to 1.63 in	5.1 to 6.5
E -- 9 to 13 in	fine sandy loam	moderately rapid	0.43 to 0.75 in	5.1 to 6.5
B/E,Bt -- 13 to 43 in	fine sandy loam	moderately rapid	2.99 to 4.49 in	5.1 to 6.5
BCd -- 43 to 80 in	fine sandy loam	very slow	1.85 to 2.96 in	5.6 to 7.3

Chetek

Extent: 20 to 50 percent of the unit

Landform(s): drumlins, moraines

Slope gradient: 3 to 8 percent

Parent material: loamy glaciofluvial deposits and/or sandy and gravelly outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat excessively drained

Soil loss tolerance (T factor): 2

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .28

Land capability, nonirrigated: 6s

Hydric soil: no

Hydrologic group: A

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 5 in	fine sandy loam	moderately rapid	0.72 to 0.92 in	5.1 to 6.0
E -- 5 to 12 in	fine sandy loam	moderately rapid	0.67 to 1.14 in	5.1 to 6.0
Bt1 -- 12 to 18 in	sandy loam	moderately rapid	0.57 to 1.20 in	5.1 to 6.0
2Bt2 -- 18 to 25 in	gravelly loamy coarse sand	rapid	0.07 to 0.64 in	5.1 to 6.5
2BC,2C -- 25 to 80 in	gravelly coarse sand	very rapid	0.55 to 3.28 in	5.1 to 6.5

Map Unit Description (MN)

Benton County, Minnesota

C91C--Milaca, stony-Chetek complex, 8 to 15 percent slopes

Milaca, stony

Extent: 30 to 70 percent of the unit

Landform(s): drumlins, moraines

Slope gradient: 8 to 15 percent

Parent material: dense loamy till

Restrictive feature(s): densic material at 40 to 60 inches

Flooding: none

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 4

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .15

Land capability, nonirrigated: 3e

Hydric soil: no

Hydrologic group: A/D

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 9 in	fine sandy loam	moderately rapid	1.27 to 1.63 in	5.1 to 6.5
E -- 9 to 13 in	fine sandy loam	moderately rapid	0.43 to 0.75 in	5.1 to 6.5
B/E,Bt -- 13 to 43 in	fine sandy loam	moderately rapid	2.99 to 4.49 in	5.1 to 6.5
BCd -- 43 to 80 in	fine sandy loam	very slow	1.85 to 2.96 in	5.6 to 7.3

Chetek

Extent: 20 to 50 percent of the unit

Landform(s): drumlins, moraines

Slope gradient: 8 to 15 percent

Parent material: loamy glaciofluvial deposits and/or sandy and gravelly outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat excessively drained

Soil loss tolerance (T factor): 2

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .28

Land capability, nonirrigated: 6s

Hydric soil: no

Hydrologic group: A

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 5 in	fine sandy loam	moderately rapid	0.72 to 0.92 in	5.1 to 6.0
E -- 5 to 12 in	fine sandy loam	moderately rapid	0.67 to 1.14 in	5.1 to 6.0
Bt1 -- 12 to 18 in	sandy loam	moderately rapid	0.57 to 1.20 in	5.1 to 6.0
2Bt2 -- 18 to 25 in	gravelly loamy coarse sand	rapid	0.07 to 0.64 in	5.1 to 6.5
2BC,2C -- 25 to 80 in	gravelly coarse sand	very rapid	0.55 to 3.28 in	5.1 to 6.5

Map Unit Description (MN)

Benton County, Minnesota

C122D--Emmert-Chetek complex, 8 to 25 percent slopes

Emmert

Extent: 30 to 60 percent of the unit

Landform(s): eskers, outwash plains, stream terraces

Slope gradient: 8 to 25 percent

Parent material: gravelly outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: excessively drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .17

Land capability, nonirrigated: 6s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 2 in	fine sandy loam	rapid	0.16 to 0.24 in	5.1 to 6.5
E -- 2 to 12 in	gravelly loamy sand	rapid	0.49 to 1.48 in	5.1 to 6.5
Bt -- 12 to 37 in	gravelly loamy coarse sand	rapid	0.76 to 2.27 in	5.1 to 6.5
C -- 37 to 80 in	very gravelly coarse sand	very rapid	0.43 to 2.57 in	5.1 to 6.5

Chetek

Extent: 30 to 40 percent of the unit

Landform(s): eskers, outwash plains, stream terraces

Slope gradient: 8 to 25 percent

Parent material: loamy glaciofluvial deposits and/or sandy and gravelly outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat excessively drained

Soil loss tolerance (T factor): 2

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .24

Land capability, nonirrigated: 6e

Hydric soil: no

Hydrologic group: A

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 5 in	fine sandy loam	moderately rapid	0.72 to 0.92 in	5.1 to 6.0
E -- 5 to 12 in	fine sandy loam	moderately rapid	0.67 to 1.14 in	5.1 to 6.0
Bt1 -- 12 to 18 in	sandy loam	moderately rapid	0.57 to 1.20 in	5.1 to 6.0
2Bt2 -- 18 to 25 in	gravelly loamy coarse sand	rapid	0.07 to 0.64 in	5.1 to 6.5
2BC,2C -- 25 to 80 in	gravelly coarse sand	very rapid	0.55 to 3.28 in	5.1 to 6.5

Map Unit Description (MN)

Benton County, Minnesota

C125A--Hulligan loam, depressional, 0 to 1 percent slopes

Hulligan, depressional

Extent: 70 to 85 percent of the unit

Landform(s): outwash plains, stream terraces

Slope gradient: 0 to 1 percent

Parent material: loamy glaciofluvial deposits and/or sandy and gravelly outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: frequent

Drainage class: very poorly drained

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .37

Land capability, nonirrigated: 6w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

Representative soil profile:

	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 9 in	loam	moderate	1.54 to 1.99 in	5.1 to 6.5
Bw -- 9 to 17 in	loam	moderate	0.79 to 1.50 in	5.1 to 6.5
Bg -- 17 to 27 in	loam	moderate	0.92 to 1.94 in	5.1 to 6.5
2Cg -- 27 to 60 in	gravelly coarse sand	very rapid	0.33 to 1.96 in	5.6 to 6.5

Map Unit Description (MN)

Benton County, Minnesota

C126B--Balmlake-Rosy complex, 1 to 6 percent slopes

Balmlake

Extent: 50 to 80 percent of the unit

Landform(s): lake plains

Slope gradient: 1 to 6 percent

Parent material: loamy glaciolacustrine deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .37

Land capability, nonirrigated: 2e

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 10 in	very fine sandy loam	moderate	1.57 to 2.17 in	5.1 to 6.5
B/E -- 10 to 15 in	loam	moderate	0.41 to 0.97 in	5.1 to 6.5
Bt -- 15 to 41 in	loam	moderate	1.82 to 5.72 in	5.1 to 7.3
2BC,2C -- 41 to 80 in	stratified loamy very fine sand to silty clay loam	rapid	3.12 to 8.57 in	5.1 to 7.3

Rosy

Extent: 15 to 30 percent of the unit

Landform(s): lake plains

Slope gradient: 1 to 4 percent

Parent material: loamy glaciolacustrine deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .43

Land capability, nonirrigated: 2s

Hydric soil: no

Hydrologic group: C

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 9 in	very fine sandy loam	moderate	1.45 to 1.99 in	5.1 to 6.5
E -- 9 to 20 in	loamy very fine sand	moderately rapid	0.99 to 2.09 in	5.1 to 6.5
Bt -- 20 to 54 in	loam	moderate	2.71 to 7.45 in	5.1 to 7.3
2C -- 54 to 80 in	stratified loamy very fine sand to silty clay loam	rapid	2.08 to 5.72 in	5.1 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

C128A--Talmoon loam, 0 to 2 percent slopes

Talmoon

Extent: 80 to 95 percent of the unit

Landform(s): moraines

Slope gradient: 0 to 2 percent

Parent material: loamy till

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .32

Land capability, nonirrigated: 3w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 7 in	loam	moderate	1.20 to 1.49 in	5.1 to 6.5
Eg -- 7 to 12 in	fine sandy loam	moderately rapid	0.61 to 0.85 in	5.1 to 6.5
Btg -- 12 to 32 in	loam	moderate	2.41 to 3.61 in	5.6 to 7.3
Cg -- 32 to 80 in	loam	moderate	5.76 to 8.65 in	7.4 to 8.4

Map Unit Description (MN)

Benton County, Minnesota

C129A--Longsiding fine sandy loam, terrace, 0 to 2 percent slopes

Longsiding, terrace

Extent: 50 to 75 percent of the unit

Landform(s): lake plains, terraces

Slope gradient: 0 to 2 percent

Parent material: silty and clayey glaciolacustrine deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .24

Land capability, nonirrigated: 2w

Hydric soil: no

Hydrologic group: C/D

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 9 in	fine sandy loam	moderately rapid	1.27 to 1.63 in	5.1 to 6.5
B/E -- 9 to 12 in	silty clay loam	moderately slow	0.30 to 0.61 in	5.1 to 7.3
Bt -- 12 to 35 in	silty clay	slow	1.86 to 4.65 in	5.1 to 7.3
Bk -- 35 to 45 in	silt loam	moderate	1.77 to 2.17 in	7.4 to 8.4
C -- 45 to 80 in	silt loam	moderate	5.96 to 7.71 in	7.4 to 8.4

Map Unit Description (MN)

Benton County, Minnesota

D1B--Anoka and Zimmerman soils, terrace, 2 to 6 percent slopes

Anoka, terrace

Extent: 30 to 60 percent of the unit

Landform(s): hills on stream terraces

Slope gradient: 2 to 6 percent

Parent material: sandy outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .20

Land capability, nonirrigated: 4s

Hydric soil: no

Hydrologic group: A

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 10 in	loamy fine sand	rapid	1.28 to 1.57 in	5.1 to 6.5
E,E&Bt -- 10 to 60 in	fine sand	rapid	3.00 to 6.00 in	5.1 to 7.3

Zimmerman, terrace

Extent: 30 to 60 percent of the unit

Landform(s): hills on stream terraces

Slope gradient: 2 to 4 percent

Parent material: sandy outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: excessively drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 1

Wind erodibility index (WEI): 250

Kw factor (surface layer) .10

Land capability, nonirrigated: 4s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 9 in	fine sand	rapid	0.63 to 0.81 in	5.1 to 6.5
E,E&Bt -- 9 to 60 in	fine sand	rapid	3.05 to 5.08 in	5.1 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

D1C--Anoka and Zimmerman soils, terrace, 6 to 12 percent slopes

Anoka, terrace

Extent: 35 to 65 percent of the unit

Landform(s): hills on stream terraces

Slope gradient: 6 to 12 percent

Parent material: sandy outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .20

Land capability, nonirrigated: 4s

Hydric soil: no

Hydrologic group: A

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 10 in	loamy fine sand	rapid	1.28 to 1.57 in	5.1 to 6.5
E,E&Bt -- 10 to 60 in	fine sand	rapid	3.00 to 6.00 in	5.1 to 7.3

Zimmerman, terrace

Extent: 35 to 65 percent of the unit

Landform(s): hills on stream terraces

Slope gradient: 6 to 12 percent

Parent material: sandy outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: excessively drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 1

Wind erodibility index (WEI): 250

Kw factor (surface layer) .10

Land capability, nonirrigated: 4s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 9 in	fine sand	rapid	0.63 to 0.81 in	5.1 to 6.5
E,E&Bt -- 9 to 60 in	fine sand	rapid	3.05 to 5.08 in	5.1 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

D2A--Elkriver fine sandy loam, 0 to 2 percent slopes, rarely flooded

Elkriver, rarely flooded

Extent: 90 to 100 percent of the unit
Landform(s): flats on benches on flood plains
Slope gradient: 0 to 2 percent
Parent material: alluvium
Restrictive feature(s): greater than 60 inches
Flooding: rare
Ponding: none
Drainage class: moderately well drained

Soil loss tolerance (T factor): 3
Wind erodibility group (WEG): 3
Wind erodibility index (WEI): 86
Kw factor (surface layer) .28
Land capability, nonirrigated: 2s
Hydric soil: no
Hydrologic group: B
Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 10 in	fine sandy loam	moderately rapid	1.57 to 1.97 in	5.1 to 7.3
A1,A2,A3 -- 10 to 35 in	fine sandy loam	moderately rapid	3.78 to 5.04 in	5.1 to 7.3
Bw -- 35 to 39 in	fine sandy loam	moderately rapid	0.59 to 0.75 in	5.6 to 7.8
2C -- 39 to 80 in	sand	rapid	0.82 to 4.09 in	5.6 to 7.8

Map Unit Description (MN)

Benton County, Minnesota

D3A--Elkriver fine sandy loam, 0 to 2 percent slopes, occasionally flooded

Elkriver, occasionally flooded

Extent: 75 to 95 percent of the unit

Landform(s): flood plains

Slope gradient: 0 to 2 percent

Parent material: loamy alluvium and/or sandy and gravelly alluvium

Restrictive feature(s): greater than 60 inches

Flooding: occasional

Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .28

Land capability, nonirrigated: 2w

Hydric soil: no

Hydrologic group: B/D

Potential for frost action: moderate

Representative soil profile:

	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 10 in	fine sandy loam	moderately rapid	1.38 to 1.77 in	5.1 to 7.3
A1,A2,A3 -- 10 to 26 in	fine sandy loam	moderately rapid	2.26 to 3.07 in	5.1 to 7.3
Bw -- 26 to 32 in	very fine sandy loam	moderate	0.77 to 1.12 in	5.6 to 7.8
2C -- 32 to 80 in	sand	very rapid	1.92 to 3.36 in	5.6 to 7.8

Map Unit Description (MN)

Benton County, Minnesota

D6A--Verndale sandy loam, acid substratum, 0 to 2 percent slopes

Verndale, acid substratum

Extent: 80 to 100 percent of the unit

Landform(s): stream terraces, outwash plains

Slope gradient: 0 to 2 percent

Parent material: loamy glaciofluvial deposits over sandy outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat excessively drained

Soil loss tolerance (T factor): 2

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .24

Land capability, nonirrigated: 3s

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

<i>Representative soil profile:</i>		<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap --	0 to 10 in	sandy loam	moderately rapid	1.28 to 1.67 in	5.1 to 7.3
Bt --	10 to 19 in	sandy loam	moderate	1.27 to 1.63 in	5.1 to 7.3
2Bw --	19 to 28 in	sand	rapid	0.54 to 0.72 in	5.1 to 7.3
2C --	28 to 80 in	sand	rapid	1.04 to 3.12 in	5.1 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

D7A--Hubbard loamy sand, 0 to 2 percent slopes

Hubbard

Extent: 90 to 100 percent of the unit

Landform(s): stream terraces, outwash plains

Slope gradient: 0 to 2 percent

Parent material: sandy outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: excessively drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .10

Land capability, nonirrigated: 4s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,AB -- 0 to 20 in	loamy sand	rapid	1.61 to 2.41 in	5.1 to 7.3
Bw -- 20 to 32 in	loamy sand	rapid	0.35 to 0.83 in	5.1 to 7.3
BC,C -- 32 to 80 in	sand	rapid	1.44 to 3.36 in	5.6 to 7.8

D7B--Hubbard loamy sand, 2 to 6 percent slopes

Hubbard

Extent: 85 to 100 percent of the unit

Landform(s): hills on stream terraces, hills on outwash plains

Slope gradient: 2 to 6 percent

Parent material: sandy outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: excessively drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .10

Land capability, nonirrigated: 4s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 18 in	loamy sand	rapid	1.45 to 2.17 in	5.1 to 7.3
Bw -- 18 to 23 in	loamy sand	rapid	0.14 to 0.33 in	5.1 to 7.3
BC,C -- 23 to 80 in	sand	rapid	1.71 to 4.00 in	5.6 to 7.8

Map Unit Description (MN)

Benton County, Minnesota

D7C--Hubbard loamy sand, 6 to 12 percent slopes

Hubbard

Extent: 85 to 95 percent of the unit

Landform(s): hills on stream terraces, hills on outwash plains

Slope gradient: 6 to 12 percent

Parent material: sandy outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: excessively drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .15

Land capability, nonirrigated: 6s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,AB -- 0 to 12 in	loamy sand	rapid	0.94 to 1.42 in	5.1 to 7.3
Bw -- 12 to 33 in	coarse sand	rapid	0.64 to 1.49 in	5.1 to 7.3
C -- 33 to 80 in	coarse sand	rapid	1.41 to 3.28 in	5.6 to 7.8

D7E--Hubbard loamy sand, 18 to 35 percent slopes

Hubbard

Extent: 75 to 95 percent of the unit

Landform(s): hills on stream terraces, hills on outwash plains

Slope gradient: 18 to 35 percent

Parent material: sandy outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: excessively drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .15

Land capability, nonirrigated: 7s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A,AB -- 0 to 12 in	loamy sand	rapid	0.94 to 1.42 in	5.1 to 7.3
Bw -- 12 to 33 in	coarse sand	rapid	0.64 to 1.49 in	5.1 to 7.3
C -- 33 to 80 in	coarse sand	rapid	1.41 to 3.28 in	5.6 to 7.8

Map Unit Description (MN)

Benton County, Minnesota

D8E--Sandberg loamy coarse sand, 18 to 35 percent slopes

Sandberg

<p><i>Extent:</i> 70 to 90 percent of the unit</p> <p><i>Landform(s):</i> escarpments on stream terraces, hills on stream terraces</p> <p><i>Slope gradient:</i> 18 to 35 percent</p> <p><i>Parent material:</i> sandy and gravelly outwash</p> <p><i>Restrictive feature(s):</i> greater than 60 inches</p> <p><i>Flooding:</i> none</p> <p><i>Ponding:</i> none</p> <p><i>Drainage class:</i> excessively drained</p>	<p><i>Soil loss tolerance (T factor):</i> 5</p> <p><i>Wind erodibility group (WEG):</i> 2</p> <p><i>Wind erodibility index (WEI):</i> 134</p> <p><i>Kw factor (surface layer):</i> .05</p> <p><i>Land capability, nonirrigated:</i> 7s</p> <p><i>Hydric soil:</i> no</p> <p><i>Hydrologic group:</i> A</p> <p><i>Potential for frost action:</i> low</p>
-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 11 in	loamy coarse sand	rapid	1.10 to 1.32 in	5.6 to 7.8
Bw -- 11 to 27 in	coarse sand	rapid	0.48 to 1.61 in	6.1 to 7.8
C -- 27 to 80 in	gravelly coarse sand	very rapid	1.06 to 3.17 in	7.4 to 8.4

Map Unit Description (MN)

Benton County, Minnesota

D9B--Stonelake-Sanburn complex, 1 to 6 percent slopes

Stonelake

Extent: 50 to 75 percent of the unit

Landform(s): moraines, outwash plains

Slope gradient: 1 to 6 percent

Parent material: gravelly outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: excessively drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .05

Land capability, nonirrigated: 6s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 4 in	gravelly loamy sand	rapid	0.20 to 0.35 in	5.1 to 6.5
Bw -- 4 to 11 in	gravelly coarse sand	very rapid	0.07 to 0.64 in	5.1 to 6.5
Bt -- 11 to 24 in	very gravelly coarse sand	very rapid	0.13 to 0.91 in	5.1 to 6.5
BC,C -- 24 to 80 in	gravelly sand	very rapid	0.56 to 3.35 in	5.1 to 7.3

Sanburn

Extent: 25 to 45 percent of the unit

Landform(s): moraines, outwash plains

Slope gradient: 1 to 4 percent

Parent material: loamy glaciofluvial deposits over sandy and gravelly outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat excessively drained

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .24

Land capability, nonirrigated: 4s

Hydric soil: no

Hydrologic group: A

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 5 in	sandy loam	moderately rapid	0.61 to 0.77 in	5.1 to 6.5
Bt -- 5 to 20 in	gravelly sandy loam	moderately rapid	1.05 to 1.80 in	5.1 to 6.5
2BC,2C -- 20 to 80 in	gravelly coarse sand	very rapid	1.20 to 2.39 in	5.1 to 6.5

Map Unit Description (MN)

Benton County, Minnesota

D9C--Stonelake-Sanburn complex, 6 to 15 percent slopes

Stonelake

Extent: 55 to 80 percent of the unit

Landform(s): moraines, outwash plains

Slope gradient: 6 to 15 percent

Parent material: gravelly outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: excessively drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .02

Land capability, nonirrigated: 6s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 4 in	gravelly loamy sand	rapid	0.20 to 0.35 in	5.1 to 6.5
Bw -- 4 to 11 in	gravelly coarse sand	very rapid	0.07 to 0.64 in	5.1 to 6.5
Bt -- 11 to 24 in	very gravelly coarse sand	very rapid	0.13 to 0.91 in	5.1 to 6.5
BC,C -- 24 to 80 in	gravelly sand	very rapid	0.56 to 3.35 in	5.1 to 7.3

Sanburn

Extent: 20 to 40 percent of the unit

Landform(s): moraines, outwash plains

Slope gradient: 6 to 12 percent

Parent material: loamy glaciofluvial deposits over sandy and gravelly outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat excessively drained

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .17

Land capability, nonirrigated: 4s

Hydric soil: no

Hydrologic group: A

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 5 in	sandy loam	moderately rapid	0.61 to 0.77 in	5.1 to 6.5
Bt -- 5 to 20 in	gravelly sandy loam	moderately rapid	1.05 to 1.80 in	5.1 to 6.5
2BC,2C -- 20 to 80 in	gravelly coarse sand	very rapid	1.20 to 2.39 in	5.1 to 6.5

Map Unit Description (MN)

Benton County, Minnesota

D9E--Stonelake-Sanburn complex, 15 to 40 percent slopes

Stonelake

Extent: 60 to 85 percent of the unit

Landform(s): moraines, outwash plains

Slope gradient: 15 to 40 percent

Parent material: gravelly outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: excessively drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .05

Land capability, nonirrigated: 7s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 2 in	loamy coarse sand	rapid	0.12 to 0.30 in	5.1 to 6.5
E -- 2 to 8 in	very gravelly loamy coarse sand	very rapid	0.18 to 0.24 in	5.1 to 6.5
Bt -- 8 to 16 in	very gravelly coarse sand	very rapid	0.25 to 0.66 in	5.1 to 6.5
C -- 16 to 80 in	gravelly coarse sand	very rapid	1.28 to 3.19 in	5.1 to 7.8

Sanburn

Extent: 15 to 30 percent of the unit

Landform(s): moraines, outwash plains

Slope gradient: 15 to 25 percent

Parent material: loamy glaciofluvial deposits over sandy and gravelly outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat excessively drained

Soil loss tolerance (T factor): 2

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .17

Land capability, nonirrigated: 6e

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 5 in	sandy loam	moderately rapid	0.61 to 0.77 in	5.1 to 6.5
Bt -- 5 to 14 in	sandy loam	moderately rapid	0.63 to 1.09 in	5.1 to 6.5
2C -- 14 to 80 in	coarse sand	rapid	1.31 to 2.63 in	5.1 to 6.5

Map Unit Description (MN)

Benton County, Minnesota

D14B--Elkriver-Mosford complex, 0 to 6 percent slopes, rarely flooded

Elkriver, rarely flooded

Extent: 60 to 80 percent of the unit

Landform(s): flats on benches on flood plains

Slope gradient: 1 to 6 percent

Parent material: alluvium

Restrictive feature(s): greater than 60 inches

Flooding: rare

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .28

Land capability, nonirrigated: 2s

Hydric soil: no

Hydrologic group: B

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 10 in	fine sandy loam	moderately rapid	1.57 to 1.97 in	5.1 to 7.3
A1,A2,A3 -- 10 to 35 in	fine sandy loam	moderately rapid	3.78 to 5.04 in	5.1 to 7.3
Bw -- 35 to 39 in	fine sandy loam	moderately rapid	0.59 to 0.75 in	5.6 to 7.8
2C -- 39 to 80 in	sand	rapid	0.82 to 4.09 in	5.6 to 7.8

Mosford, rarely flooded

Extent: 25 to 35 percent of the unit

Landform(s): rises on benches on flood plains

Slope gradient: 0 to 2 percent

Parent material: alluvium

Restrictive feature(s): greater than 60 inches

Flooding: rare

Ponding: none

Drainage class: somewhat excessively drained

Soil loss tolerance (T factor): 2

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .28

Land capability, nonirrigated: 3s

Hydric soil: no

Hydrologic group: B

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 11 in	fine sandy loam	moderate	1.43 to 1.98 in	5.1 to 7.3
Bw -- 11 to 16 in	fine sandy loam	moderate	0.46 to 0.87 in	5.1 to 7.3
2BC,2C1 -- 16 to 57 in	fine sand	rapid	2.05 to 2.87 in	5.1 to 7.3
2C2 -- 57 to 80 in	gravelly sand	very rapid	0.46 to 1.60 in	5.1 to 7.8

Map Unit Description (MN)

Benton County, Minnesota

D17A--Duelm loamy sand, 0 to 2 percent slopes

Duelm

Extent: 75 to 95 percent of the unit

Landform(s): outwash plains, stream terraces

Slope gradient: 0 to 2 percent

Parent material: sandy outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: moderately well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .15

Land capability, nonirrigated: 4s

Hydric soil: no

Hydrologic group: A/D

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,AB -- 0 to 16 in	loamy sand	rapid	1.29 to 1.94 in	5.6 to 7.3
Bw -- 16 to 30 in	coarse sand	rapid	0.83 to 1.52 in	5.1 to 7.3
C -- 30 to 80 in	coarse sand	rapid	1.00 to 3.50 in	5.6 to 7.8

D20A--Isan sandy loam, 0 to 2 percent slopes

Isan

Extent: 80 to 95 percent of the unit

Landform(s): outwash plains, stream terraces

Slope gradient: 0 to 2 percent

Parent material: sandy outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 2

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .20

Land capability, nonirrigated: 4w

Hydric soil: yes

Hydrologic group: A/D

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 14 in	sandy loam	moderately rapid	1.42 to 2.13 in	5.6 to 7.3
AB,Bg -- 14 to 34 in	loamy sand	rapid	1.18 to 1.97 in	5.1 to 6.5
Cg -- 34 to 80 in	coarse sand	rapid	1.84 to 2.76 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

D21A--Isan sandy loam, depressional, 0 to 1 percent slopes

Isan, depressional

Extent: 70 to 90 percent of the unit

Landform(s): depressions on outwash plains, depressions on stream terraces

Slope gradient: 0 to 1 percent

Parent material: sandy outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: frequent

Drainage class: very poorly drained

Soil loss tolerance (T factor): 2

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .20

Land capability, nonirrigated: 6w

Hydric soil: yes

Hydrologic group: A/D

Potential for frost action: moderate

Representative soil profile:

	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 14 in	sandy loam	moderately rapid	1.42 to 2.13 in	5.6 to 7.3
AB,Bg -- 14 to 34 in	loamy sand	rapid	1.18 to 1.97 in	5.1 to 6.5
Cg -- 34 to 80 in	coarse sand	rapid	1.84 to 2.76 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

D22A--Hubbard-Mosford complex, 0 to 3 percent slopes

Hubbard

Extent: 60 to 85 percent of the unit

Landform(s): stream terraces, outwash plains

Slope gradient: 0 to 3 percent

Parent material: sandy outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: excessively drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .10

Land capability, nonirrigated: 4s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,AB -- 0 to 20 in	loamy sand	rapid	1.61 to 2.41 in	5.1 to 7.3
Bw -- 20 to 32 in	loamy sand	rapid	0.35 to 0.83 in	5.1 to 7.3
BC,C -- 32 to 80 in	sand	rapid	1.44 to 3.36 in	5.6 to 7.8

Mosford

Extent: 15 to 40 percent of the unit

Landform(s): swales on stream terraces, swales on outwash plains

Slope gradient: 0 to 2 percent

Parent material: loamy alluvium over sandy outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat excessively drained

Soil loss tolerance (T factor): 2

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .20

Land capability, nonirrigated: 3s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 13 in	sandy loam	moderately rapid	1.69 to 2.34 in	5.1 to 7.3
Bw -- 13 to 16 in	coarse sandy loam	moderately rapid	0.38 to 0.54 in	5.1 to 7.3
2Bw -- 16 to 35 in	coarse sand	rapid	0.57 to 2.08 in	5.1 to 7.3
2C -- 35 to 80 in	sand	rapid	0.90 to 3.14 in	5.1 to 7.8

Map Unit Description (MN)

Benton County, Minnesota

D30A--Seelyeville and Markey soils, depressional, 0 to 1 percent slopes

Markey, depressional

<p><i>Extent:</i> 0 to 100 percent of the unit</p> <p><i>Landform(s):</i> lake plains, outwash plains</p> <p><i>Slope gradient:</i> 0 to 1 percent</p> <p><i>Parent material:</i> muck herbaceous organic material over sandy outwash</p> <p><i>Restrictive feature(s):</i> greater than 60 inches</p> <p><i>Flooding:</i> none</p> <p><i>Ponding:</i> occasional</p> <p><i>Drainage class:</i> very poorly drained</p>	<p><i>Soil loss tolerance (T factor):</i> 1</p> <p><i>Wind erodibility group (WEG):</i> 2</p> <p><i>Wind erodibility index (WEI):</i> 134</p> <p><i>Kw factor (surface layer)</i> .02</p> <p><i>Land capability, nonirrigated:</i> 7w</p> <p><i>Hydric soil:</i> yes</p> <p><i>Hydrologic group:</i> A/D</p> <p><i>Potential for frost action:</i> high</p>
-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Oa -- 0 to 36 in	muck	moderately rapid	12.54 to 16.12 in	
A -- 36 to 42 in	loamy sand	rapid	0.38 to 0.94 in	5.6 to 7.3
Cg -- 42 to 80 in	sand	rapid	1.13 to 3.02 in	5.6 to 7.3

Seelyeville, depressional

<p><i>Extent:</i> 0 to 100 percent of the unit</p> <p><i>Landform(s):</i> lake plains, outwash plains</p> <p><i>Slope gradient:</i> 0 to 1 percent</p> <p><i>Parent material:</i> highly decomposed organic material</p> <p><i>Restrictive feature(s):</i> greater than 60 inches</p> <p><i>Flooding:</i> none</p> <p><i>Ponding:</i> occasional</p> <p><i>Drainage class:</i> very poorly drained</p>	<p><i>Soil loss tolerance (T factor):</i> 2</p> <p><i>Wind erodibility group (WEG):</i> 2</p> <p><i>Wind erodibility index (WEI):</i> 134</p> <p><i>Kw factor (surface layer)</i> .02</p> <p><i>Land capability, nonirrigated:</i> 7w</p> <p><i>Hydric soil:</i> yes</p> <p><i>Hydrologic group:</i> A/D</p> <p><i>Potential for frost action:</i> high</p>
--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Oa1 -- 0 to 10 in	muck	moderately rapid	3.44 to 4.43 in	
Oa2,Oa3 -- 10 to 80 in	muck	moderately rapid	24.53 to 31.54 in	

Map Unit Description (MN)

Benton County, Minnesota

D32A--Mosford sandy loam, 0 to 2 percent slopes

Mosford

<i>Extent:</i> 90 to 100 percent of the unit	<i>Soil loss tolerance (T factor):</i> 2
<i>Landform(s):</i> swales on stream terraces, swales on outwash plains	<i>Wind erodibility group (WEG):</i> 3
<i>Slope gradient:</i> 0 to 2 percent	<i>Wind erodibility index (WEI):</i> 86
<i>Parent material:</i> loamy alluvium over sandy outwash	<i>Kw factor (surface layer)</i> .20
<i>Restrictive feature(s):</i> greater than 60 inches	<i>Land capability, nonirrigated:</i> 3s
<i>Flooding:</i> none	<i>Hydric soil:</i> no
<i>Ponding:</i> none	<i>Hydrologic group:</i> A
<i>Drainage class:</i> somewhat excessively drained	<i>Potential for frost action:</i> low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 13 in	sandy loam	moderately rapid	1.69 to 2.34 in	5.1 to 7.3
Bw -- 13 to 16 in	coarse sandy loam	moderately rapid	0.38 to 0.54 in	5.1 to 7.3
2Bw -- 16 to 35 in	coarse sand	rapid	0.57 to 2.08 in	5.1 to 7.3
2C -- 35 to 80 in	sand	rapid	0.90 to 3.14 in	5.1 to 7.8

Map Unit Description (MN)

Benton County, Minnesota

D36B--Eagleview loamy sand, 2 to 6 percent slopes

Eagleview

Extent: 80 to 100 percent of the unit

Landform(s): moraines, outwash plains, stream terraces

Slope gradient: 2 to 6 percent

Parent material: sandy outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat excessively drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .20

Land capability, nonirrigated: 4s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 9 in	loamy sand	rapid	0.91 to 1.09 in	5.6 to 7.3
E -- 9 to 36 in	sand	rapid	2.41 to 2.94 in	5.6 to 7.3
E&Bt -- 36 to 54 in	loamy sand	rapid	1.09 to 1.45 in	6.1 to 7.3
C -- 54 to 80 in	sand	rapid	1.30 to 1.82 in	6.1 to 8.4

Map Unit Description (MN)

Benton County, Minnesota

D36C--Eagleview loamy sand, 6 to 12 percent slopes

Eagleview

Extent: 80 to 100 percent of the unit

Landform(s): moraines, outwash plains, stream terraces

Slope gradient: 6 to 12 percent

Parent material: sandy outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat excessively drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .15

Land capability, nonirrigated: 4s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 3 in	loamy sand	rapid	0.31 to 0.38 in	5.6 to 7.3
E -- 3 to 37 in	sand	rapid	3.05 to 3.72 in	5.6 to 7.3
E&Bt -- 37 to 60 in	stratified sand to loamy sand	rapid	1.37 to 1.83 in	6.1 to 7.3
C -- 60 to 80 in	sand	rapid	1.00 to 1.41 in	6.1 to 8.4

D38A--Cantlin loamy fine sand, 0 to 3 percent slopes

Cantlin

Extent: 75 to 95 percent of the unit

Landform(s): lake plains, outwash plains

Slope gradient: 0 to 3 percent

Parent material: sandy glaciofluvial deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .20

Land capability, nonirrigated: 4s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 8 in	loamy fine sand	rapid	0.71 to 0.87 in	5.1 to 6.5
Bw -- 8 to 22 in	loamy fine sand	rapid	0.71 to 1.42 in	5.1 to 7.3
BC,C -- 22 to 80 in	fine sand	rapid	2.89 to 4.05 in	5.1 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

D39A--Fordum loam, 0 to 2 percent slopes, occasionally flooded

Fordum, occasionally flooded

Extent: 85 to 95 percent of the unit

Landform(s): flood plains

Slope gradient: 0 to 2 percent

Parent material: loamy alluvium over sandy and gravelly alluvium

Restrictive feature(s): greater than 60 inches

Flooding: occasional

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 3

Wind erodibility group (WEG): 5

Wind erodibility index (WEI): 56

Kw factor (surface layer) .32

Land capability, nonirrigated: 4w

Hydric soil: yes

Hydrologic group: B/D

Potential for frost action: high

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 9 in	loam	moderate	1.36 to 1.99 in	5.1 to 7.3
Cg -- 9 to 38 in	loam	moderate	2.91 to 6.41 in	5.1 to 7.3
2Cg -- 38 to 80 in	stratified sand to silt loam	rapid	2.09 to 8.35 in	5.6 to 7.3

D44A--Isanti loamy fine sand, 0 to 2 percent slopes

Isanti

Extent: 80 to 90 percent of the unit

Landform(s): lake plains, outwash plains

Slope gradient: 0 to 2 percent

Parent material: sandy glaciofluvial deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .28

Land capability, nonirrigated: 4w

Hydric soil: yes

Hydrologic group: A/D

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 12 in	loamy fine sand	rapid	0.94 to 1.30 in	5.1 to 6.5
Bg -- 12 to 36 in	loamy sand	rapid	1.44 to 3.60 in	5.6 to 7.3
Cg -- 36 to 80 in	sand	rapid	1.32 to 3.53 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

D46A--Lino loamy fine sand, 0 to 2 percent slopes

Lino

Extent: 80 to 90 percent of the unit

Landform(s): lake plains, outwash plains

Slope gradient: 0 to 2 percent

Parent material: sandy glaciofluvial deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .32

Land capability, nonirrigated: 4s

Hydric soil: no

Hydrologic group: A/D

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 7 in	loamy fine sand	rapid	0.64 to 0.78 in	5.1 to 6.5
Bw -- 7 to 45 in	fine sand	rapid	2.27 to 4.16 in	5.1 to 7.3
C -- 45 to 60 in	fine sand	rapid	0.75 to 1.05 in	5.1 to 7.3

D47A--Kost loamy fine sand, 0 to 2 percent slopes

Kost

Extent: 85 to 100 percent of the unit

Landform(s): outwash plains

Slope gradient: 0 to 2 percent

Parent material: sandy glaciofluvial deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: excessively drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .17

Land capability, nonirrigated: 4s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 16 in	loamy fine sand	rapid	1.45 to 1.94 in	5.1 to 7.3
Bw -- 16 to 34 in	fine sand	rapid	1.06 to 1.42 in	5.1 to 7.3
C -- 34 to 60 in	fine sand	rapid	1.30 to 1.82 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

D47B--Kost loamy fine sand, 2 to 6 percent slopes

Kost

Extent: 85 to 100 percent of the unit

Landform(s): outwash plains

Slope gradient: 2 to 6 percent

Parent material: sandy glaciofluvial deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: excessively drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .17

Land capability, nonirrigated: 4s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 14 in	loamy fine sand	rapid	1.28 to 1.70 in	5.1 to 7.3
Bw -- 14 to 33 in	fine sand	rapid	1.13 to 1.51 in	5.1 to 7.3
C -- 33 to 60 in	fine sand	rapid	1.34 to 1.87 in	5.6 to 7.3

D47C--Kost loamy fine sand, 6 to 12 percent slopes

Kost

Extent: 90 to 100 percent of the unit

Landform(s): outwash plains

Slope gradient: 6 to 12 percent

Parent material: sandy glaciofluvial deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: excessively drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .17

Land capability, nonirrigated: 4s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 14 in	loamy fine sand	rapid	1.28 to 1.70 in	5.1 to 7.3
Bw -- 14 to 33 in	fine sand	rapid	1.13 to 1.51 in	5.1 to 7.3
C -- 33 to 60 in	fine sand	rapid	1.34 to 1.87 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

D48A--Cantlin loamy fine sand, thick surface, 0 to 2 percent slopes

Cantlin, thick surface

Extent: 85 to 95 percent of the unit

Landform(s): outwash plains

Slope gradient: 0 to 2 percent

Parent material: sandy glaciofluvial deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: well drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .17

Land capability, nonirrigated: 4s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 14 in	loamy fine sand	rapid	1.28 to 1.56 in	5.1 to 6.5
Bw -- 14 to 30 in	loamy fine sand	rapid	0.79 to 1.57 in	5.1 to 7.3
BC,C -- 30 to 80 in	fine sand	rapid	2.50 to 3.50 in	5.1 to 7.3

D50A--Isanti fine sandy loam, depressional, 0 to 1 percent slopes

Isanti, depressional

Extent: 80 to 90 percent of the unit

Landform(s): depressions on outwash plains

Slope gradient: 0 to 1 percent

Parent material: sandy glaciofluvial deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: frequent

Drainage class: very poorly drained

Soil loss tolerance (T factor): 2

Wind erodibility group (WEG): 3

Wind erodibility index (WEI): 86

Kw factor (surface layer) .28

Land capability, nonirrigated: 6w

Hydric soil: yes

Hydrologic group: A/D

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 16 in	fine sandy loam	moderately rapid	1.94 to 2.42 in	5.1 to 6.5
Bg -- 16 to 28 in	loamy fine sand	rapid	0.71 to 1.30 in	5.1 to 7.3
Cg -- 28 to 80 in	fine sand	rapid	2.60 to 3.64 in	5.6 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

D51A--Kost loamy fine sand, banded substratum, 0 to 2 percent slopes

Kost, banded substratum

Extent: 85 to 100 percent of the unit

Landform(s): outwash plains

Slope gradient: 0 to 2 percent

Parent material: sandy glaciofluvial deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: excessively drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .17

Land capability, nonirrigated: 4s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 16 in	loamy fine sand	rapid	1.45 to 1.94 in	5.1 to 7.3
Bw -- 16 to 25 in	fine sand	rapid	0.54 to 0.72 in	5.1 to 7.3
E&Bt -- 25 to 80 in	fine sand	rapid	2.74 to 5.47 in	5.1 to 7.3

D51B--Kost loamy fine sand, banded substratum, 2 to 6 percent slopes

Kost, banded substratum

Extent: 85 to 100 percent of the unit

Landform(s): outwash plains

Slope gradient: 2 to 6 percent

Parent material: sandy glaciofluvial deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: excessively drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .17

Land capability, nonirrigated: 4s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 14 in	loamy fine sand	rapid	1.28 to 1.70 in	5.1 to 7.3
Bw -- 14 to 25 in	fine sand	rapid	0.66 to 0.88 in	5.1 to 7.3
E&Bt -- 25 to 80 in	fine sand	rapid	2.74 to 5.47 in	5.1 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

D51C--Kost loamy fine sand, banded substratum, 6 to 12 percent slopes

Kost, banded substratum

Extent: 85 to 100 percent of the unit

Landform(s): outwash plains

Slope gradient: 6 to 12 percent

Parent material: sandy glaciofluvial deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: excessively drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .17

Land capability, nonirrigated: 4s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 14 in	loamy fine sand	rapid	1.28 to 1.70 in	5.1 to 7.3
Bw -- 14 to 25 in	fine sand	rapid	0.66 to 0.88 in	5.1 to 7.3
E&Bt -- 25 to 80 in	fine sand	rapid	2.74 to 5.47 in	5.1 to 7.3

D52A--Glendorado loamy fine sand, 0 to 2 percent slopes

Glendorado

Extent: 75 to 90 percent of the unit

Landform(s): outwash plains

Slope gradient: 0 to 2 percent

Parent material: sandy glaciofluvial deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .17

Land capability, nonirrigated: 4s

Hydric soil: no

Hydrologic group: A/D

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 15 in	loamy fine sand	rapid	1.35 to 1.80 in	5.1 to 7.3
Bw1 -- 15 to 22 in	fine sand	rapid	0.43 to 0.57 in	5.1 to 7.3
Bw2 -- 22 to 48 in	sand	rapid	1.56 to 2.86 in	5.1 to 7.3
C -- 48 to 80 in	sand	rapid	0.64 to 2.23 in	5.6 to 7.8

Map Unit Description (MN)

Benton County, Minnesota

D54B--Hubbard, bedrock substratum-Rock outcrop complex, 1 to 8 percent slopes

Hubbard, bedrock substratum

Extent: 60 to 80 percent of the unit

Landform(s): stream terraces

Slope gradient: 1 to 8 percent

Parent material: sandy outwash over bedrock

Restrictive feature(s): lithic bedrock at 40 to 80 inches

Flooding: none

Ponding: none

Drainage class: excessively drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .10

Land capability, nonirrigated: 4s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 18 in	loamy sand	rapid	1.45 to 2.17 in	5.1 to 7.3
Bw -- 18 to 23 in	loamy sand	rapid	0.14 to 0.33 in	5.1 to 7.3
BC,C -- 23 to 60 in	sand	rapid	1.11 to 2.59 in	5.6 to 7.8
2R -- 60 to 80 in	unweathered bedrock	impermeable		

Rock outcrop

Extent: 15 to 25 percent of the unit

Landform(s): stream terraces

Slope gradient: 1 to 8 percent

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated:

Hydric soil:

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>

Map Unit Description (MN)

Benton County, Minnesota

D55B--Zimmerman fine sand, banded substratum, 1 to 6 percent slopes

Zimmerman, banded substratum

Extent: 80 to 90 percent of the unit

Landform(s): outwash plains

Slope gradient: 1 to 6 percent

Parent material: sandy glaciofluvial deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: excessively drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 1

Wind erodibility index (WEI): 250

Kw factor (surface layer) .10

Land capability, nonirrigated: 4s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 7 in	fine sand	rapid	0.43 to 0.57 in	5.1 to 6.5
Bw -- 7 to 63 in	fine sand	rapid	3.35 to 6.15 in	5.1 to 7.3
E&Bt -- 63 to 80 in	fine sand	rapid	0.85 to 1.69 in	5.1 to 7.3

D55C--Zimmerman fine sand, banded substratum, 6 to 12 percent slopes

Zimmerman, banded substratum

Extent: 80 to 95 percent of the unit

Landform(s): outwash plains

Slope gradient: 6 to 12 percent

Parent material: sandy glaciofluvial deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: excessively drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 1

Wind erodibility index (WEI): 250

Kw factor (surface layer) .10

Land capability, nonirrigated: 4s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 7 in	fine sand	rapid	0.43 to 0.57 in	5.1 to 6.5
Bw -- 7 to 63 in	fine sand	rapid	3.35 to 6.15 in	5.1 to 7.3
E&Bt -- 63 to 80 in	fine sand	rapid	0.85 to 1.69 in	5.1 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

D55E--Zimmerman fine sand, banded substratum, 12 to 35 percent slopes

Zimmerman, banded substratum

Extent: 80 to 90 percent of the unit

Landform(s): outwash plains

Slope gradient: 12 to 35 percent

Parent material: sandy glaciofluvial deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: excessively drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 1

Wind erodibility index (WEI): 250

Kw factor (surface layer) .10

Land capability, nonirrigated: 6e

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 3 in	fine sand	rapid	0.22 to 0.28 in	5.1 to 6.5
Bw -- 3 to 63 in	fine sand	rapid	3.59 to 6.58 in	5.1 to 7.3
E&Bt -- 63 to 80 in	fine sand	rapid	0.85 to 1.69 in	5.1 to 7.3

D56A--Lino loamy fine sand, stratified substratum, 0 to 2 percent slopes

Lino, stratified substratum

Extent: 70 to 90 percent of the unit

Landform(s): lake plains, outwash plains

Slope gradient: 0 to 2 percent

Parent material: sandy glaciofluvial deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .32

Land capability, nonirrigated: 4s

Hydric soil: no

Hydrologic group: A/D

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 7 in	loamy fine sand	rapid	0.64 to 0.78 in	5.1 to 6.5
Bw -- 7 to 65 in	loamy fine sand	rapid	3.47 to 6.37 in	5.1 to 7.3
2C -- 65 to 80 in	stratified loamy very fine sand to silty clay loam	rapid	1.20 to 3.29 in	5.1 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

D61A--Glendorado loamy sand, 0 to 2 percent slopes

Glendorado

Extent: 80 to 95 percent of the unit

Landform(s): outwash plains, stream terraces

Slope gradient: 0 to 2 percent

Parent material: sandy outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: somewhat poorly drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 2

Wind erodibility index (WEI): 134

Kw factor (surface layer) .15

Land capability, nonirrigated: 4s

Hydric soil: no

Hydrologic group: A/D

Potential for frost action: moderate

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap,A -- 0 to 15 in	loamy sand	rapid	1.20 to 1.80 in	5.6 to 7.3
Bw1 -- 15 to 22 in	loamy sand	rapid	0.43 to 0.78 in	5.1 to 7.3
Bw2 -- 22 to 48 in	sand	rapid	1.56 to 2.86 in	5.1 to 7.3
C -- 48 to 80 in	sand	rapid	0.64 to 2.23 in	5.6 to 7.8

D64A--Zimmerman fine sand, 0 to 3 percent slopes

Zimmerman

Extent: 80 to 95 percent of the unit

Landform(s): outwash plains

Slope gradient: 0 to 3 percent

Parent material: sandy glaciofluvial deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: excessively drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 1

Wind erodibility index (WEI): 250

Kw factor (surface layer) .10

Land capability, nonirrigated: 4s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 7 in	fine sand	rapid	0.43 to 0.57 in	5.1 to 6.5
Bw -- 7 to 27 in	fine sand	rapid	1.20 to 2.21 in	5.1 to 7.3
E&Bt -- 27 to 80 in	fine sand	rapid	2.64 to 5.28 in	5.1 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

D64B--Zimmerman fine sand, 3 to 6 percent slopes

Zimmerman

Extent: 95 to 100 percent of the unit

Landform(s): outwash plains

Slope gradient: 3 to 6 percent

Parent material: sandy glaciofluvial deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: excessively drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 1

Wind erodibility index (WEI): 250

Kw factor (surface layer) .10

Land capability, nonirrigated: 4s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 7 in	fine sand	rapid	0.43 to 0.57 in	5.1 to 6.5
Bw -- 7 to 27 in	fine sand	rapid	1.20 to 2.21 in	5.1 to 7.3
E&Bt -- 27 to 80 in	fine sand	rapid	2.64 to 5.28 in	5.1 to 7.3

D64C--Zimmerman fine sand, 6 to 12 percent slopes

Zimmerman

Extent: 90 to 100 percent of the unit

Landform(s): outwash plains

Slope gradient: 6 to 12 percent

Parent material: sandy glaciofluvial deposits

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: excessively drained

Soil loss tolerance (T factor): 5

Wind erodibility group (WEG): 1

Wind erodibility index (WEI): 250

Kw factor (surface layer) .10

Land capability, nonirrigated: 4s

Hydric soil: no

Hydrologic group: A

Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
Ap -- 0 to 7 in	fine sand	rapid	0.43 to 0.57 in	5.1 to 6.5
Bw -- 7 to 27 in	fine sand	rapid	1.20 to 2.21 in	5.1 to 7.3
E&Bt -- 27 to 80 in	fine sand	rapid	2.64 to 5.28 in	5.1 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

D64E--Zimmerman fine sand, 12 to 30 percent slopes

Zimmerman

Extent: 90 to 100 percent of the unit
Landform(s): outwash plains
Slope gradient: 12 to 30 percent
Parent material: sandy glaciofluvial deposits
Restrictive feature(s): greater than 60 inches
Flooding: none
Ponding: none
Drainage class: excessively drained

Soil loss tolerance (T factor): 5
Wind erodibility group (WEG): 1
Wind erodibility index (WEI): 250
Kw factor (surface layer) .10
Land capability, nonirrigated: 6e
Hydric soil: no
Hydrologic group: A
Potential for frost action: low

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
A -- 0 to 3 in	fine sand	rapid	0.22 to 0.28 in	5.1 to 6.5
Bw -- 3 to 25 in	fine sand	rapid	1.32 to 2.43 in	5.1 to 7.3
E&Bt -- 25 to 80 in	fine sand	rapid	2.74 to 5.47 in	5.1 to 7.3

Map Unit Description (MN)

Benton County, Minnesota

GP--Pits, gravel-Udipsamments complex

Pits, gravel

Extent: 70 to 90 percent of the unit

Landform(s): moraines, stream terraces, outwash plains

Slope gradient: 0 to 45 percent

Parent material: outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated:

Hydric soil:

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Udipsamments

Extent: 10 to 30 percent of the unit

Landform(s): moraines, stream terraces, outwash plains

Slope gradient: 0 to 6 percent

Parent material: outwash

Restrictive feature(s): greater than 60 inches

Flooding: none

Ponding: none

Drainage class: excessively drained

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated:

Hydric soil:

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

Map Unit Description (MN)

Benton County, Minnesota

M-W--Water, miscellaneous

Water, miscellaneous

Extent: 100 percent of the unit

Landform(s):

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated:

Hydric soil:

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

W--Water

Water

Extent: 100 percent of the unit

Landform(s):

Slope gradient:

Parent material:

Restrictive feature(s): greater than 60 inches

Flooding:

Ponding:

Drainage class:

Soil loss tolerance (T factor):

Wind erodibility group (WEG):

Wind erodibility index (WEI):

Kw factor (surface layer)

Land capability, nonirrigated:

Hydric soil:

Hydrologic group:

Potential for frost action:

<i>Representative soil profile:</i>	<i>Texture</i>	<i>Permeability</i>	<i>Available water capacity</i>	<i>pH</i>
-------------------------------------	----------------	---------------------	---------------------------------	-----------

This report provides a semitabular listing of some soil and site properties and interpretations that are valuable in communicating the concept of a map unit. The report also provides easy access to the commonly used conservation planning information in one place. The major soil components in each map unit are displayed. Minor components may be displayed if they are included in the database and are selected at the time the report is generated.