

Ecological Site Description

Wet Terrace Forest

F115CY003MO


- (*Quercus macrocarpa* - *Quercus shumardii*/*Ilex decidua*/*Chasmanthium latifolium*)
- (bur oak – Shumard oak/possum haw/river oats)

An Ecological Site Description (ESD) is a reference document of ecological knowledge regarding a particular land area (ecological site). An ESD describes ecological potential and ecosystem dynamics of land areas and their potential management. Ecological sites are linked to soil survey map unit components, which allows for mapping of ecological sites. (**NOTE:** *This is a “provisional” ESD, and is subject to change. It contains basic ecological information sufficient for conservation planning and land management in Missouri. After additional information is developed and reviewed, a “Correlated” ESD will be published and will be available via the Web Soil Survey <http://websoilsurvey.nrcs.usda.gov> .)*

Major Land Resource Area: 115C – Central Mississippi Valley Wooded Slopes, Northern Part

Introduction

The Central Mississippi Valley Wooded Slopes, Northern Part (area outlined in red on the map) is centered on the dissected, loess-covered hills bordering the Mississippi and Illinois River floodplain and terrace systems, and associated tributaries. Local physiographic regions include the Galesburg




Plain and the Springfield Plain in Illinois, and the Lincoln Hills in Missouri. Elevation ranges from about 420 feet along the Mississippi River just upstream from St. Louis, to nearly 900 feet on the highest upland summits in Illinois. Local relief varies from 10 to 20 feet in the major river floodplains, to 50 to 100 feet in the dissected uplands, with bluffs of up to 250 feet along the Mississippi and Illinois Rivers. Loess caps both stream and glacial outwash terraces along the major rivers, as well as both Illinoian and Pre-Illinoian till near the edges of the area. The underlying Mississippian-aged limestone influences soils on lower, steeper slopes.

Wet Terrace Forests are within the green areas on the map (Missouri portion only; Illinois distributions are currently under review). They are in the southern part of the area, in Pike county, Missouri, along tributary streams of the Mississippi River. Soils are very deep with clayey subsoils, have a high water table in the spring months, and are subject to flooding.

Physiographic Features

This site is on low stream terraces with slopes of less than 3 percent. The site generates some runoff to adjacent lower floodplain sites, and receives some runoff from higher stream terraces and uplands. This site is subject to rare flooding. Scour is uncommon in these flood events, and deposition is minimal, so ecological processes more closely resemble those of stream terrace systems.

The adjacent figure (adapted from Love, 1997) shows the typical landscape position of this



ecological site, and landscape relationships among the major ecological sites in the floodplains, stream terraces, and adjacent uplands. The site is within the area labeled “3”, on higher positions than the active floodplain along the stream channel. More than one terrace step and soil may be present, as indicated by the dashed line within the Wet Terrace Forest on the diagram.

Soil Features

These soils have no rooting restriction. They were formed

under a mixture of herbaceous wetland and woodland vegetation. Organic matter content is variable. Parent material is alluvium. They have silt loam surface horizons, and clayey subsoils with argillic horizons. They are affected by a seasonal high water table during the spring months. Soil series associated with this site include Okaw.

Ecological Dynamics

Information contained in this section was developed using historical data, professional experience, field reviews, and scientific studies. The information presented is representative of very complex vegetation communities. Key indicator plants, animals and ecological processes are described to help inform land management decisions. Plant communities will differ across the MLRA because of the naturally occurring variability in weather, soils, and aspect. The Reference Plant Community is not necessarily the management goal. The species lists are representative and are not botanical descriptions of all species occurring, or potentially occurring, on this site. They are not intended to cover every situation or the full range of conditions, species, and responses for the site.

The reference plant community is dominated by a wide variety of deciduous hardwood tree species, tolerant of seasonally wet conditions including bur oak, Shumard oak, swamp white oak, American elm, black cherry. Trees are generally large and tall forming a dense, closed canopy. These forests are structurally and compositionally diverse, with occasional tree-fall gaps and natural mortality providing opportunities for regeneration of overstory species. The understory is also complex, with multiple layers of shade tolerant species such as blue beech, spicebush, and Ohio buckeye. Grape vines, greenbriar, and trumpet creeper are also present along with a diverse array of ground flora species that carpets the forest floor.

Today, the Wet Terrace Forests are largely converted to pasture and cropland. Where they do still occur, they often occur as a rather narrow band of forest traversing the riverfront forest or stream edge. These bands of forest play an important role as a source of food and shelter for migrating birds. In addition, they are very important in stream bank stabilization. Most sites have suffered from extensive hydrological alterations (ditches, levees, etc.). This ecological site can intermix with

Wet Terrace Prairie and Savannas. Uncontrolled grazing by domestic livestock in these remaining areas of forest damages and kills smaller trees and removes the ground cover. Carefully planned timber harvests can be tolerated on these sites, but high grading of the timber will ultimately degrade the sites. Re-establishment of these terrace forests is important for stream quality and stream health, and as critical habitat for migratory birds. Planting of later successional species on the appropriate landscape position and soils has proven to be an effective means for restoration.

Reference State Plant Community

Canopy Trees

Common Name	Botanical Name	Cover % (low-high)	Canopy Height (ft)
BUR OAK	<i>Quercus macrocarpa</i>	20-40	80
GREEN ASH	<i>Fraxinus pennsylvanica</i>	10-20	80
SWAMP WHITE OAK	<i>Quercus bicolor</i>	10-20	70
HACKBERRY	<i>Celtis occidentalis</i>	10-20	70
SHELLBARK HICKORY	<i>Carya lacinosa</i>	10-20	70
SLIPPERY ELM	<i>Ulmus rubra</i>	10-20	60
SYCAMORE	<i>Platanus occidentalis</i>	10-20	90
BLACK CHERRY	<i>Prunus serotina</i>	10-20	70
SHUMARD OAK	<i>Quercus shumardii</i>	20-40	90
PECAN	<i>Carya illinoensis</i>	10-20	70

Understory Trees

Common Name	Botanical Name	Cover % (low-high)	Canopy Height (ft)
OHIO BUCKEYE	<i>Aesculus glabra</i>	10-20	40
BLUE BEECH	<i>Carpinus caroliniana</i>	10-20	20

Shrubs

Common Name	Botanical Name	Cover % (low-high)	Canopy Height (ft)
POSSUM HAW	<i>Ilex decidua</i>	10-20	8
SPICE BUSH	<i>Lindera benzoin</i>	10-20	6

Vines

Common Name	Botanical Name	Cover % (low-high)
TRUMPET CREEPER	<i>Campis radicans</i>	10-20
RACCOON GRAPE	<i>Ampelopsis cordata</i>	10-20
FOX GRAPE	<i>Vitis vulpina</i>	10-20
GREENBRIAR	<i>Smilax glauca</i>	10-20

Forbs

Common Name	Botanical Name	Cover % (low-high)
PALE TOUCH-ME-NOT	<i>Impatiens pallida</i>	5-20
FALSE NETTLE	<i>Boehmeria cylindrica</i>	10-20
CLEARWEED	<i>Pilea pumila</i>	5-20
WHITE WOODLAND ASTER	<i>Aster lateriflorus</i>	5-20
WOOD NETTLE	<i>Laportea canadensis</i>	10-20
WATERLEAF	<i>Hydrophyllum virginianum</i>	5-20
HISPID BUTTERCUP	<i>Ranunculus hispidus</i>	5-20
SKULLCAP	<i>Scutellaria nervosa</i>	5-20

Grasses and sedges

Common Name	Botanical Name	Cover % (low-high)
RIVER OATS	<i>Chasmanthium latifolium</i>	10-20
HOP SEDGE	<i>Carex lupulina</i>	5-10
WOOD REED GRASS	<i>Cinna arundinacea</i>	5-10

Site Interpretations

Wildlife

- Moist conditions with abundant coarse woody debris make this type of ecological site important for many herptiles.
- Ephemeral pools provide important amphibian breeding habitat. Periodic inundation and acorns provide important habitat and food for migrating ducks (especially mallards) and breeding ducks including wood ducks and hooded mergansers.
- Tall emergent trees along with an uneven canopy structure and canopy gaps are important for heron colonies, eagle nesting, Mississippi kites, cerulean warblers and other bird species.
- Birds associated with late-successional to mature forests are Wood Duck, Hooded Merganser, Barred Owl, Cerulean Warbler, Yellow-throated Warbler, Prothonotary Warbler, Pileated Woodpecker, Yellow-throated Vireo, Brown Creeper, and Yellow-crowned Night Heron.
- Reptiles and amphibians associated with ecological site include: small-mouthed salamander, central newt, midland brown snake, gray treefrog, northern spring peeper, Blanchard’s cricket frog, southern leopard frog, western painted turtle, and red-eared slider.

Forestry

- **Management:** Estimated site index values range from 50 to 70. On the wettest sites, timber management opportunities may be limited. Management of these groups is often difficult because of the great variation in species, age, stocking levels and seasonal wetness. Use seed-tree, group selection, or clear cutting regeneration methods. Harvest favoring reproduction of the less-shade tolerant species such as bur oak, sycamore, and cottonwood. Maintain adequate riparian buffer areas.
- **Limitations:** Wetness from flooding; high water table. Use of equipment may be restricted in spring and other excessively wet periods. Restrict activities to dry periods or surfaced areas. Equipment use when wet may compact soil and damage tree roots. Unsurfaced roads and traffic areas tend to be slippery and form ruts easily. Access to forests is easiest during periods in late summer or winter when soils are frozen or dry. Planting is extremely difficult during spring periods. Seedling mortality may be high due to excess wetness. Unsurfaced roads and skid trails may be impassable during rainy periods.

Glossary

Backslope – a hillslope profile position that forms the steepest and generally linear, middle portion of the slope.

Backswamp – marshy or swampy, depressed areas of flood plains between natural levees and valley sides or terraces

Calcareous – the presence of calcium carbonate in the soil parent material within the rooting zone; relatively alkaline

Claypan – a dense, compact, slowly permeable layer in the subsoil having much higher clay content than the overlying material

Chert – hard, extremely dense or compact crystalline sedimentary rock, consisting dominantly of interlocking crystals of quartz

Cliff – a significant vertical, or near vertical, rock exposure

Dolomite – a type of sedimentary rock that is a carbonate mineral composed of calcium magnesium carbonate

Drainageway – the upper most reach of a stream channel system characterized by little meandering

Dry – a site where soil moisture is limiting during the growing season; low available water capacity

Dune – a low mound, ridge, bank or hill of loose, wind-blown sand

Exposed – steep, south and west-facing slopes, which are warmer and drier than other slope aspects

Flatwoods – a type of woodland that occurs on soils with a root restricting subsoil layer within 20 to 30 inches, resulting in very slow runoff and ponding that remains saturated for most of the winter and early spring months but dries out and becomes very dry in the summer months; plants that grow there must be adapted to both conditions

Floodplain – the nearly level plain that borders a stream and is subject to inundation under flood-stage conditions

Footslope – a hillslope position at the base of a slope where hillslope sediment (colluvium) accumulates

Forest – a vegetative community dominated by trees forming a closed canopy and interspersed with shade-tolerant understory species

Fragipan – a dense, brittle subsoil horizon that is extremely hard and compact when dry

Glade – open, rocky, barren vegetative community dominated by drought-adapted forbs and grasses, typically with scattered, stunted woody plants

Igneous –bedrock formed by cooling and solidification of magma. Granite and rhyolite are typical igneous bedrocks in Missouri

Limestone – a type of sedimentary rock composed largely of calcium carbonate

Loess – material transported and deposited by wind and consisting predominantly of silt-size particles

Loamy – soil material containing a relatively equal mixture of sand and silt and a somewhat smaller proportion of clay

Marsh – a type of wetland that is dominated by herbaceous rather than woody plant species

Moist – a site that is moderately well to well drained and has high available water capacity, resulting in a well-balanced supply of moisture (neither too dry nor too wet).

Mudstone – blocky or massive, fine-grained sedimentary rock in which the proportions of clay and silt are approximately equal

Natric – a soil horizon that displays a blocky, columnar, or prismatic structure and has a subhorizon with an exchangeable-sodium saturation of over 15%

Outwash – stratified sediments of sand and gravel removed or “washed out” from a glacier by melt-water streams

Prairie – a vegetative community dominated by perennial grasses and forbs with scattered shrubs and very few trees

Protected – steep, north- and east-facing slopes, which are cooler and moister than other slope aspects

Residuum - unconsolidated, weathered, or partly weathered mineral material that accumulates by disintegration of bedrock in place

Riser – a component of terraces and flood-plain steps consisting of the steep side slope; the escarpment

Riverfront – a vegetative community in the floodplain immediately adjacent and generally parallel to a river or stream channel

River hills – a geographic area characterized by thick, dissected loess deposits, formed immediately adjacent to the edges of the Missouri and Mississippi River floodplains

Sandy – a coarse-sized soil containing a large mixture of sand and gravels and a somewhat smaller proportion of silts and clays with excessive drainage

Sandstone – a sedimentary rock containing dominantly sand-size particles

Savanna – grasslands interspersed with open-grown scattered trees, groupings of trees, and shrubs

Shale – a sedimentary rock formed from clay, silty clay, or silty clay loam deposits and having the tendency to split into thin layers

Shallow – a site with bedrock within 20 inches of the surface

Shoulder – the slope profile position that forms the convex surface near the top of a hill slope; it comprises the transition zone from summit to backslope

Sinkhole – a closed, circular or elliptical depression, commonly funnel-shaped, characterized by subsurface drainage and formed either by dissolution of the surface of underlying bedrock or by collapse of underlying caves within bedrock

Summit – the top or highest area of a hillslope

Swale –shallow, closed depressions irregularly spaced across a floodplain or terrace with an irregularly undulating surface.

Swamp – an area of low, saturated ground, intermittently or permanently covered with water, and predominantly vegetated by shrubs and trees.

Talus – rock fragments of any size or shape (usually coarse and angular) derived from and lying at the base of a cliff or very steep rock slope.

Terrace – a step-like surface, bordering a valley floor that represents the former position of a flood plain

Till – dominantly unsorted and unstratified soil material deposited directly by a glacier

Upland – a general term for the higher ground of a region, in contrast with a low-lying, adjacent land such as a valley or floodplain

Wet – a somewhat poorly, poorly or very poorly drained site that has an oversupply of moisture during the growing season

Woodland – a highly variable vegetative community with a canopy of trees ranging from 30 to 100 percent closure with a sparse midstory and a dense ground flora of grasses, sedges and forbs

References

Love, Charles L. 1997. Soil Survey of Pike County, Missouri. U.S. Dept. of Agric. Natural Resources Conservation Service.

Missouri Department of Conservation. 2006. Missouri Forest and Woodland Community Profiles. Missouri Department of Conservation, Jefferson City, Missouri.

Natural Resources Conservation Service. 2002. Woodland Suitability Groups. Missouri FOTG, Section II, Soil Interpretations and Reports. 30 pgs.

NatureServe. 2010. Vegetation Associations of Missouri (revised). NatureServe, St. Paul, Minnesota.

Nelson, Paul W. 2010. The Terrestrial Natural Communities of Missouri. Missouri Department of Conservation, Jefferson City, Missouri.

Nigh, Timothy A., & Walter A. Schroeder. 2002. Atlas of Missouri Ecoregions. Missouri Department of Conservation, Jefferson City, Missouri.