

Practice: 340 - Cover Crop

Scenario # 1 Legumes

Missouri

Scenario Description:

A legume will be planted as a cover crop immediately after harvest of a row crop, and will be followed by a row crop that will utilize fixed nitrogen and cover crop biomass as a mulch. This scenario assumes that seed will be planted with a no-till drill. Legume seeds must be inoculated with the proper inoculant prior to planting. The cover crop should be allowed to reach early to mid-bloom before it is terminated, using approved chemical and/or mechanical methods, in order to maximize nitrogen fixation. The legume will promote biological nitrogen fixation and reduce energy use by reducing the need for commercial nitrogen fertilizer in following crops.

Before Practice Situation:

Row crops such as corn, soybeans, or cotton are grown and harvested in mid-late fall. Fields are disked immediately following harvest. Residue amounts after harvest average 30% or less, resulting in bare soil being exposed to intense rainfall during the fall, winter, and early spring. Over the winter residue degrades and sediment/nutrient runoff from fields increases. Sheet and rill erosion occurs with visible rills by spring. Runoff from the fields flows into streams, water courses or other water bodies causing degradation to the receiving waters. Soil health (soil organic matter) declines over time as a result of tillage practices and long periods of bare soil. Only commercial nitrogen fertilizer is used for following crops.

After Practice Situation:

Within 30 days after harvest fields are planted with a legume cover crop, typically a clover or vetch species. The cover crop is seeded with a no-till drill. No fertilizer is applied with the cover crop. The cover crop provides soil cover by late fall, throughout the winter, and into the early spring. Runoff and erosion are reduced and no rills are visible on the soil surface in the spring. The cover crop is terminated chemically and/or mechanically prior to spring planting as late as feasible to maximize plant biomass production. Over time, soil health is improved due to the additional biomass, ground cover, and plant diversity introduced to the cropping system. Cover crop residues left on the surface may maximize weed control by increasing allelopathic and mulching effect. By utilizing the nitrogen that is fixed by the legume cover crop, the amount of energy is reduced by reducing the amount of commercial fertilizer that will be needed for the following crop.

Scenario Feature Measure:

Area Planted

| | | | | |
|-------------------------------|-----|------|---------------|---------|
| Scenario Typical Size: | 100 | Acre | Tot Unit Cost | \$51.40 |
|-------------------------------|-----|------|---------------|---------|

| Cost Category | Component Name | Quantity | Unit | Unit Cost | Cost |
|--------------------|--|----------|-------|-----------|-------------------|
| Materials | Red Clover (<i>Trifolium pratense</i>) | 800 | Pound | \$2.60 | \$2,080.00 |
| Materials | Herbicide, Glyphosate | 100 | Acre | \$11.04 | \$1,104.00 |
| Equip./Install. | Chemical, ground application | 100 | Acre | \$4.57 | \$457.00 |
| Equip./Install. | Seeding Operation, No Till/Grass Drill | 100 | Acre | \$14.99 | \$1,499.00 |
| Total Cost: | | | | | \$5,140.00 |

Payment types:

| PayType | Unit Payment | PayType | Unit Payment |
|------------|--------------|--------------|--------------|
| EQIP | \$38.55 | EQIP-HU | \$46.26 |
| EQIP-MRBI | \$38.55 | EQIP-HUMRBI | \$46.26 |
| EQIP-CCPI | \$38.55 | EQIP-HUCCPI | \$46.26 |
| EQIP-NOFEI | \$38.55 | EQIP-HUNOFEI | \$46.26 |

Practice: 340 - Cover Crop

Scenario # 2 Grass or Cereal Grains

Missouri

Scenario Description:

Typically a grass or small grain monoculture will be planted as a cover crop immediately after harvest of a row crop, and will be followed by a row crop that will utilize the residue as a mulch. This scenario assumes that seed will be planted with a no-till drill. The cover crop should be allowed to generate as much biomass as possible, without delaying planting of the following crop. The cover crop will be terminated using approved chemical and/or mechanical methods.

Before Practice Situation:

Row crops such as corn, soybeans, or cotton are grown and harvested in mid-late fall. Fields are disked immediately following harvest. Residue amounts after harvest average 30% or less, resulting in bare soil being exposed to wind erosion and/or intense rainfall during the fall, winter, and early spring. Over the winter residue degrades and sediment/nutrient runoff from fields increases. Sheet and rill erosion occurs with visible rills by spring. Runoff from the fields flows into streams, water courses or other water bodies causing degradation to the receiving waters. Soil health (soil organic matter) declines over time as a result of tillage practices, low residue crops, and long periods of bare soil.

After Practice Situation:

Within 30 days after harvest of row crop, fields are planted with a grass or small grain cover crop, typically ryegrass and clover. The cover crop is seeded with a no-till drill. No additional fertilizer is applied with the cover crop. The cover crop provides soil cover by late fall, throughout the winter, and into the early spring. Runoff and erosion are reduced and no rills are visible on the soil surface in the spring. Wind erosion is reduced by standing residues. The cover crop is terminated with approved chemical and/or mechanical methods prior to spring planting as late as feasible to maximize plant biomass production. Over time, soil health is improved due to the additional biomass, ground cover, soil infiltration, and plant diversity introduced to the cropping system. Cover crop residues left on the surface may maximize weed control by increasing allelopathic and mulching effect.

Scenario Feature Measure:

Area Planted

| | | | | |
|-------------------------------|-----|------|---------------|---------|
| Scenario Typical Size: | 100 | Acre | Tot Unit Cost | \$49.35 |
|-------------------------------|-----|------|---------------|---------|

| Cost Category | Component Name | Quantity | Unit | Unit Cost | Cost |
|-----------------|--|----------|-------|-----------|------------|
| Materials | Ryegrass, Annual (Lolium multiflorum) | 1500 | Pound | \$1.25 | \$1,875.00 |
| Materials | Herbicide, Glyphosate | 100 | Acre | \$11.04 | \$1,104.00 |
| Equip./Install. | Chemical, ground application | 100 | Acre | \$4.57 | \$457.00 |
| Equip./Install. | Seeding Operation, No Till/Grass Drill | 100 | Acre | \$14.99 | \$1,499.00 |

Total Cost: \$4,935.00

Payment types:

| PayType | Unit Payment | PayType | Unit Payment |
|-----------|--------------|-------------|--------------|
| EQIP | \$37.01 | EQIP-HU | \$44.42 |
| EQIP-MRBI | \$37.01 | EQIP-HUMRBI | \$44.42 |
| EQIP-CCPI | \$37.01 | EQIP-HUCCPI | \$44.42 |

Practice: 340 - Cover Crop

Scenario # 3 Winter Kill Species

Missouri

Scenario Description:

Typically a small grain or small grain-brassica mix (may also use forage sorghum, legumes, buckwheat, etc) will be planted as a cover crop immediately after harvest of a row crop, and will be followed by a row crop that will utilize the residue as a mulch. This scenario assumes that seed will be planted with a no-till drill. The cover crop species established under this scenario will winter kill, meaning no species termination is required.

Before Practice Situation:

Row crops such as corn, soybeans, or cotton are grown and harvested in mid-late fall. Fields are disked immediately following harvest. Residue amounts after harvest average 30% or less, resulting in bare soil being exposed to wind erosion and/or intense rainfall during the fall, winter, and early spring. Over the winter residue degrades and sediment/nutrient runoff from fields increases. Sheet and rill erosion occurs with visible rills by spring. Runoff from the fields flows into streams, water courses or other water bodies causing degradation to the receiving waters. Soil health (soil organic matter) declines over time as a result of tillage practices, low residue crops, and long periods of bare soil.

After Practice Situation:

Within 30 days after harvest of row crop, fields are planted with a small grain-brassica mix cover crop, typically oats and oilseed radish. The cover crop is seeded with a no-till drill. No additional fertilizer is applied with the cover crop. The cover crop provides soil cover by late fall, throughout the winter, and into the early spring. Runoff and erosion are reduced and no rills are visible on the soil surface in the spring. The cover crop is established using winter kill species which should not require termination in the spring. Over time, soil health is improved due to the additional biomass, ground cover, and plant diversity introduced to the cropping system. Wind erosion is reduced by standing residues. Cover crop residues left on the surface may maximize weed control by increasing allelopathic and mulching effect.

Scenario Feature Measure:

Area planted

| | | | | |
|-------------------------------|-----|------|---------------|---------|
| Scenario Typical Size: | 100 | Acre | Tot Unit Cost | \$33.61 |
|-------------------------------|-----|------|---------------|---------|

| Cost Category | Component Name | Quantity | Unit | Unit Cost | Cost |
|--------------------|--|----------|-------|-----------|-------------------|
| Materials | Oats (Avena sativa) | 2500 | Pound | \$0.46 | \$1,150.00 |
| Materials | Oilseed Radish (Raphanus sativus var. | 200 | Pound | \$3.56 | \$712.00 |
| Equip./Install. | Seeding Operation, No Till/Grass Drill | 100 | Acre | \$14.99 | \$1,499.00 |
| Total Cost: | | | | | \$3,361.00 |

Payment types:

| PayType | Unit Payment | PayType | Unit Payment |
|-----------|--------------|-------------|--------------|
| EQIP | \$25.21 | EQIP-HU | \$30.25 |
| EQIP-MRBI | \$25.21 | EQIP-HUMRBI | \$30.25 |
| EQIP-CCPI | \$25.21 | EQIP-HUCCPI | \$30.25 |

Practice: 340 - Cover Crop

Scenario # 4 Species Mix

Missouri

Scenario Description:

This scenario reflects the establishment of a diverse mix of cover crops consisting of two or more species which can include a combination of grasses, legumes, forbs, or other herbaceous plants. Cover crop will be planted immediately after harvest of a row crop, and will be followed by a row crop that will utilize the residue as a mulch. This scenario assumes that seed will be planted with a no-till drill. The cover crop should be allowed to generate as much biomass as possible, without delaying planting of the following crop. The cover crop will be terminated using approved chemical and/or mechanical methods.

Before Practice Situation:

Row crops such as corn, soybeans, or cotton are grown and harvested in mid-late fall. Fields are disked immediately following harvest. Residue amounts after harvest average 30% or less, resulting in bare soil being exposed to wind erosion and/or intense rainfall during the fall, winter, and early spring. Over the winter residue degrades and sediment/nutrient runoff from fields increases. Sheet and rill erosion occurs with visible rills by spring. Runoff from the fields flows into streams, water courses or other water bodies causing degradation to the receiving waters. Soil health (soil organic matter) declines over time as a result of tillage practices, low residue crops, and long periods of bare soil.

After Practice Situation:

Within 30 days after harvest of row crop, fields are planted with a diverse mix of cover crop species. The cover crop is seeded with a no-till drill. No additional fertilizer is applied with the cover crop. The cover crop provides soil cover by late fall, throughout the winter, and into the early spring. Runoff and erosion are reduced and no rills are visible on the soil surface in the spring. Wind erosion is reduced by standing residues. The cover crop is terminated with approved chemical and/or mechanical methods prior to spring planting as late as feasible to maximize plant biomass production. Over time, soil health is improved at an accelerated pace due to the diversity in additional biomass and ground cover which provides increased soil infiltration, and plant diversity introduced to the cropping system. Cover crop residues left on the surface may maximize weed control by increasing allelopathic and mulching effect.

Scenario Feature Measure:

Area planted

| | | | | |
|-------------------------------|-----|------|---------------|---------|
| Scenario Typical Size: | 100 | Acre | Tot Unit Cost | \$50.00 |
|-------------------------------|-----|------|---------------|---------|

| Cost Category | Component Name | Quantity | Unit | Unit Cost | Cost |
|-----------------|--|----------|-------|-----------|------------|
| Materials | Red Clover (<i>Trifolium pratense</i>) | 400 | Pound | \$2.60 | \$1,040.00 |
| Materials | Rye, Cereal (<i>Secale cereale</i> L.) | 3000 | Pound | \$0.30 | \$900.00 |
| Materials | Herbicide, Glyphosate | 100 | Acre | \$11.04 | \$1,104.00 |
| Equip./Install. | Chemical, ground application | 100 | Acre | \$4.57 | \$457.00 |
| Equip./Install. | Seeding Operation, No Till/Grass Drill | 100 | Acre | \$14.99 | \$1,499.00 |

Total Cost: \$5,000.00

Payment types:

| PayType | Unit Payment | PayType | Unit Payment |
|------------|--------------|--------------|--------------|
| EQIP | \$37.50 | EQIP-HU | \$45.00 |
| EQIP-MRBI | \$37.50 | EQIP-HUMRBI | \$45.00 |
| EQIP-CCPI | \$37.50 | EQIP-HUCCPI | \$45.00 |
| EQIP-NOFEI | \$37.50 | EQIP-HUNOFEI | \$45.00 |

Practice: 340 - Cover Crop

Scenario # 5 Organic Cover Crop

Scenario Description:

Missouri

Typically a small grain or small grain-legume mix (may also use forage sorghum, radishes, turnips, buckwheat, etc) will be planted as a cover crop immediately after harvest of an organically grown crop, and will be followed by an organically grown crop that will utilize the residue as a mulch. This scenario assumes that seed will be planted with a no-till drill. The cover crop should be allowed to generate as much biomass as possible, without delaying planting of the following crop. The cover crop will be terminated using a mechanical kill method (mowing, rolling, undercutting, etc.), a minimum of 3 weeks prior to planting the subsequent crop. This scenario **REQUIRES** use of Certified Organic Seed.

Before Practice Situation:

Organically grown crops such as various vegetable and fruit crops (along with organically produced row crops) are grown and harvested in mid-late fall. Fields are disked immediately following harvest. Residue amounts after harvest average 30% or less, resulting in bare soil being exposed to wind erosion and/or intense rainfall during the fall, winter, and early spring. Over the winter residue degrades and sediment/nutrient runoff from fields increases. Sheet and rill erosion occurs with visible rills by spring. Runoff from the fields flows into streams, water courses or other water bodies causing degradation to the receiving waters. Soil health (soil organic matter) declines over time as a result of tillage practices, low residue crops, and long periods of bare soil.

After Practice Situation:

Within 30 days after harvest of organic crop, fields are planted with a small grain-legume mix cover crop, typically rye and clover. The cover crop is seeded with a no-till drill. No additional fertilizer is applied with the cover crop. The cover crop provides soil cover by late fall, throughout the winter, and into the early spring. Runoff and erosion are reduced and no rills are visible on the soil surface in the spring. The cover crop is terminated with using a mechanical kill method (mowing, rolling, undercutting, etc.), prior to spring planting as late as feasible to maximize plant biomass production. Over time, soil health is improved due to the additional biomass, ground cover, and plant diversity introduced to the cropping system. Wind erosion is reduced by standing residues. Cover crop residues left on the surface may maximize weed control by increasing allelopathic and mulching effect.

Scenario Feature Measure:

Area planted

| | | | | |
|-------------------------------|----|------|---------------|----------|
| Scenario Typical Size: | 25 | Acre | Tot Unit Cost | \$100.02 |
|-------------------------------|----|------|---------------|----------|

| Cost Category | Component Name | Quantity | Unit | Unit Cost | Cost |
|--------------------|--|----------|-------|-----------|-------------------|
| Materials | Certified Organic, Crimson Clover (Trifolium | 200 | Pound | \$5.35 | \$1,070.00 |
| Materials | Certified Organic, Oats (Avena sativa) | 450 | Pound | \$0.62 | \$279.00 |
| Equip./Install. | Tillage, Light | 25 | Acre | \$10.29 | \$257.25 |
| Equip./Install. | Mechanical weed control, Vegetation | 25 | Acre | \$20.78 | \$519.50 |
| Equip./Install. | Seeding Operation, No Till/Grass Drill | 25 | Acre | \$14.99 | \$374.75 |
| Total Cost: | | | | | \$2,500.50 |

Payment types:

| PayType | Unit Payment | PayType | Unit Payment |
|----------|--------------|------------|--------------|
| EQIP-NOI | \$75.02 | EQIP-HUNOI | \$90.02 |