

## *Ecological Site Description*

### **Loamy Footslope Savanna**

**R107BY008MO**


- (*Quercus macrocarpa*/*Salix humilis*/*Andropogon gerardii* - *Tripsacum dactyloides*)
- (bur oak/prairie willow/big bluestem – Eastern gama grass)

An Ecological Site Description (ESD) is a reference document of ecological knowledge regarding a particular land area (ecological site). An ESD describes ecological potential and ecosystem dynamics of land areas and their potential management. Ecological sites are linked to soil survey map unit components, which allows for mapping of ecological sites. (**NOTE:** *This is a “provisional” ESD, and is subject to change. It contains basic ecological information sufficient for conservation planning and land management in Missouri. After additional information is developed and reviewed, a “Correlated” ESD will be published and will be available via the Web Soil Survey <http://websoilsurvey.nrcs.usda.gov> .)*

**Major Land Resource Area:** 107B – Iowa and Missouri Deep Loess Hills

### **Introduction**

The Iowa and Missouri Deep Loess Hills (area outlined in red on the map) encompass the Missouri River floodplain and associated loess-covered uplands, from about Sioux City Iowa in the north to




central Missouri. Elevation is about 1,565 feet on the highest ridges, to about 600 feet along the Missouri River near Glasgow in central Missouri. Local relief varies from 10 to 20 feet in the major river floodplains, to 50 to 100 feet in the dissected uplands, with loess bluffs of 200 to 300 feet along the Missouri River. The loess thins with distance from the Missouri river, and local relief decreases. The loess caps pre-Illinoian till, which crops out on lower hillslopes near the edges of the MLRA. The underlying bedrock is mainly Pennsylvanian and Cretaceous-aged shale, mudstone and sandstone.

Loamy Footslope Savannas are within the green areas on the map (Missouri portion only; Iowa distributions are currently under review). These sites are primarily in the central part of the MLRA. Soils are very deep, and loamy throughout.

### **Physiographic Features**

This site is on footslopes, with slopes of 1 to 7 percent. The site receives runoff from adjacent upland sites. This site does not flood.

The following figure (adapted from Young & Kowalewycz, 1994) shows the typical landscape position of this ecological site, and landscape relationships with other ecological sites. The site is within the area labeled “1”, on footslopes. Floodplain sites, such as the Loamy Floodplain Forest shown in the diagram, are typically downslope. A variety of upland sites are upslope, especially Deep Loess sites.



**Soil Features**

These soils have no rooting restriction. The soils were formed under prairie vegetation, and have dark, organic-rich surface horizons. Parent material is colluvium derived from loess. Surface horizons are primarily loam. Subsurface horizons are loamy. Soil series associated with this site include Judson and Napier.

**Ecological Dynamics**

*Information contained in this section was developed using historical data, professional experience, field reviews, and scientific studies. The information presented is representative of very complex vegetation communities. Key indicator plants, animals and ecological processes are described to help inform land management decisions. Plant communities will differ across the MLRA because of the naturally occurring variability in weather, soils, and aspect. The Reference Plant Community is not necessarily the management goal. The species lists are representative and are not botanical descriptions of all species occurring, or potentially occurring, on this site. They are not intended to cover every situation or the full range of conditions, species, and responses for the site.*

The reference plant community is characterized as a savanna unit with a ground cover dominated by big and little bluestem, Indian grass, switch grass, eastern gamagrass and a wide variety of prairie wildflowers while other species such as Culver’s root, Michigan lily, and bunchflower added to the mix of footslope prairie species. Slightly higher areas in the landscape support scattered bur oak, wild plum and prairie willow.

While not as typically dry and fire prone as the Loess Upland Prairies, this ecological site still likely burned every 1 to 3 years. Fire removed dead plant litter and provided room for a lush growth of prairie vegetation. Fire also kept woody species at bay. Grazing by native large herbivores, such as bison, elk, and deer, also impacted these sites. Their activities altered the composition, fuel loads and structure of the vegetation, creating a diversity of structure and composition. The partially wooded draws would have burned less intensely and less frequently. During fire free intervals woody species would have increased in abundance and spread out onto the prairie.

These sites are very productive. Today, Loamy Footslope Savannas are nearly extirpated from the region as the former footslope prairies and savannas have been converted to intensive agriculture. Few quality remnants exist. While re-establishing prairie and savanna on converted agriculture sites is beneficial to wildlife, restoration to the reference state from agricultural land is a long term proposition with uncertain outcomes.

**Reference State Plant Community**

## Canopy Trees

Common Name	Botanical Name	Cover % (low-high)	Canopy Height (ft)
BUR OAK	<i>Quercus macrocarpa</i>	5-10	60

## Shrubs

Common Name	Botanical Name	Cover % (low-high)	Canopy Height (ft)
AMERICAN HAZELNUT	<i>Corylus americana</i>	5-10	5
NEW JERSEY TEA	<i>Ceanothus americanus</i>	5-10	4
LEAD PLANT	<i>Amorpha canescens</i>	5-10	3
PRAIRIE WILLOW	<i>Salix humilis</i>	5-20	5
WILD PLUM	<i>Prunus americana</i>	0-10	8

## Forbs

Common Name	Botanical Name	Cover % (low-high)
WILD QUININE	<i>Parthenium integrifolium</i>	5-20
BUTTERFLY WEED	<i>Asclepias tuberosa</i>	5-20
BLAZING STAR	<i>Liatris pycnostachya</i>	5-20
MISSOURI GOLDENROD	<i>Solidago missouriensis</i>	5-20
ASHY SUNFLOWER	<i>Helianthus mollis</i>	5-20
RATTLESNAKE MASTER	<i>Eryngium yuccifolium</i>	5-20
WHITE INDIGO	<i>Baptisia alba</i>	5-20
PRAIRIE MILKWEED	<i>Asclepias sullivantii</i>	5-20
FIELD MILKWORT	<i>Polygala sanguinea</i>	5-10
ROSWINE	<i>Silphium integrifolium</i>	5-20
PURPLE PRAIRIE CLOVER	<i>Dalea purpurea</i>	5-20
PURPLE CONEFLOWER	<i>Echinacea purpurea</i>	5-20
COMPASS PLANT	<i>Silphium laciniatum</i>	5-20
HOARY PUCCOON	<i>Lithospermum canescens</i>	5-20
WILD BERGAMOT	<i>Monarda fistulosa</i>	5-20
CULVER'S ROOT	<i>Veronicastrum virginicum</i>	5-20
BUNCHFLOWER	<i>Melanthium virginicum</i>	5-20
MICHIGAN LILY	<i>Lilium michiganensis</i>	5-10

## Grasses and sedges

Common Name	Botanical Name	Cover % (low-high)
LITTLE BLUESTEM	<i>Schizachyrium scoparium</i>	10-20
EASTEN GAMA GRASS	<i>Tripsacum dactyloides</i>	10-30
PRAIRIE DROPSEED	<i>Sporobolus heretoeipis</i>	5-10
BIG BLUESTEM	<i>Andropogon gerardii</i>	30-50
INDIAN GRASS	<i>Sorghastrum nutans</i>	10-20
SWITCHGRASS	<i>Panicum virgatum</i>	10-20

**Site Interpretations***Wildlife\***Prairie Phase:*

- Game species that utilize this ecological site include:  
Northern Bobwhite will utilize this ecological site for food (seeds, insects) and cover needs (escape, nesting and roosting cover).

Cottontail rabbits will utilize this ecological site for food (seeds, soft mast) and cover needs.

Turkey will utilize this ecological site for food (seeds, green browse, soft mast, insects) and nesting and brood-rearing cover. Turkey poult feed heavily on insects provided by this site type.

White-tailed Deer will utilize this ecological site for browse (plant leaves in the growing season, seeds and soft mast in the fall/winter). This site type also can provide escape cover.

- Bird species associated with this ecological site's reference state condition: Breeding birds as related to vegetation structure (related to time since fire, grazing, haying, and mowing):

Vegetation Height Short (< 0.5 meter, low litter levels, bare ground visible):

Grasshopper Sparrow, Horned Lark, Upland Sandpiper, Greater Prairie Chicken, Northern Bobwhite

Mid-Vegetation Height (0.5 – 1 meter, moderate litter levels, some bare ground visible):

Eastern Meadowlark, Dickcissel, Field Sparrow, Upland Sandpiper, Greater Prairie Chicken, Northern Bobwhite, Eastern Kingbird, Bobolink, Lark Sparrow

Tall Vegetation Height (> 1 meter, moderate-high litter levels, little bare ground visible):

Henslow's Sparrow, Dickcissel, Greater Prairie Chicken, Field Sparrow, Northern Bobwhite, Sedge Wren, Northern Harrier

Brushy – Mix of grasses, forbs, native shrubs (e.g., *Rhus copallina*, *Prunus americana*, *Rubus* spp., *Rosa carolina*) and small trees (e.g., *Cornus racemosa*): Bell's Vireo, Yellow-breasted Chat, Loggerhead Shrike, Brown Thrasher, Common Yellowthroat

Winter Resident: Short-Eared Owl, Le Conte's Sparrow

- Amphibian and reptile species associated with this ecological site's reference state condition: prairies with or nearby to fishless ponds/pools (may be ephemeral) may have Eastern Tiger Salamander (*Ambystoma tigrinum tigrinum*) and Western Chorus Frog (*Pseudacris triseriata triseriata*); prairies with crawfish burrows may have Northern Crawfish Frog (*Rana areolata circulosa*); other species include Northern Prairie Skink (*Eumeces septentrionalis septentrionalis*), Ornate Box Turtle (*Terrapene ornata ornata*), Western Slender Glass Lizard (*Ophisaurus attenuatus attenuatus*), Eastern Yellow-bellied Racer (*Coluber constrictor flaviventris*), Prairie Ring-necked Snake (*Diadophis punctatus arnyi*), and Bullsnake (*Pituophis catenifer sayi*).
- Small mammals associated with this ecological site's reference state condition: Least Shrew (*Cryptotis parva*), Franklin's Ground Squirrel (*Spermophilus franklinii*), Plains Pocket Gopher (*Geomys bursarius*), Prairie Vole (*Microtus ochrogaster*), Southern Bog Lemming (*Synaptomys cooperi*), Meadow Jumping Mouse (*Zapus hudsonius*), Thirteen-lined Ground Squirrel (*Spermophilus tridecemlineatus*) and Badger (*Taxidea taxus*).

- Invertebrates:  
Many native insect species are likely associated with this ecological site, especially native bees, ants, beetles, butterflies and moths, and crickets, grasshoppers and katydids. However information on these groups is often lacking enough resolution to assign them to individual ecological sites.

Insect species known to be associated with this ecological site's reference state condition: Regal Fritillary butterfly (*Speyeria idalia*) whose larvae feed primarily on native prairie violets (*Viola pedata*, *V. pedatifida*, and *V. sagittata*); Mottled Dusky Wing butterfly (*Erynnis martialis*), Golden Byssus butterfly (*Problema byssus kumskaka*), Delaware Skipper butterfly (*Atryone logan logan*), and Crossline Skipper butterfly (*Polites origenes*). The larvae of the moth *Eucosma bipunctella* bore into compass plant (*Silphium laciniatum*) roots and feed and the larvae of the moth *Eucosma giganteana* bore into a number of *Silphium* species roots and feed. Native bees, important pollinators, that may be associated with this ecological site's reference condition include: *Colletes brevicornis*, *Andrena beameri*, *A. helianthiformis*, *Protandrena rudbeckiae*, *Halictus parallelus*, *Lasioglossum albipennis*, *L. coreopsis*, *L. disparilis*, *L. nymphaeum*, *Ashmeadiella buconis*, *Megachile addenda*, *Anthidium psoraleae*, *Eucera hamata*, *Melissodes coloradensis*, *M. coreopsis*, and *M. vernoniae*. The Short-winged Katydid (*Amblycorypha parvipennis*), Green Grasshopper (*Hesperotettix speciosus*) and Two-voiced Conehead katydid (*Neoconcephalus bivocatus*) are possible orthopteran associates of this ecological site.

Other invertebrate associates include the Grassland Crayfish (*Procambarus gracilis*).

#### Savanna Phase:

- Oaks and hickories provide an important food source for many animals including White-tailed Deer, Wild Turkey, and Fox Squirrel.
- Both snags and live cavity or den trees provide important food and cover for vertebrate wildlife. Snags are also very important to invertebrate species. Fox Squirrel, Red-headed Woodpecker and Eastern Bluebird utilize snags and den trees for foraging, nesting or shelter. "Wolf" trees are a particularly valuable type of live cavity tree. These large diameter, often open-grown, old-ages, hollow trees provide both cavities for wildlife and usually hard or soft mast food sources. Large diameter snags and den trees are particularly important wildlife habitat features to retain.
- Game species that utilize this ecological site include:

Northern Bobwhite will utilize this ecological site for food (seeds, insects) and cover needs (escape, nesting and roosting cover).

Cottontail rabbits will utilize this ecological site for food (seeds, soft mast) and cover needs.

Turkey will utilize this ecological site for food (seeds, green browse, soft mast, insects) and nesting and brood-rearing cover. Turkey poults feed heavily on insects provided by this site type.

White-tailed Deer will utilize this ecological site for browse (plant leaves in the growing season, seeds and soft mast in the fall/winter). This site type also can provide escape cover.

- Bird species associated with this ecological site's reference state condition:  
Breeding birds: Northern Bobwhite, Eastern Kingbird, Eastern Bluebird, Brown Thrasher, White-eyed Vireo, Prairie Warbler, Field Sparrow, Eastern Towhee, Red-headed Woodpecker, Great Crested Flycatcher, Loggerhead Shrike

Winter resident: American Tree Sparrow, Harris' Sparrow

- Amphibian and reptile species likely associated with this ecological site's reference state condition: Ornate Box Turtle (*Terrapene ornata ornata*), Northern Fence Lizard (*Sceloporus undulatus hyacinthinus*), Five-lined Skink (*Eumeces fasciatus*), Western Slender Glass Lizard (*Ophisaurus attenuatus attenuatus*), Eastern Yellow-bellied Racer (*Coluber constrictor flaviventris*), Prairie Ring-necked Snake (*Diadophis punctatus arnyi*), and Rough Green Snake (*Ophedryx aestivus aestivus*). Sites containing or nearby to fishless or ephemeral ponds/pools may support the Eastern Tiger Salamander (*Ambystoma tigrinum tigrinum*).
- Small mammals likely associated with this ecological site's reference state condition: Fox Squirrel (*Sciurus niger*), Woodland Vole (*Microtus pinetorum*), Least Shrew (*Cryptotis parva*), and Indiana Bat (*Myotis sodalis*). Indiana bats utilize suitable live, dying or dead roost trees for summer habitat and raising young. Suitable roost trees typically have exfoliating or flaking bark and are larger in diameter.
- Invertebrates – Many native insect species are likely associated with this phase of this ecological site's reference state condition, especially native bees, ants, beetles, butterflies and moths, and crickets, grasshoppers and katydids. However we don't have enough information on these groups to assign them to this phase of this ecological site's reference state condition at this time.

\*This section prepared by Mike Leahy, Natural Areas Coordinator, Missouri Department of Conservation, 2013

#### Forestry

- **Management:** **This ecological site is not recommended for traditional timber management activity.** Historically this site was dominated by a ground cover of native prairie grasses and forbs. Some scattered open grown trees may have also been present. May be suitable for non-traditional forestry uses such as windbreaks, environmental plantings, alley cropping (a method of planting, in which rows of trees or shrubs are interspersed with rows of crops) or woody bio-fuels.

**Glossary**

*Backslope* – a hillslope profile position that forms the steepest and generally linear, middle portion of the slope.

*Backswamp* – marshy or swampy, depressed areas of flood plains between natural levees and valley sides or terraces

*Calcareous* – the presence of calcium carbonate in the soil parent material within the rooting zone; relatively alkaline

*Claypan* – a dense, compact, slowly permeable layer in the subsoil having much higher clay content than the overlying material

*Chert* – hard, extremely dense or compact crystalline sedimentary rock, consisting dominantly of interlocking crystals of quartz

*Cliff* – a significant vertical, or near vertical, rock exposure

*Dolomite* – a type of sedimentary rock that is a carbonate mineral composed of calcium magnesium carbonate

*Drainageway* – the upper most reach of a stream channel system characterized by little meandering

*Dry* – a site where soil moisture is limiting during the growing season; low available water capacity

*Dune* – a low mound, ridge, bank or hill of loose, wind-blown sand

*Exposed* – steep, south and west-facing slopes, which are warmer and drier than other slope aspects

*Flatwoods* – a type of woodland that occurs on soils with a root restricting subsoil layer within 20 to 30 inches, resulting in very slow runoff and ponding that remains saturated for most of the winter and early spring months but dries out and becomes very dry in the summer months; plants that grow there must be adapted to both conditions

*Floodplain* – the nearly level plain that borders a stream and is subject to inundation under flood-stage conditions

*Footslope* – a hillslope position at the base of a slope where hillslope sediment (colluvium) accumulates

*Forest* – a vegetative community dominated by trees forming a closed canopy and interspersed with shade-tolerant understory species

*Fragipan* – a dense, brittle subsoil horizon that is extremely hard and compact when dry

*Glade* – open, rocky, barren vegetative community dominated by drought-adapted forbs and grasses, typically with scattered, stunted woody plants

*Igneous* – bedrock formed by cooling and solidification of magma. Granite and rhyolite are typical igneous bedrocks in Missouri

*Limestone* – a type of sedimentary rock composed largely of calcium carbonate

*Loess* – material transported and deposited by wind and consisting predominantly of silt-size particles

*Loamy* – soil material containing a relatively equal mixture of sand and silt and a somewhat smaller proportion of clay

*Marsh* – a type of wetland that is dominated by herbaceous rather than woody plant species

*Moist* – a site that is moderately well to well drained and has high available water capacity, resulting in a well-balanced supply of moisture (neither too dry nor too wet).

*Mudstone* – blocky or massive, fine-grained sedimentary rock in which the proportions of clay and silt are approximately equal

*Natric* – a soil horizon that displays a blocky, columnar, or prismatic structure and has a subhorizon with an exchangeable-sodium saturation of over 15%

*Outwash* – stratified sediments of sand and gravel removed or “washed out” from a glacier by melt-water streams

*Prairie* – a vegetative community dominated by perennial grasses and forbs with scattered shrubs and very few trees

*Protected* – steep, north- and east-facing slopes, which are cooler and moister than other slope aspects

*Residuum* - unconsolidated, weathered, or partly weathered mineral material that accumulates by disintegration of bedrock in place

*Riser* – a component of terraces and flood-plain steps consisting of the steep side slope; the escarpment

*Riverfront* – a vegetative community in the floodplain immediately adjacent and generally parallel to a river or stream channel

*River hills* – a geographic area characterized by thick, dissected loess deposits, formed immediately adjacent to the edges of the Missouri and Mississippi River floodplains

*Sandy* – a coarse-sized soil containing a large mixture of sand and gravels and a somewhat smaller proportion of silts and clays with excessive drainage

*Sandstone* – a sedimentary rock containing dominantly sand-size particles

*Savanna* – grasslands interspersed with open-grown scattered trees, groupings of trees, and shrubs

*Shale* – a sedimentary rock formed from clay, silty clay, or silty clay loam deposits and having the tendency to split into thin layers

*Shallow* – a site with bedrock within 20 inches of the surface

*Shoulder* – the slope profile position that forms the convex surface near the top of a hill slope; it comprises the transition zone from summit to backslope

*Sinkhole* – a closed, circular or elliptical depression, commonly funnel-shaped, characterized by subsurface drainage and formed either by dissolution of the surface of underlying bedrock or by collapse of underlying caves within bedrock

*Summit* – the top or highest area of a hillslope

*Swale* – shallow, closed depressions irregularly spaced across a floodplain or terrace with an irregularly undulating surface.

*Swamp* – an area of low, saturated ground, intermittently or permanently covered with water, and predominantly vegetated by shrubs and trees.

*Talus* – rock fragments of any size or shape (usually coarse and angular) derived from and lying at the base of a cliff or very steep rock slope.

*Terrace* – a step-like surface, bordering a valley floor that represents the former position of a flood plain

*Till* – dominantly unsorted and unstratified soil material deposited directly by a glacier

*Upland* – a general term for the higher ground of a region, in contrast with a low-lying, adjacent land such as a valley or floodplain

*Wet* – a somewhat poorly, poorly or very poorly drained site that has an oversupply of moisture during the growing season

*Woodland* – a highly variable vegetative community with a canopy of trees ranging from 30 to 100 percent closure with a sparse midstory and a dense ground flora of grasses, sedges and forbs

## References

Fitzgerald, J.A. and D.N. Pashley. 2000a. Partners in Flight bird conservation plan for the Ozark/Ouachitas. American Bird Conservancy.

Fitzgerald, J.A. and D.N. Pashley. 2000b. Partners in Flight bird conservation plan for the Dissected Till Plains. American Bird Conservancy.

Heitzman, J.R. and J.E. Heitzman. 1996. Butterflies and moths of Missouri. 2<sup>nd</sup> ed. Missouri Department of Conservation, Jefferson City.

Jacobs, B. 2001. Birds in Missouri. Missouri Department of Conservation, Jefferson City.

Johnson, T.R. 2000. The amphibians and reptiles of Missouri. 2<sup>nd</sup> ed. Missouri Department of Conservation, Jefferson City.

Nelson, Paul W. 2010. The Terrestrial Natural Communities of Missouri. Missouri Department of Conservation, Jefferson City, Missouri.

Nigh, Timothy A., & Walter A. Schroeder. 2002. Atlas of Missouri Ecoregions. Missouri Department of Conservation, Jefferson City, Missouri.

Pitts, D.E. and W.D. McGuire. 2000. Wildlife management for Missouri landowners. 3<sup>rd</sup> ed. Missouri Department of Conservation, Jefferson City.

Schwartz, C.W., E.R. Schwartz and J.J. Conley. 2001. The wild mammals of Missouri. University of Missouri Press, Columbia and Missouri Department of Conservation, Jefferson City.

Young, Fred J., & Patricia Kowalewycz. 1994. Soil Survey of Atchison County, Missouri. U.S Dept. of Agric. Soil Conservation Service.