

Author(s)/participant(s): W. Lujan, N. Svendsen, J. Alexander, K. Walstad, J. Siddoway, M. Hansen

Contact: Missoula Area Office

Date: 5/2010

Reference site used? **Yes**

MLRA/LRU: 44A

Ecological Site: Sandy LRU B

This *must* be verified based on soils and

climate (see Ecological Site Description). Current plant community *cannot* be used to identify the ecological site.

<p>Indicators. For each indicator, describe the potential for the site. Where possible, (1) use numbers, (2) include expected range of values for above- and below-average years for each community within the reference state (when appropriate), and (3) cite data.</p>
<p>1. Number and extent of rills: Rills are unlikely to occur in the Taller Bunchgrass State.</p>
<p>2. Presence of water flow patterns: Water flow patterns are generally not evident in the reference state. Following occasional (5-30% probability), heavy thunderstorms and winter thaw events, short, sinuous, discontinuous flow patterns may be apparent, but rare, on slopes ranging from 4-15%. Water flow patterns should not be evident on slopes lower than 4%.</p>
<p>3. Number and height of erosional pedestals or terracettes: None to very slight. Rarely pedestals up to 0.5 inches may be encountered.</p>
<p>4. Bare ground from Ecological Site Description or other studies (rock, litter, lichen, moss, and plant canopy are <i>not</i> bare ground): Bare ground should not exceed 5% – bare areas tend to be inconspicuous and not connected.</p>
<p>5. Number of gullies and erosion associated with gullies: Gullies should not occur in the Taller Bunchgrass State. If there is evidence of past erosion that has created gullies, these areas should be stabilized and have no active erosion.</p>
<p>6. Extent of wind scoured, blowouts and/or depositional areas: Appearance or evidence of these erosional features on the landscape would not be present on this site.</p>
<p>7. Amount of litter movement (describe size and distance expected to travel): Litter will be evident across this site representing organic debris from the vegetation of the functional/structural groups and will not move. A severe convection storm or a significant thaw event could cause litter to move short distances, especially on slopes greater than 6%.</p>
<p>8. Soil surface (top few millimeters) resistance to erosion (stability values are averages – most sites will show a range of values for both plant canopy and interspaces, if different): Resistance to erosion will be high with soil stability values of 5 or 6; areas of bare soil on this site may have values between 3 and 5 if not under plant canopy.</p>
<p>9. Soil surface structure and Soil Organic Matter (SOM) content (include type and strength of structure, and A-horizon color and thickness for both plant canopy and interspaces, if different): Structure is granular at the soil surface. Organic matter is about 1.5%. The surface horizon is 4 to 8 inches thick.</p>
<p>10. Effect of plant community composition (relative proportion of different functional groups) and spatial distribution on infiltration and runoff: The reference plant community (1.1) is dominated by rough fescue which will maximize infiltration and minimize runoff throughout the site. With the increase of Idaho fescue in Plant community (1.2) infiltration may slightly decrease and runoff may slightly increase but overall this plant community will have only minor affects on infiltration and runoff.</p>
<p>11. Presence and thickness of compaction layer (usually none; describe soil profile features which may be mistaken for compaction on this site): A compaction layer would not be expected on this ecological site.</p>
<p>12. Functional/Structural Groups (list in order of descending dominance by above-ground weight using symbols: >>, >, = to indicate much greater than, greater than, and equal to): Plant community 1.1 – Taller cool season bunchgrasses (rough fescue) >> mid-stature cool season bunchgrasses (Idaho fescue) > cool season rhizomatous grasses (western wheatgrass), shortgrasses (prairie junegrass) and grasslikes (sedges) = perennial forbs > shrubs. Plant community 1.2 – rough fescue and Idaho fescue share dominance – the other functional/structural groups will remain the same in descending order.</p>
<p>13. Amount of plant mortality and decadence (include which functional groups are expected to show mortality or decadence): Plant mortality for all functional groups will be low, but there will be some natural mortality of functional groups over time. Prolonged droughts and/or excessive rest may show increases in mortality and decadence for all plant groups.</p>
<p>14. Average percent litter cover (65-80%) and depth (0 to 0.5 inches). Note: The majority of the litter in the plant community in the Taller Bunchgrass State will be non-persistent.</p>
<p>15. Expected annual production (this is TOTAL annual above-ground production, not just forage production): 1,100 #/acre – 2,200 #/acre for the reference community (1.1) with a Representative Value (RV) of 1,650 #/acre. Production varies based on effective precipitation and natural variability of soil properties for this ecological site.</p>
<p>16. Potential invasive (including noxious) species (native and non-native). List species which characterize degraded states and which, after a threshold is crossed, “will continue to increase regardless of the management of the site” and may eventually dominate the site: Cheatgrass, knapweed spp., leafy spurge, sulphur cinquefoil, dalmatian toadflax, houndstongue, whitetop, Canada thistle, Japanese brome, broom snakeweed, fringed sagewort, salsify and dandelion.</p>
<p>17. Perennial plant reproductive capability: All native plants are capable of reproducing sexually and/or vegetatively.</p>