

**NATURAL RESOURCES CONSERVATION SERVICE
MONTANA CONSERVATION PRACTICE STANDARD
FORAGE HARVEST MANAGEMENT (ACRE)**

CODE 511

DEFINITION

The timely cutting and removal of forages from the field as hay, green-chop or ensilage.

Stage of Maturity. Harvest forage at the stage of maturity that provides the desired quality and quantity without compromising plant vigor and stand longevity.

PURPOSE

- Optimize yield and quality of forage at the desired levels
- Promote vigorous plant re-growth
- Manage for the desired species composition
- Use forage plant biomass as a soil nutrient uptake tool
- Control insects, diseases and weeds
- Maintain and/or improve wildlife habitat

A final hay harvesting hay following a fall killing frost is acceptable provided adequate stubble is left to protect the soil from erosion and over-winter damage.

Native hay meadows with a high water table throughout the growing season or meadows that receive supplemental water during the growing season may be harvested for hay more than once per year. Upland range sites will be harvested for hay no more than every other year to allow adequate recovery of plant vigor. Grazing on rangeland harvested for hay is limited to the dormant season in both the harvest and following year.

CONDITIONS WHERE PRACTICE APPLIES

This practice applies to all land uses where machine harvested forage crops are grown **including seeded hayland, native grass/sedge meadows, and rangeland harvested for hay.**

If noxious weeds are present in the stand, harvest the hay prior to weed seed maturity to prevent the spread of viable seed. If herbicides are used, follow label directions for the interval of time from chemical application to safe harvest.

CRITERIA

General Criteria Applicable to All Purposes

Forage will be harvested at a frequency and height that optimizes the desired forage stand, plant community, and stand life. Follow State Cooperative extension Service (CES) recommendations or refer to **Field Office Technical Guide, Section IV, Specification, Forage Harvest Management (Code 511)** for forage harvest based on stage of maturity, moisture content, length of cut, stubble height and harvest interval. The following criteria must be met:

Moisture Content. Harvest silage/haylage crops within the optimum moisture range for the type of storage method(s) or structure(s) being utilized.

CES recommendations must be followed for optimum moisture content and levels as well as methods and techniques to monitor and/or determine moisture content and levels.

Avoid fermentation and seepage losses of digestible dry matter from direct cut hay crop silage (moisture content >70%) by treatment with chemical preservatives or add dry feedstuffs.

For optimal dry hay quality, rake hay at 30 to 40 percent moisture and turn or invert swaths when moisture is above 40 percent.

**NRCS, MT
October 2011**

Conservation practice standards are reviewed periodically and updated if needed. To obtain the current version of this standard contact the Natural Resources Conservation Service.

NOTE: This type of font (**AaBbCcDdEe 123..**) indicates NRCS National Standards.
This type of font (**AaBbCcDdEe 123..**) indicates Montana Supplement.

To preserve forage quality and quantity, bale field cured hay at 15-20 percent moisture and bale force air-dried hay at 20-35 percent moisture.

Length of Cut. When harvested for ensilage, forage will be chopped to a size appropriate for the type of storage structure used and optimal effective fiber. The length of chop selected will allow adequate packing to produce the anaerobic conditions necessary to ensure the proper ensiling process.

A shorter chop length on very dry silage may help to ensure good packing and adequate silage density.

Stubble Height. Cut forage plants at a height that will promote the vigor and health of the desired species. Cutting heights will provide adequate residual leaf area; adequate numbers of terminal, basal or auxiliary tillers or buds; insulation from extreme heat or cold; and/or un-severed stem bases that store food reserves needed for full, vigorous recovery. Follow CES recommendations for proper stubble heights to avoid winterkill of forage species in cold climates.

Contaminants. Forage shall not contain contaminants or debris that can cause illness or death to the animal being fed or rejection of the offered forage. Check CES contaminant notices, cautions, and recommendations for the specific harvest site location and area.

Additional Criteria to Improve or Maintain Stand Life, Plant Vigor and Forage Species Mix

Stage of Maturity and Harvest Interval. Cut forage plants at a stage of maturity or harvest interval range that will provide adequate food reserves and/or basal or auxiliary tillers or buds for re-growth and/or reproduction to occur without loss of plant vigor (see FOTG, Section IV, Specification, Forage Harvest Management (Code 511)).

Cut reseeding annuals at a stage of maturity and frequency that ensures the production of ample viable seed or carryover of hard seed to maintain desired stand density.

If plants show signs of short-term environmental stress, harvests will be adjusted in a manner that encourages the continued health and vigor of the

stand. Follow CES recommendations in these cases.

Manipulate timing and cutting heights of harvest to ensure germination and establishment of reseeding or seeded annuals.

Additional Criteria for Use as a Nutrient Uptake Tool

Employ a harvest regime that utilizes the maximum amount of available or targeted nutrients. Using this practice for this purpose may require more frequent harvests to increase uptake instead of managing for stand longevity.

Additional Criteria to Control Disease, Insect, Weed and Invasive Plant Infestations

Follow CES guidelines when available for control of disease, insect, weed and invasive plant infestations to forage.

Schedule harvest periods to control disease, insect, and weed infestations. When a pesticide is used to control disease, insects or weeds, adhere to the specified days to harvest period stated on the pesticide label. Evaluate pest management options by planning conservation practice standard, Pest Management (595) for all forage areas to be harvested. Also plan and schedule removal of invasive plants and noxious weeds.

Lessen incidence of disease, insect damage, and weed infestation by managing harvests to maintain a full, vigorous, dense forage stand.

Cut forages after dew, rain, or irrigation water on the leaves has evaporated.

Additional Criteria to Improve Wildlife Habitat Values

If client objectives include providing suitable habitat for desired wildlife species then appropriate harvest schedule(s), cover patterns, and minimum plant heights to provide suitable habitat for the desired specie(s) should be implemented and maintained.

Time harvests to benefit the desired wildlife species by **delaying haying (if possible) on at least part of the fields until after the peak of the nesting season (after July 15).**

Avoid cutting hay by mowing around the field. Instead, mow back and forth from one end of the

field to the other, or mow from the inside of the field to the outside. Consider using a flushing bar on swather.

Coordinate this practice with conservation practice standard, Upland Wildlife Habitat Management (645) and accompanying job sheets.

CONSIDERATIONS

Where applicable coordinate this practice with NRCS practice standard, Prescribed Grazing (528).

When nutrients or other soil amendments are applied coordinate forage harvests with NRCS practice standard, Nutrient Management (590) and/or Waste Recycling (633) as appropriate. An excess or improper balance of nutrients such as nitrogen can produce plant material that causes toxicity in some animals.

Produce stored forages of the quality needed for optimum performance of the animal being fed. Legume forages too low in fiber can lead to metabolic disorders in ruminants and an economic loss to the producer due to lowered animal performance. Consider analyzing harvested forages for feed quality. Coordinate this practice with NRCS practice standard, Feed Management (592).

Direct cut grass and legume silage can create silage leachate (seepage) in storage. Consider use of practice standards, Runoff Management System (570) and Waste Storage Facility (313).

In conjunction with harvest options, consider storage and feeding options that will retain acceptable forage quality and minimize digestible dry matter loss.

Where weather conditions make it difficult to harvest the desired quality of forage consider use of mechanical or chemical conditioners, forced air barn curing and/or ensile.

Consider delaying harvest if prolonged or heavy precipitation is forecast that would reduce forage quality.

In regions where rainfall and/or humidity levels cause unacceptable forage quality losses consider green chopping or ensiling the forage to reduce or eliminate field drying time. Other options are: the use of desiccants, preservatives, or macerating implements to reduce field-drying time.

To reduce safety hazards, avoid operating harvesting and hauling equipment on field slopes over 25 percent, particularly on cross slope traffic patterns.

Consider harvesting forages in the afternoon to optimize water soluble carbohydrates and nutritional quality.

PLANS AND SPECIFICATIONS

Place the detailed specifications in a site-specific job or design sheet or in the practice narrative in the conservation plan.

Plans and specifications must include as minimum for the forage harvest operations:

1. Goals, objectives, specific purpose (such as high forage quantity and quality or nutrient uptake, etc.)
 2. Forage species to be harvested
- By each dominant forage species harvested show:
3. Method of harvest
 4. Stage of maturity
 5. Optimal harvest moisture content
 6. Length of cut
 7. Stubble height to be left
 8. Harvest interval including late harvest if applicable
 9. Contaminant avoidance recommendations.

These plans and specifications shall be available through appropriate job sheets and other materials for applying the practice to achieve its intended purpose.

OPERATION AND MAINTENANCE

Before forage harvest, clear fields of debris that could damage machinery or if ingested by livestock, lead to sickness (for example, hardware disease) or death.

Operate all forage harvesting equipment at the optimum settings and speeds to minimize loss of leaves.

To control forage plant diseases, insects, and movement of weeds, clean harvesting equipment after harvest and before storing.

Set shear-plate on forage chopper to the proper theoretical cut for the crop being harvested. Keep knives well sharpened. Do not use re-cutters or screens unless forage moisture levels fall below recommended levels for optimum chopping action.

Follow all agricultural equipment manufacturer's safety measures when operating forage harvesting equipment.

Regardless of silage/haylage storage method, ensure good compaction and an airtight seal to exclude oxygen and mold or bacterial formations.

Dispose of the plastic wrap or bags used to store forage in an environmentally sound manner.

Fall grazing is not recommended to maintain stand productivity. However, if haylands are fall grazed, grazing should be delayed until after a killing frost.

REFERENCES

Ball, D. M., C. S. Hoveland, & G. D. Lacefield. 2002. Southern Forages. (Third Ed.). Potash & Phosphate Institute.

Barnes, R. F., C. J. Nelson., K.J. Moore & M. Collins. 2007. Forages, The Science of Grassland Agriculture, Sixth Edition.

Hanson, A. A., D. K. Barnes, & R. R. Hill, Jr. 1988. Alfalfa and Alfalfa Improvement.

Henning, J.C., M. Collins, D. Ditsch, G.D. Lacefield. 1998. Baling Forage Crops for Silage. Univ. of Ky., Ext. Pub. AGR-173, Lexington, KY.

Ishler, V. A., etal. 1991. Harvesting and Utilizing Silage. Pennsylvania State University Circular 396.

Leep R., J. Andresen, D.H. Min & A. Pollyea. 2006. Late Summer and Fall Harvest Management of Alfalfa. First Alfalfa Technology Conference. Michigan State University, East Lansing, MI.

Matches, A. G. 1973. Anti-Quality Components of Forages. Crop Science Society of America Special Pub. No. 4.

J.S. Jacobsen, S.H. Lorbeer, H.A.R. Houlton, and G.R. Carlson, Nitrogen Fertilization of Dryland Grasses in the Northern Great Plains, 1985, J. Range Management 49:340-345.

Montana State University, Animal and Range Science Department, Forage Extension Program.

<http://www.animalrangeextension.montana.edu/Forage/forage.htm>

Montana State University Extension Service, Bulletin #EB 161 "Fertilizer Guidelines for Montana", January 2003.

Montana-Utah-Wyoming Weed Management Handbook, 2006-2007. Extension Service of Montana State University, Utah State University, and University of Wyoming.

Pitt, R. E. 1990. Silage and Hay Preservation. Northeast Regional Agricultural Engineering Service.

Serotkin, N., Ed. The Penn State Agronomy Guide, 1995-1996. Pennsylvania State University. 1994. University Park, PA.

Smith, D. 1975. Forage Management in the North, Third Edition. Kendall/Hunt Publishing Company.

Taylor, N. L. 1985. Clover Science and Technology. American Society of Agronomy.