

## United States Department of Agriculture Natural Resources Conservation Service

### Ecological Site Description

#### Site Stage: Provisional

**Provisional:** an ESD at the provisional status represents the lowest tier of documentation that is releasable to the public. It contains a grouping of soil units that respond similarly to ecological processes. The ESD contains 1) enough information to distinguish it from similar and associated ecological sites and 2) a draft state and transition model capturing the ecological processes and vegetative states and community phases as they are currently conceptualized. The provisional ESD has undergone both quality control and quality assurance protocols. It is expected that the provisional ESD will continue refinement towards an approved status.

**Site Name:** Sands


**Site Type:** Rangeland

**Site ID:** R056AY090ND

**Major Land Resource Area:** 56A – Red River Valley of the North

For more information on MLRAs, refer to the following web site:

<https://www.nrcs.usda.gov/resources/data-and-reports/major-land-resource-area-mlra>



Location of MLRA 56A within Minnesota, North Dakota, and South Dakota

The Red River Valley of the North MLRA is an expansive and agriculturally important region consisting of 10,400,000 acres and including a portion of 25 counties in eastern North Dakota and northwestern Minnesota along with a small portion of the northeast corner (Roberts County) of South Dakota.

Although MLRA 56A is currently called the Red River Valley of the North, the landscape does not fit the common understanding of “valley” as the transition out of the Valley is very gradual in most places. The extent of the MLRA corresponds to the area covered by Glacial Lake Agassiz including lacustrine sediments, beach ridges, and deltas where rivers flowed into the glacial lake. Also included are island areas of glacial till which were surrounded by the lake waters. Some of the lacustrine deposits are very deep and some have glacial till within a few feet of the surface. The glaciolacustrine materials range from clayey to sandy.

The primary river in the MLRA is the Red River of the North flowing northward into Canada where it empties into Lake Winnipeg. The river is formed by the confluence of the Bois de Sioux River (flowing from northeastern South Dakota) and the Ottertail River flowing from west-central Minnesota. Numerous tributaries in MLRA 56A contribute additional water to the Red River. In Minnesota these include the Two Rivers, Snake, Marsh, Middle, Red Lake, Wild Rice, and Buffalo. In North Dakota, the Pembina, Tongue, Park, Forest, Turtle, Goose, Elm, Rush, Maple, Sheyenne, and Wild Rice are tributaries to the Red River. There are also smaller streams and coulees along with many legal drains.

The relative flatness of much of the MLRA contributes to a flooding hazard for large areas of agricultural land in the spring months. Soil salinity, while variable, also impacts land management on many areas within the MLRA. Extensive surface and subsurface (tile) drainage systems have been constructed/installed to manage excess water and/or salinity on cropland. This extensive drainage has apparently reduced ground water recharge regionally, thus impacting seasonal water table level/fluctuation and its influence on plant communities. Soils that were poorly drained prior to wide-spread drainage may now function as somewhat poorly drained or even moderately well drained soils. For example, undrained Fargo soils are Wet Meadow ecological sites; with surface drainage they may function as Subirrigated sites; and with tile drainage, they commonly function as Clayey sites. Because of the extensive alteration of the hydrology, restoration to the natural conditions of the reference state dynamics would not be possible.

MLRA 56A is an ecotone between grassland dominated MLRAs 55A and 55B to the west and forest dominated MLRAs 56B and 102A to the east. This region is utilized mostly by farms; about 80 percent is non-irrigated cropland, but some irrigated fields exist on the beach areas. Cash-grain, bean, sugar beets, potatoes, and oil production crops are the principal enterprise on many farms, but other feed grains and hay are also grown. Currently about 6 percent of this area is forested, mostly in areas along rivers that are difficult to access with farm equipment. Another 6 percent is grassland used for ranching and/or wildlife habitat. Grazing lands occur primarily in the Sand Hills area of the Sheyenne River delta, on beach areas, and on other areas too wet, saline, sodic, steep, or inaccessible to be productive cropland.

## **Ecological Site Concept**

The Sands ecological site is located on uplands – typically on sandy lake plains and deltas and on glacial lake beaches and beach ridges. Some areas are wind-worked. The soils are very deep. Surface textures typically are fine sand, loamy fine sand, or loamy sand; however, fine sandy loam and sandy loam surface layers are allowable if  $\leq 10$  inches thick. Also, on beaches coarse sandy loam surface textures may occur. Below the surface layer (to depth of more than 20 inches), the textures typically range from loamy fine sand to coarse sand. The subsoil may form a ball, but it does not ribbon. Some soils have loamy or silty material at a depth below 20 inches. Soil on this site is well drained to excessively drained. Slopes range from 0 to 15 percent. On the landscape, this site is above the Subirrigated Sands, Subirrigated, and Limy Subirrigated ecological sites. The Sandy ecological site occurs on similar landscape positions; it is fine

sandy loam or sandy loam to a depth >10 inches. The Choppy Sands site occurs on adjacent dunes or moderately steep to steep ridges where slope exceeds 15 percent.



Figure 1. Sands ecological site in association with Choppy Sands ecological site. Sands ecological site can be found down slope from Choppy Sands ecological sites.

## Physiographic Features

This site occurs on uplands – typically on sandy lake plains and deltas and on glacial lake beaches and beach ridges. The parent materials are coarse textured to a depth of 20 inches or more. Slopes range from 0 to 15 percent.

**Landform:** delta, lake plain, beach, beach ridge

	<u>Minimum</u>	<u>Maximum</u>
<b>Elevation (feet):</b>	750	1475
<b>Slope (percent):</b>	0	15
<b>Water Table Depth (inches):</b>	48	>80
<b>Flooding:</b>		
Frequency:	None	None
<b>Ponding:</b>		
Frequency:	None	None
<b>Runoff Class:</b>	Negligible	Low
<b>Aspect:</b>	No influence on this site	

## Climatic Features

MLRA 56A is considered to have a continental climate – cold winters and relatively hot summers, low to moderate humidity, light rainfall, and much sunshine. Extremes in temperature may also abound. The climate is the result of this MLRA's location near the geographic center of North America. There are few natural barriers on the northern Great Plains and air masses move freely across the plains and account for rapid changes in temperature.

Annual precipitation typically ranges from 18 to 23 inches per year. The average annual temperature is about 40°F. January is the coldest month with average temperatures ranging from about 1°F (Pembina, North Dakota (ND) to about 11°F (Wheaton, Minnesota (MN). July is the warmest month with temperatures averaging from about 68°F (Pembina, ND) to about 73°F (Wheaton, MN). The range of normal average monthly temperatures between the coldest and warmest months is about 65°F. This large annual range attests to the continental nature of this area's climate. Winds are estimated to average about 13 miles per hour annually, ranging from about 15 miles per hour during the spring to about 11 miles per hour during the summer. Daytime winds are generally stronger than nighttime and occasional strong storms may bring brief periods of high winds with gusts to more than 50 miles per hour.


Growth of cool season plants begins in early to mid-March, slowing or ceasing in late June. Warm season plants begin growth about mid-May and continue to early or mid-September. Greening up of cool season plants may occur in September and October when adequate soil moisture is present.

### Climate normals


	Representative		Actual			Average
	High	Low	High	Low		
Mean annual precipitation (in):	24	21	25	20		22
Frost free period (days):	126	102	131	87		112
Freeze free period (days):	145	132	150	126		138

	Normal monthly precipitation (in)											
	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Representative high:	0.7	0.6	1.3	2.1	3	4	3.4	2.8	3	2.2	1.1	0.7
Representative low:	0.5	0.5	0.8	1.1	2.7	3.7	3.1	2.6	1.8	1.7	0.9	0.5
Actual high:	0.8	0.7	1.5	2.2	3.2	4.1	3.4	3	3.1	2.3	1.2	0.8
Actual low:	0.5	0.4	0.8	1	2.7	3.6	3	2.5	1.8	1.6	0.8	0.5


### Normal Monthly Precipitation (in)



	Normal monthly maximum temperature (°F)											
	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Representative high:	19.4	25	36.7	55.3	69.1	77.6	82.5	80.8	71.1	56.9	38.6	22.9
Representative low:	15.7	22.1	34.5	54.3	68.5	76.7	81.3	80	69.8	53.9	34.8	19.9
Actual high:	19.9	25.1	36.8	55.9	69.7	77.7	82.7	80.8	71.4	57	38.6	23.9
Actual low:	13.4	19.3	32	51.8	65.8	74.5	79.1	78.4	67.9	52.3	33.1	18.1
Average:	17	23	35.2	54.5	68.3	76.7	81.5	80.1	70.2	55.1	36.1	21.1



	Normal monthly minimum temperature (°F)											
	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Representative high:	0.6	5.6	18.4	32	45.7	56.1	60.5	58.2	47.6	34.5	20.6	6
Representative low:	-4.4	0.5	15.1	30.8	43.3	53.9	58.4	55.8	46.2	33.1	17.9	1.7
Actual high:	0.6	5.7	18.9	33.3	46.1	56.2	60.7	58.4	48.4	35.6	20.7	6.3
Actual low:	-5.4	-0.5	13.9	29.7	41.9	52.4	56.6	53.8	43.8	31.5	16.9	1.2
Average:	-1.5	3.6	16.9	31.5	44.2	54.5	58.9	56.3	46.5	33.9	19.4	4.3



## Climate stations used

- (1) VICTOR 4 NNE [USC00398652], Rosholt, SD
- (2) PARK RIVER [USC00326857], Park River, ND
- (3) GRAFTON [USC00323594], Grafton, ND
- (4) WHEATON [USC00218907], Wheaton, MN
- (5) AGASSIZ REFUGE [USC00210050], Grygla, MN
- (6) PEMBINA [USW00014924], Pembina, ND

## Influencing Water Features

This site does not receive significant additional water, either as runoff from adjacent slopes or from a seasonal high-water table. Depth to the water table exceeds 4 feet in the spring and commonly exceeds 6 feet in the summer months. Surface infiltration is moderately rapid to very rapid. Permeability through the profile is rapid or very rapid in the sandy materials; where a contrasting texture is present below a depth 20 inches, it is moderately slow to moderate in that layer. Water loss is through percolation below the root zone and through evapotranspiration.

## Representative Soil Features

Soils associated with Sands ES are in the Mollisol and Entisol orders. The Mollisols are classified further as Calcic Hapludolls and Entic Hapludolls. The Entisols are classified further as Typic Udipsammments. These soils were developed under prairie vegetation. They formed in glaciolacustrine deposits, deltaic deposits, beach deposits, and eolian sands.

The common features of soils in this site are coarse textures within a depth of 10 inches (soil may form a ball, but it does not form a ribbon) that extend to a depth exceeding 20 inches and limited available water capacity. Most soils in the site have <5% gravel (predominantly fine gravel); however, soils on beaches can have as much as 10 percent in the upper 20 inches and higher amounts below that depth. The soils are very deep; some have medium or moderately fine textured soil materials below a depth of 20 inches and within a depth of 40 inches. They are well drained to excessively drained. The surface layer is most

commonly fine sand, loamy fine sand or loamy sand; coarse sandy loam also occurs. A few soils included in this site have a fine sandy loam or sandy loam surface layer that is <10 inches thick. Wind-worked areas may have buried A horizons.

Salinity is none in the sandy materials; where a loamy or silty substratum occurs, it may increase to slight (E.C. <4 dS/m). Sodicity is none. Soil reaction is slightly acid to moderately alkaline (pH 6.1 to 8.4). Calcium carbonate content is none to moderately low in the sandy materials; but where loamy or silty materials occur deep in the profile, it can be as high as 30 percent.

The soil surface is stable and intact. These soils are susceptible to wind erosion. Loss of the soil surface layer can result in a shift in species composition and/or production.

Major soil series correlated to the Sands site are Dickey, Lohnes, Maddock, and Serden. Also, currently included are Radium and Sandberg soils (see **Site Development and Testing Plan**).

Access Web Soil Survey (<https://websoilsurvey.sc.egov.usda.gov/App/WebSoilSurvey.aspx>) for specific local soils information.

**Parent Material Kind:** glaciolacustrine deposits, beach deposits, glaciofluvial deposits, eolian deposits

**Parent Material Origin:** lacustrine, outwash

**Surface Texture:** fine sand, loamy fine sand, loamy sand, coarse sandy loam

**Surface Texture Modifier:** None

**Subsurface Texture Group:** sandy

**Surface Fragments <3" (% Cover):** 0-10

**Surface Fragments ≥3" (% Cover):** 0

**Subsurface Fragments <3" (% Volume):** 0-10

**Subsurface Fragments ≥3" (% Volume):** 0-3

	<b>Minimum</b>	<b>Maximum</b>
<b>Drainage Class:</b>	well	excessive
<b>Permeability Class*:</b>	rapid	very rapid
<b>Depth to first restrictive layer (inches):</b>	80	>80
<b>Electrical Conductivity (dS/m)*:</b>	0	2
<b>Sodium Absorption Ratio*:</b>	0	0
<b>Soil Reaction (1:1 Water)*:</b>	6.1	8.4
<b>Soil Reaction (0.1M CaCl<sub>2</sub>):</b>	NA	NA
<b>Available Water Capacity (inches)**:</b>	2	5
<b>Calcium Carbonate Equivalent (percent)*:</b>	0	5

\*These attributes represent from 0 to >20 inches (sandy materials). Electrical Conductivity (E.C.) values are based on Saturated Paste method; the commonly used 1:1 field method will likely have E.C. values ≤1.

\*\*This attribute represents from 0-40 inches.

## Plant Communities

### Ecological Dynamics of the Site:

This ecological site description is based on nonequilibrium ecology and resilience theory and utilizes a State-and-Transition Model (STM) diagram to organize and communicate information about ecosystem change as a basis for management. The ecological dynamics characterized by the STM diagram reflect how changes in ecological drivers, feedback mechanisms, and controlling variables can maintain or induce changes in plant community composition (phases and/or states). The application of various management actions, combined with weather variables, impact the ecological processes which influence the competitive interactions, thereby maintaining or alter plant community structure.

Prior to European influence, the historical disturbance regime for MLRA 56A included frequent fires, both anthropogenic and natural in origin. Most fires, however, were anthropogenic fires set by Native Americans. Native Americans set fires in all months except perhaps January. These fires occurred in two peak periods, one from March-May with the peak in April and another from July-November with the peak occurring in October. Most of these fires were scattered and of small extent and duration. The grazing history would have involved grazing and browsing by large herbivores (such as American bison, elk, and whitetail deer). Herbivory by small mammals, insects, nematodes, and other invertebrates are also important factors influencing the production and composition of the communities. Grazing and fire interaction, particularly when coupled with drought events, influenced the dynamics discussed and displayed in the following state and transition diagram and descriptions.

Following European influence, this ecological site generally has had a history of grazing by domestic livestock, particularly cattle, which along with other related activities (e.g., fencing, water development, fire suppression) has changed the disturbance regime of the site. Changes will occur in the plant communities due to these and other factors.

Weather fluctuations coupled with managerial factors may lead to changes in the plant communities and may, under adverse impacts, result in a slow decline in vegetative vigor and composition. However, under favorable conditions the botanical composition may resemble that prior to European influence.

Five vegetative states have been identified for the site (Reference, Native/Invaded, Invaded, Wooded, and Go-Back). Within each state, one or more community phases have been identified. These community phases are named based on the more dominant and visually conspicuous species; they have been determined by study of historical documents, relict areas, scientific studies, and ecological aspects of plant species and plant communities. Transitional pathways and thresholds have been determined through similar methods.

**State 1: Reference State** represents the natural range of variability that dominated the dynamics of this ecological site prior to European influence. Dynamics of the state were largely determined by variations in climate and weather (e.g., drought), as well as that of fire (e.g., timing, frequency) and grazing by native herbivores (e.g., frequency, intensity, selectivity). Due to those variations, the Reference State is thought to have shifted temporally and spatially between two Plant Community Phases.

Presently the primary disturbances include widespread introduction of exotic plants, concentrated livestock grazing, lack of fire, and perhaps long-term non-use or very light grazing and no fire. Because of these changes, particularly the widespread occurrence of exotic species, as well as other environmental changes, the Reference State is considered to no longer exist. Thus, the presence of exotic plants on the site precludes it from being placed in the Reference State. It must then be placed in one of the other states, commonly State 2: Native/Invaded State (T1A).

**State 2: Native/Invaded State.** Colonization of the site by exotic plants results in a transition from State 1: Reference State to State 2: Native/Invaded State (T1A). This transition was inevitable; it often resulted from colonization by exotic cool-season grasses (e.g., Kentucky bluegrass, smooth brome, crested wheatgrass, quackgrass) which have been particularly and consistently invasive under long-term non-use or very light grazing and no fire. Other exotic plants (e.g., Canada thistle, leafy spurge) are also known to invade the site.

Three community phases have been identified for this state; they are similar to the community phases in the Reference State but have now been invaded by exotic cool-season grasses. These exotic cool-season grasses can be expected to increase. As that increase occurs, plants more desirable to wildlife and livestock may decline. A decline in forb diversity can also be expected. Under non-use or minimal

use management, mulch increases and may become a physical barrier to plant growth. This also changes the micro-climate near the soil surface and may alter infiltration, nutrient cycling, and biological activity near the soil surface. As a result, these factors, coupled with shading, cause desirable native plants to have increasing difficulty remaining viable and recruitment declines.

To slow or limit the invasion of these exotic grasses or other exotic plants, it is imperative that managerial techniques (e.g., prescribed grazing, prescribed burning) be carefully constructed, monitored, and evaluated with respect to that objective. If management does not include measures to control or reduce these exotic plants, the transition to State 3: Invaded State should be expected (T2B). This state may also transition to State 4: Invaded Wooded State during long-term non-use or very light grazing and no fire (T2A).

**State 3: Invaded State.** The threshold for this state is reached when both the exotic cool-season grasses exceed 30% of the plant community and native grasses represent less than 40% of the community. One plant community phase has been identified for this state.

The exotic cool-season grasses can be quite invasive and often form monotypic stands. As they increase, both forage quantity and quality of the annual production becomes increasingly restricted to late spring and early summer, even though annual production may increase. Forb diversity often declines. Under non-use or minimal use management, mulch can increase and become a physical barrier to plant growth, which alters nutrient cycling, infiltration, and soil biological activity. As such, desirable native plants become increasingly displaced.

Once the state is well established, prescribed burning and prescribed grazing techniques have been largely ineffective in suppressing or eliminating the exotic cool-season grasses, even though some short-term reductions may appear successful. However, assuming there is an adequate component of native grasses to respond to treatments, a restoration pathway to State 2: Native/Invaded State (R3A) may be accomplished with the implementation of long-term prescribed grazing in conjunction with prescribed burning. Long-term non-use or very light grazing and no fire may transition to State 4: Invaded Wooded State T3A).

**State 4: Invaded Wooded State.** This state historically existed as small patches of trees and/or shrubs scattered across the site when precipitation, fire frequency, and other factors enabled woody species to colonize or encroach on the site. This often resulted in a mosaic of patches of woody vegetation interspersed within the grass dominated vegetation. A marked increase in non-use management and active fire suppression since European influence has enabled this state to expand and become more widespread. One community phase has been identified and often results from long-term non-use or very light grazing and no fire (T2A, R3A).

Brush control (i.e., prescribed burning, chemical, or mechanical treatments) may lead to State 2: Native/Invaded State (R4A). However, depending upon the abundance of exotic cool-season grasses, a range planning may be necessary following the brush control to complete the restoration effort. Brush control can also lead to State 3: Invaded State (R4B).

**State 5: Go-Back State** often results following cropland abandonment and consists of only one plant community phase. This weedy assemblage may include noxious weeds that need control. Over time, the exotic cool-season grasses (e.g., Kentucky bluegrass, smooth brome, crested wheatgrass, quackgrass) will likely predominate.

Initially, due to extensive bare ground and a preponderance of shallow rooted annual plants, infiltration is low and the potential for soil erosion is high. Plant species richness may be high, but overall diversity (i.e., equitability) is typically low, with the site dominated by a relatively small assemblage of species. Due to


the lack of native perennials and other factors, restoring the site with the associated ecological processes is difficult. However, a successful range planting may result in something approaching State 2: Native/Invaded State (R5A). Following seeding, prescribed grazing, prescribed burning, haying, and the use of herbicides will generally be necessary to achieve the desired result and control weeds, some of which may be noxious weeds. A failed range planting and/or secondary succession will lead to State 3: Invaded State (R5B).

The following state and transition model diagram illustrates the common states, community phases, community pathways, and transition and restoration pathways that can occur on the site. These are the most common plant community phases and states based on current knowledge and experience; changes may be made as more data are collected. Pathway narratives describing the site's ecological dynamics reference various management practices (e.g., prescribed grazing, prescribed fire, brush management, herbaceous weed treatment) which, if properly designed and implemented, will positively influence plant community competitive interactions. The design of these management practices will be site specific and should be developed by knowledgeable individuals; based upon management goals and a resource inventory; and supported by an ongoing monitoring protocol.

When the management goal is to maintain an existing plant community phase or restore to another phase within the same state, modification of existing management to ensure native species have the competitive advantage may be required. To restore a previous state, the application of two or more management practices in an ongoing manner will be required. Whether using prescribed grazing, prescribed burning, or a combination of both with or without additional practices (e.g., brush management), the timing and method of application needs to favor the native species over the exotic species. Adjustments to account for variations in annual growing conditions and implementing an ongoing monitoring protocol to track changes and adjust management inputs to ensure desired outcome will be necessary.

The plant community phase composition table(s) has been developed from the best available knowledge including research, historical records, clipping studies, and inventory records. As more data are collected, plant community species composition and production information may be revised.

## Plant Communities and Transitional Pathways



**Diagram Legend - MLRA 56A Sands**

T1A	Introduction of exotic plants
T2A	Long-term non-use or very light grazing, no fire, long-term above average precipitation
T2B	Light grazing, heavy season-long grazing, or no use and no fire
T3A	Long-term non-use or very light grazing and no fire
T6A	Cessation of annual cropping
R3A	Long-term prescribed grazing and prescribed burning
R4A	Brush control and perhaps range planting
R4B	Brush control
R5A	Successful range planting
R5B	Failed range planting and/or secondary succession
CP 1.1 - 1.2 (1.1A)	Periods of below average precipitation and increased disturbance
CP 1.2 - 1.1 (1.2A)	Return to average precipitation and disturbance regime
CP 2.1 - 2.2 (2.1A)	Heavy grazing
CP 2.2 - 2.1 (2.2A)	Long-term prescribed grazing and prescribed burning
CP 2.2 - 2.3 (2.2B)	Heavy grazing
CP 2.3 - 2.2 (2.3A)	Long-term prescribed grazing and prescribed burning

**State 1: Reference State**

This state represented the natural range of variability that dominated the dynamics of this ecological site prior to European influence. The primary disturbance mechanisms for this site in the reference condition included frequent fire and grazing by large herding ungulates. Timing of fires and grazing, coupled with weather events, dictated the dynamics that occurred within the natural range of variability. These factors likely caused the community to shift both spatially and temporally between two community phases.

**Characteristics and indicators** (i.e., characteristics and indicators that can be used to distinguish this state from others). Because of changes in disturbances and other environmental factors (particularly the widespread occurrence of exotic species), the Reference State is considered to no longer exist.

**Resilience management** (i.e., management strategies that will sustain a state and prevent a transition). If intact, the reference state should probably be managed with current disturbance regimes which has permitted the site to remain in reference condition as well as maintaining the quality and integrity of associated ecological sites. Maintenance of the reference state is contingent upon a monitoring protocol to guide management.

**Community Phase 1.1: Prairie Sandreed-Porcupinegrass-Bluestems (*Calamovilfa longifolia*-*Hesperostipa spartea*-*Andropogon* spp., *Schizachyrium scoparium*)**

This community phase was historically the most dominant both temporally and spatially. The major grasses and sedges included prairie sandreed, porcupinegrass, sand bluestem, and big bluestem. Other associated grasses included needle and thread, little bluestem, sideoats grama, and sand dropseed. Common forbs included Cuman ragweed, field sagewort, white sagebrush, milkweed, field chickweed, Canada horseweed, and purple prairie clover. Leadplant, western sandcherry, prairie sagewort, and prairie rose were common shrubs.

Annual production likely varied from about 1800-3200 pounds per acre with grasses and grass-likes, forbs, and shrubs contributing about 85%, 10% and 5%, respectively. Because both warm-season and cool-season grasses were well represented in the community, production would have been distributed throughout the growing season. This community represents the plant community phase upon which

Site Type: Rangeland  
MLRA: 56A – Red River Valley of the North

**Sands**  
**R056AY090ND**

interpretations are primarily based and is described in the “Plant Community Composition and Group Annual Production” portion of this ecological site description.

Plant Community Composition and Group Annual Production				
1.1 Prairie Sandreed-Porcupinegrass-Bluestems				
COMMON/GROUP NAME	SYMBOL	Group	lbs./acre	% Comp
<b>GRASSES &amp; GRASS-LIKES</b>			1950 - 2210	75 - 85
<b>TALL WARM-SEASON GRASSES</b>		1	<b>780 - 1170</b>	<b>30 - 45</b>
big bluestem	ANGE	1	130 - 260	5 - 10
prairie sandreed	CALO	1	130 - 390	5 - 15
sand bluestem	ANHA	1	260 - 520	10 - 20
<b>MID COOL-SEASON BUNCH</b>		2	<b>390 - 520</b>	<b>15 - 20</b>
Canada wildrye	ELCA4	2	26 - 52	1 - 2
needle and thread	HECOC8	2	130 - 390	5 - 15
porcupinegrass	HESP11	2	26 - 78	1 - 3
<b>MID WARM-SEASON</b>		3	<b>130 - 390</b>	<b>5 - 15</b>
little bluestem	SCSC	3	130 - 260	5 - 10
sand dropseed	SPCR	3	26 - 130	1 - 5
sideoats grama	BOCU	3	130 - 260	5 - 10
<b>SHORT WARM-SEASON</b>		4	<b>26 - 130</b>	<b>1 - 5</b>
blue grama	BOGR2	4	26 - 130	1 - 5
<b>OTHER NATIVE GRASSES</b>		5	<b>26 - 130</b>	<b>1 - 5</b>
prairie Junegrass	KOMA	5	26 - 130	1 - 5
Canada wildrye	ELCA4	5	0 - 78	0 - 3
western wheatgrass	PASM	5	0 - 78	0 - 3
Scribner rosette grass	DIOLS	5	0 - 78	0 - 3
other perennial grasses	2GP	5	0 - 130	0 - 5
<b>GRASS-LIKES</b>		6	<b>130 - 260</b>	<b>5 - 10</b>
Pennsylvania sedge	CAPE6	6	26 - 78	1 - 3
sun sedge	CAINH2	6	26 - 130	1 - 5
sedge	CAREX	6	0 - 78	0 - 3
Schweinitz's flatsedge	CYSC5	6	0 - 26	0 - 1
other grass-likes	2GL	6	0 - 130	0 - 5
<b>FORBS</b>		7	<b>130 - 260</b>	<b>5 - 10</b>
Cuman ragweed	AMPS	7	26 - 52	1 - 2
field sagewort	ARCA12	7	26 - 52	1 - 2
white sagebrush	ARLU	7	26 - 52	1 - 2
milkweed	ASCLE	7	26 - 52	1 - 2
field chickweed	CLEAR4	7	26 - 52	1 - 2
Canadian horseweed	COCAC3	7	26 - 52	1 - 2
purple prairie clover	DAPU5	7	26 - 52	1 - 2
silky prairie clover	DAVI	7	26 - 52	1 - 2
western wallflower	ERAS2	7	26 - 52	1 - 2
hairy false goldenaster	HEVIV	7	26 - 52	1 - 2
dotted blazing star	LIPU	7	26 - 52	1 - 2
hoary puccoon	LICA12	7	26 - 52	1 - 2
narrowleaf stoneseed	LIIN2	7	26 - 52	1 - 2
rush skeletonweed	LYJU	7	26 - 52	1 - 2
silverleaf Indian breadroot	PEAR6	7	26 - 52	1 - 2
large beardtongue	PEGR7	7	26 - 52	1 - 2
lilac beardtongue	PEGRG3	7	26 - 52	1 - 2
groundcherry	PHHI8	7	26 - 52	1 - 2
upright prairie coneflower	RACO3	7	26 - 52	1 - 2
Missouri goldenrod	SOMI2	7	26 - 52	1 - 2
white heath aster	SYER	7	26 - 52	1 - 2
prairie spiderwort	TROC	7	26 - 52	1 - 2
wild onion	ALLU	7	0 - 26	0 - 1
thymeleaf sandmat	CHSE6	7	0 - 26	0 - 1
Flodman's thistle	CIFL	7	0 - 26	0 - 1
smooth horsetail	EQLA	7	0 - 26	0 - 1
flat-top goldenrod	EUGR5	7	0 - 26	0 - 1
heartleaf four o'clock	MINY	7	0 - 26	0 - 1
cinquefoil	POTEN	7	0 - 26	0 - 1
primrose	PRIMU	7	0 - 26	0 - 1
gray goldenrod	SONE	7	0 - 26	0 - 1
hoary verbena	VEST	7	0 - 26	0 - 1
prairie violet	VIPE2	7	0 - 26	0 - 1
native forbs	2FN	7	26 - 130	1 - 5
<b>SHRUBS</b>		8	<b>26 - 130</b>	<b>1 - 5</b>
leadplant	AMCA6	8	26 - 78	1 - 3
western sandcherry	PRPUB	8	26 - 78	1 - 3
prairie sagewort	ARFR4	8	26 - 52	1 - 2
prairie rose	ROAR3	8	26 - 52	1 - 2
western snowberry	SYOC	8	26 - 52	1 - 2
white meadowsweet	SPAL2	8	0 - 52	0 - 2
poison ivy	TORY	8	0 - 26	0 - 1
other shrubs	ZSHRUB	8	26 - 78	1 - 3
<b>Annual Production lbs./acre</b>		LOW	RV	HIGH
<b>GRASSES &amp; GRASS-LIKES</b>		1650 -	2327	2790
<b>FORBS</b>		125 -	195	275
<b>SHRUBS</b>		25 -	78	135
<b>TOTAL</b>		1800 -	2600	3200

This list of plants and their relative proportions are based on near normal years. Fluctuations in species composition and relative production may change from year to year dependent upon precipitation or other climatic factors. RV = Representative Value.

### **Community Phase Pathway 1.1A**

Community Phase Pathway 1.1 to 1.2 occurred during periods of below average precipitation with increased disturbance resulting in increases of needle and thread and blue grama with corresponding decreases in bluestems and porcupinegrass.

### **Community Phase 1.2: Needle and Thread-Prairie Sandreed-Blue Grama (*Hesperostipa comata*-*Calamovilfa longifolia*-*Bouteloua gracilis*)**

This plant community resulted from periods of below average precipitation with increased disturbance (such as heavy, frequent grazing over a period of several years and/or several consecutive years of below average precipitation). This increase in grazing pressure may have resulted from proximity to a water source, changes in fire frequency, and/or prolonged drought.

Compared to Community Phase 1.1, needle and thread would have displaced porcupinegrass to become the dominant needlegrass while blue grama, sand dropseed, and sedges would have also increased. Prairie sandreed and the bluestems would have decreased. Forb species (such as field sagewort, goldenrod, Cuman ragweed, and common yarrow) would have increased. Annual production would have been somewhat reduced compared to Community Phase 1.1.

### **Community Phase Pathway 1.2A**

Community Phase Pathway 1.2 to 1.1 occurred upon the return to average precipitation and disturbance regime, which allowed for the recovery of the bluestems, porcupinegrass and prairie sandreed.

### **Transition T1A**

This is the transition from the State 1: Reference State to the State 2: Native/Invaded State due to the introduction and establishment of exotic cool-season grasses (e.g., Kentucky bluegrass, smooth brome, crested wheatgrass, quackgrass). This transition was inevitable and corresponded to a decline in native warm-season and cool-season grasses. It may have been exacerbated by chronic season-long or heavy late season grazing. Complete rest from grazing and suppression of fire could also have hastened the transition. The threshold between states was crossed when Kentucky bluegrass, smooth brome, crested wheatgrass, quackgrass, or other exotic plants became established on the site.

**Constraints to recovery** (i.e., variables or processes that preclude recovery of the former state). Current knowledge and technology will not facilitate a successful restoration to Reference State.

### **State 2: Native/Invaded State**

This state is similar to State 1: Reference State but has now been colonized by the exotic cool-season grasses (e.g., Kentucky bluegrass, smooth brome, crested wheatgrass, quackgrass) which are now present in small amounts. Although the state is still dominated by native grasses, an increase in these exotic cool-season grasses can be expected.

These exotic cool-season grasses can be quite invasive on the site and are particularly well adapted to heavy grazing. They also often form monotypic stands. As these exotic cool-season grasses increase, both forage quantity and quality become increasingly restricted to late spring and early summer due to the monotypic nature of the stand, even though annual production may increase. Native forbs generally decrease in production, abundance, diversity, and richness compared to that of State 1: Reference State.

These exotic cool-season grasses have been particularly and consistently invasive under extended periods of non-use and no fire. To slow or limit the invasion of these exotic grasses, it is imperative that managerial techniques (e.g., prescribed grazing, prescribed burning) be carefully constructed, monitored,

and evaluated with respect to that objective. If management does not include measures to control or reduce these exotic cool-season grasses, the transition to State 3: Invaded State should be expected.

Annual production of this state can be quite variable, in large part due to the amount of exotic cool-season grasses. However, annual production may be similar to that of State 1: Reference State (i.e., 1800-3200 pounds per acre).

**Characteristics and indicators** (i.e., characteristics that can be used to distinguish this state from others). The presence of trace amounts of exotic cool-season grasses indicates a transition from State 1 to State 2. The presence of exotic biennial or perennial leguminous forbs (i.e., sweet clover, black medic) may not, on their own, indicate a transition from State 1 to State 2 but may facilitate that transition.

**Resilience management** (i.e., management strategies that will sustain a state and prevent a transition). To slow or limit the invasion of these exotic grasses, it is imperative that managerial techniques (e.g., prescribed grazing, prescribed burning) be carefully constructed, monitored, and evaluated with respect to that objective.

Grazing management should be applied that enhances the competitive advantage of native grass and forb species. This may include: (1) grazing when exotic cool-season grasses are actively growing and native cool-season grasses are dormant; (2) applying proper deferment periods allowing native grasses to recover and maintain or improve vigor; (3) adjusting overall grazing intensity to reduce excessive plant litter (above that needed for rangeland health indicator #14 – see Rangeland Health Reference Worksheet); (4) incorporating early heavy spring utilization which focuses grazing pressure on exotic cool-season grasses and reduces plant litter, provided that livestock are moved when grazing selection shifts from exotic cool-season grasses to native grasses.

Prescribed burning should be applied in a manner that maintains or enhances the competitive advantage of native grass and forb species. Prescribed burns should be applied as needed to adequately reduce/remove excessive plant litter and maintain the competitive advantage for native species. Timing of prescribed burns (spring vs. summer vs. fall) should be adjusted to account for differences in annual growing conditions and applied during windows of opportunity to best shift the competitive advantage to the native species.

### **Community Phase 2.1: Prairie Sandreed-Porcupinegrass-Bluestems (*Calamovilfa longifolia*-*Hesperostipa spartea*-*Andropogon* spp., *Schizachyrium scoparium*)**

This community phase is similar to Community Phase 1.1 but has been colonized by exotic cool-season grasses (e.g., Kentucky bluegrass, smooth brome, crested wheatgrass, quackgrass). However, these exotics are present in smaller amounts with the community still dominated by native grasses. Annual production is in the range of 1800-3200 pounds per acre.

Grazing systems that allow for adequate recovery periods following grazing events and, potentially, the combination of prescribed grazing and prescribed burning which closely mimics the natural disturbance regime will help maintain this community phase.



Figure 2. Community Phase 2.1: Prairie Sandreed-Porcupinegrass-Bluestems

### **Community Phase Pathway 2.1A**

Community Phase Pathway 2.1 to 2.2 occurs during periods of heavy grazing which causes a shift from the dominance of prairie sandreed and porcupinegrass to the more grazing-tolerant needle and thread, blue grama, sun sedge, and minor amounts of Kentucky Bluegrass.

### **Community Phase 2.2: Needle and Thread-Prairie Sandreed-Blue Grama/Sun Sedge/Forbs (*Hesperostipa comata*-*Calamovilfa longifolia*-*Bouteloua gracilis*/*Carex inops*/Forbs)**

This plant community phase is characterized by a decline in porcupinegrass, sand bluestem, and corresponding increases in the more grazing tolerant needle and thread, sand dropseed, sedges and Kentucky bluegrass. Forbs (such as common yarrow, goldenrod, field sagewort, and white sagebrush) will increase while the leguminous forbs may decrease. The shrub component remains similar; however, prairie sagewort may increase.



Figure 3. Community Phase 2.2: Needle and Thread-Prairie Sandreed-Blue Grama/Sun Sedge/Forbs

### **Community Phase Pathway 2.2A**

Community Phase Pathway 2.2 to 2.1 is initiated with the implementation of long-term prescribed grazing and prescribed burning. If properly implemented, this will shift the competitive advantage away from the exotic cool-season species to the mid statured bunchgrasses and tall statured rhizomatous grasses. Properly timed prescribed burning may expedite this shift.

### **Community Phase Pathway 2.2B**

Community Phase Pathway 2.2 to 2.3 occurs during periods heavy grazing, leading to marked increases in short statured grasses and sedges (i.e., blue grama, sun sedge) with corresponding decreases in prairie sandreed and the needlegrasses.

### **Community Phase 2.3: Blue Grama/Sun Sedge/Exotic Cool-Season Grasses/Forbs (*Bouteloua gracilis/Carex inops*/Exotic Cool-Season Grasses/Forbs)**

This community phase results from heavy grazing over a period of several years. Blue grama becomes the dominant grass with lesser amounts of upland sedges and Kentucky bluegrass. Grazing tolerant forbs also increase. The needlegrasses and tall statured warm-season grasses are likely still present but may not be readily observable. Conspicuous forbs often include goldenrod, field sagewort, yellow salsify, white heath aster, common yarrow, and Cuman ragweed. Prairie sagewort and prairie rose are the principal shrubs.

This community phase is approaching the threshold leading to a transition to State 3: Invaded State. As a result, it is an “at risk” community. If management does not include measures to control or reduce these exotic cool-season grasses, the transition to State 3: Invaded State should be expected.

### **Community Phase Pathway 2.3A**

This community phase pathway 2.3 to 2.2 is initiated by implementation of long-term prescribed grazing and prescribed burning. If properly implemented, this will shift the competitive advantage from the exotic cool-season species to the remnant native cool-season bunchgrass species and tall statured warm-season grasses. Kentucky bluegrass will remain in this community at varying amounts depending on the level of management. Caution should be exercised when initiating this restoration pathway to ensure that management actions do not favor Kentucky bluegrass resulting in unexpectedly crossing the threshold to State 3: Invaded State.

### **Transition T2A**

This transition from the State 2: Native/Invaded to State 4: Invaded Wooded State generally occurs during long-term non-use or very light grazing, and no fire. It may also occur with long-term above average precipitation, particularly when an abundant shrub component is present. It has also become more frequent following European settlement when the historic fire regime was markedly reduced.

**Constraints to recovery** (i.e., variables or processes that preclude recovery of the former state). The extended fire interval may make recovery doubtful due to the abundance of exotic cool-season grasses and lack of native grasses. Fire intensity along with consumption of available fuels may cause incomplete or patchy burns. Continued recruitment of tree seeds from adjacent sites will hamper site restoration.

Reticence to undertake tree removal and the perception that trees may be a desirable vegetation component for wildlife habitat, carbon sequestration, aesthetics, etc. are some of the constraints to recovery. Managers wanting to manage the site for deer, livestock, or grassland nesting birds will need to consider the intensive management required to restore and maintain the site in State 2. The disturbance regime necessary to restore this site to State 2: Native/Invaded State is very labor intensive and costly; therefore, addressing woody removal earlier in the encroachment phase is the most cost-effective treatment for woody control.

### **Transition T2B**

This transition from the State 2: Native/Invaded State to State 3: Invaded State generally occurs with light grazing, heavy season-long grazing, or periods of non-use and no fire especially when the shrub component is sparse. Exotic cool-season grasses, such as Kentucky bluegrass and/or smooth brome, often become the dominant graminoids. Studies indicate that a threshold may exist in this transition when both the exotic cool-season grasses exceed 30% of the plant community and native grasses represent less than 40% of the plant community composition.

**Constraints to recovery** (i.e., variables or processes that preclude recovery of the former state).

Variations in growing conditions (e.g., cool, wet spring) will influence effects of various management activities on exotic cool-season grass populations. The opportunity for high intensity spring burns (which can serve to reduce the introduced cool-season species) is severely reduced by early green-up and the lack of fuel.

### **State 3: Invaded State**

This state is the result of invasion and dominance by exotic cool-season grasses (e.g., Kentucky bluegrass, smooth brome, crested wheatgrass, quackgrass). Other exotic plants (e.g., leafy spurge, Canada thistle) are also known to invade the site. The exotic cool-season grasses can be quite invasive on the site and are particularly well adapted to heavy grazing. They often form monotypic stands. As

these exotic cool-season grasses increase, both forage quantity and quality become increasingly restricted to late spring and early summer due to the monotypic nature of the stand, even though annual production may increase. Native forbs generally decrease in production, abundance, diversity, and richness compared to that of State 1: Reference State. Common forbs often include white heath aster, goldenrod, common yarrow, and white sagebrush. Shrubs, such as western snowberry and rose, may show marked increases. Once the state is well established, prescribed burning and grazing techniques have been largely ineffective in suppressing or eliminating these species, even though some short-term reductions may appear successful.

Annual production of this state may vary widely, in part due to variations in the extent of invasion by exotic cool-season grasses. However, annual production may be in the range of 2800-3800 pounds per acre with the exotic cool-season grasses contributing the bulk of the production.

**Characteristics and indicators** (i.e., characteristics that can be used to distinguish this state from others). This site is characterized by exotic cool-season grasses constituting greater than 30 percent of the annual production and native grasses constituting less than 40 percent of the annual production.

**Resilience management** (i.e., management strategies that will sustain a state and prevent a transition). Light or moderately stocked continuous, season-long grazing or a prescribed grazing system which incorporates adequate deferment periods between grazing events and proper stocking rate levels will maintain this State. Application of herbaceous weed treatment, occasional prescribed burning, and/or brush management may be needed to manage noxious weeds and increasing shrub (e.g., western snowberry) populations.

### **Community Phase 3.1: Exotic Cool-Season Grasses/Forbs**

This community phase is typically dominated by exotic grasses (often Kentucky bluegrass, but perhaps smooth brome, crested wheatgrass, and/or quackgrass). Exotic forbs, such as leafy spurge or Canada thistle, may also be present. Grazing tolerant native forbs (such as white sagebrush, Cuman ragweed, and common yarrow) are often noticeably present. Left untreated, leafy spurge will continue to increase which will reduce the remnant native grasses and eventually reduce the Kentucky bluegrass. The longer this community phase exists, the more resilient it becomes. Natural or management disturbances that reduce the cover of Kentucky bluegrass are short-lived.



Figure 4. Foreground - Community Phase 3.1: Exotic Cool-Season Grasses/Forbs

### **Restoration R3A**

This restoration pathway from State 3: Invaded State to State 2: Native/Invaded State may be accomplished with the implementation of long-term prescribed grazing and prescribed burning, assuming there is an adequate component of native grasses to respond to the treatments.

Both prescribed grazing and prescribed burning are likely necessary to successfully initiate this restoration pathway, the success of which depends upon the presence of a remnant population of native grasses in Community Phase 3.1. That remnant population, however, may not be readily apparent without close inspection. The application of several prescribed burns may be needed at relatively short intervals in the early phases of this restoration process, in part because many of the shrubs (e.g., western snowberry) sprout profusely following one burn. Early season prescribed burns have been successful, however, fall burning may also be an effective technique.

The prescribed grazing should include adequate recovery periods following each grazing event and stocking levels which match the available resources. If properly implemented, this will shift the competitive advantage from the exotic cool-season grasses to the native cool-season grasses.

**Context dependence** (i.e., factors that cause variations in plant community shifts, restoration likelihood, and contribute to uncertainty). Grazing management should be applied in a manner that enhances/maximizes the competitive advantage of native grass and forb species over the exotic species. This may include the use of prescribed grazing to reduce excessive plant litter accumulations above that needed for rangeland health indicator #14 (see Rangeland Health Reference Worksheet). Increasing livestock densities may facilitate the reduction in plant litter provided length and timing of grazing periods are adjusted to favor native species.

Grazing prescriptions designed to address exotic grass invasion and favor native species may involve earlier, short, intense grazing periods with proper deferment to improve native species health and vigor. Fall (e.g., September, October) prescribed burning followed by an intensive, early spring graze period with adequate deferment for native grass recovery may shift the competitive advantage to the native species, facilitating the restoration to State 2: Native/Invaded.

Prescribed burning should be applied in a manner that enhances the competitive advantage of native grass and forb species over the exotic species. Prescribed burns should be applied at a frequency which mimics the natural disturbance regime, or more frequently as is ecologically (e.g., available fuel load) and economically feasible. Burn prescriptions may need adjustment to: (1) account for change in fine fuel orientation (e.g., “flopped” Kentucky bluegrass); (2) fire intensity and duration by adjusting ignition pattern (e.g., backing fires vs head fires); (3) account for plant phenological stages to maximize stress on exotic species while favoring native species (both cool- and warm-season grasses).

The longer this community phase exists, the more resilient it becomes. Natural or management disturbances that reduce the cover of Kentucky bluegrass or smooth brome are typically short-lived.

### **Transition T3A**

This transition from State 3: Invaded State to State 4: Invaded Wooded State occurs with long-term non-use or very light grazing and no fire. It frequently occurs when the site is in close proximity to wooded areas where the shrubs may encroach vegetatively upon the site and/or serve as a seed source for these species to colonize the site. It has also become more frequent following European settlement when the historic fire regime was markedly reduced.

**Constraints to recovery** (i.e., variables or processes that preclude recovery of the former state). The extended fire interval may make recovery doubtful due to the abundance of exotic cool-season grasses and lack of native grasses. Fire intensity along with consumption of available fuels may cause incomplete or patchy burns. Continued recruitment of tree seeds from adjacent sites will hamper site restoration.

Constraints to recovery include reticence to undertake tree removal and the perception that trees may be a desirable vegetation component for wildlife habitat, carbon sequestration, aesthetics, etc. Managers wanting to manage the site for deer, livestock, or grassland nesting birds will need to consider the intensive management required to restore and maintain the site in State 2. The disturbance regime necessary to restore this site to State 2: Native/Invaded State is very labor intensive and costly; therefore, addressing woody removal earlier in the encroachment phase is the most cost-effective treatment for woody control.

### **State 4: Invaded Wooded State**

This state historically existed as small patches of shrubs (e.g., smooth sumac, western snowberry, chokecherry, and/or wild rose) scattered across the site, particularly where these shrubs could have encroached onto the site vegetatively (e.g., rhizomes, root sprouts) or by seed from nearby woody vegetation.

**Characteristics and indicators** (i.e., characteristics and indicators that can be used to distinguish this state from others). The dominance of woody species (by cover and production) distinguishes this state from other herbaceously dominated states.

**Resilience management** (i.e., management strategies that will sustain a state and prevent a transition). This state is resistant to change in the long-term absence of fire. Restoration efforts would require the use of prescribed fire, mechanical treatment, and prescribed grazing. Considerable time and effort will be required to restore to other States.

### **Community Phase 4.1: Shrubs/Exotic Grasses/Forbs**

This community phase results from long-term non-use or very light grazing and no fire, particularly during times of above average precipitation. Common woody species often include smooth sumac, western snowberry, chokecherry, and wild rose. A marked increase in non-use management and active fire

suppression since European influence has enabled this state to expand and become more widespread. Exotic cool-season grasses such as Kentucky bluegrass generally increase along with woody species.

### **Restoration R4A**

This restoration pathway from State 4: Invaded Wooded State to State 2: Native/Invaded State can be accomplished with brush control. However, depending upon level of remnant native grasses and forbs, a range planting may also be necessary to re-establish the herbaceous plant community, thus completing the restoration.

A combination of brush control methods may be necessary (e.g., chemical, mechanical, prescribed burning). However, initially, low fuel loads may preclude the application of prescribed burns. As a result, herbicides and/or mechanical brush control may be necessary to initiate the restoration effort.

Subsequently, several prescribed burns may be needed at relatively short intervals, in part because many of the shrubs sprout profusely following one burn.

If a range planting is necessary, it may be possible using selected plant materials and agronomic practices to approach something very near the functioning of State 2: Native/Invaded State. Application of chemical herbicides and the use of mechanical planting methods using adapted varieties of the dominant native grasses are possible and can be successful.

**Context dependence** (i.e., factors that cause variations in plant community shifts, restoration likelihood, and contribute to uncertainty). Prescribed burning should be applied in a manner that enhances the competitive advantage of native grass and forb species over the exotic species. Prescribed burns should be applied at a frequency which mimics the natural disturbance regime or more frequently as is ecologically (e.g., available fuel load) and economically feasible. Burn prescriptions may need adjustment to: (1) account for change in fuel type (herbaceous vs. shrub vs. tree), fine fuel amount and orientation; (2) fire intensity and duration by adjusting ignition pattern (e.g., backing fires vs head fires); (3) account for plant phenological stages to maximize stress on woody and exotic species while favoring native species (both cool- and warm-season grasses).

The method of brush management will be site specific, but generally the goal would be to apply the pesticide, mechanical control or biological control (either singularly or in combination) in a manner that shifts the competitive advantage from the targeted species to the native grasses and forbs. The control method(s) should be as specific to the targeted species as possible to minimize impacts to non-target species.

A successful range planting will include proper seedbed preparation, weed control (both prior to and after the planting), selection of adapted native species representing functional/structural groups inherent to the State 1, and proper seeding technique. Management (e.g., prescribed grazing, prescribed burning) during and after establishment must be applied in a manner that maintains the competitive advantage for the seeded native species. Adding non-native species can impact the above and below ground biota. Some evidence suggests the addition of exotic legumes to the seeding mixture may favor exotic cool-season grass expansion/invasion.

### **Restoration R4B**

This restoration from State 4: Invaded Wooded State to State 3: Invaded State can be accomplished with brush control. Initial use of herbicides and/or mechanical brush control to reduce smooth sumac and other shrubs may be necessary to increase fine fuel loads to enable the application of prescribed burning to further control sprouting shrubs species.

**Context dependence** (i.e., factors that cause variations in plant community shifts, restoration likelihood, and contribute to uncertainty). The method of brush management will be site specific, but generally the goal would be to apply the pesticide, mechanical control or biological control (either singularly or in combination) in a manner that shifts the competitive advantage from the targeted species to the grasses and forbs. The control method(s) should be as specific to the targeted species as possible to minimize impacts to non-target species.

Continued recruitment of seeds of trees and shrubs from adjacent sites will hamper site restoration. Intensive management is required to restore and maintain the site in State 3: Invaded State.

### **State 5: Go-Back State**

This state is highly variable depending on the level and duration of disturbance related to the T6A transitional pathway. In this MLRA, the most probable origin of this state is plant succession following cropland abandonment. This plant community will initially include a variety of annual forbs and grasses, some of which may be noxious weeds needing control. Over time, however, the exotic cool-season grasses (e.g., Kentucky bluegrass, smooth brome, quackgrass, crested wheatgrass) will likely predominate.

**Characteristics and indicators** (i.e., characteristics that can be used to distinguish this state from others). Tillage has destroyed the native plant community, altered soil structure and biology, reduced soil organic matter, and resulted in the formation of a tillage induced compacted layer which is restrictive to root growth. Removal of perennial grasses and forbs results in decreased infiltration and increased runoff.

**Resilience management** (i.e., management strategies that will sustain a state and prevent a transition). Continued tillage will maintain the state. Control of noxious weeds will be required.

### **Community Phase 5.1: Annual/Pioneer Perennial/Exotics**

This state is highly variable depending on the level and duration of disturbance related to the T6A transitional pathway. In this MLRA, the most probable origin of this state is plant succession following cropland abandonment. This plant community will initially include a variety of annual forbs and grasses, some of which may be noxious weeds needing control. Over time, however, the exotic cool-season grasses (Kentucky bluegrass, smooth brome, crested wheatgrass, and/or quackgrass) will likely predominate.

### **Restoration R5A**

This Restoration Pathway from State 5: Go-Back State to the State 2: Native/Invaded State can be accomplished with a successful range planting. Following planting, prescribed grazing, prescribed burning, haying, or use of herbicides will generally be necessary to achieve the desired result and control any noxious weeds.

It may be possible using selected plant materials and agronomic practices to approach something very near the functioning of State 2: Native/Invaded State. Application of chemical herbicides and the use of mechanical planting methods using adapted varieties of the dominant native grasses are possible and can be successful. After establishment of the native plant species, prescribed grazing should include adequate recovery periods following each grazing event and stocking levels which match the available resources; management objectives must include the maintenance of those species, the associated reference state functions, and continued treatment of exotic grasses.

**Context dependence** (i.e., factors that cause variations in plant community shifts, restoration likelihood, and contribute to uncertainty). A successful range planting will include proper seedbed preparation, weed control (both prior to and after the planting), selection of adapted native species representing

functional/structural groups inherent to the State 1, and proper seeding technique. Management (e.g., prescribed grazing, prescribed burning) during and after establishment must be applied in a manner that maintains the competitive advantage for the seeded native species. Adding non-native species can impact the above and below ground biota. Elevated soil nitrogen levels have been shown to benefit smooth brome and Kentucky bluegrass more than some native grasses. As a result, fertilization, exotic legumes in the seeding mix, and other techniques that increase soil nitrogen may promote smooth brome and Kentucky bluegrass invasion.

The method or methods of herbaceous weed treatment will be site specific to each situation but, generally, the goal would be to apply the pesticide, mechanical control, or biological control (either singularly or in combination) in a manner that shifts the competitive advantage from the targeted species to the native grasses and forbs. The control method(s) should be as specific to the targeted species as possible to minimize impacts to non-target species.

### **Restoration R5B**

A failed range planting and/or secondary succession will lead to State 3: Invaded State.

**Context dependence** (i.e., factors that cause variations in plant community shifts, restoration likelihood, and contribute to uncertainty). Failed range plantings can result from many causes, both singularly and in combination, including drought, poor seedbed preparation, improper seeding methods, seeded species not adapted to the site, insufficient weed control, herbicide carryover, poor seed quality (purity & germination), improper management.

### **Transition T6A**

This is the Transition from any plant community to State 5: Go-Back State. It is most commonly associated with the cessation of cropping without the benefit of range planting, resulting in a “go-back” situation. Soil conditions can be quite variable on the site, in part due to variations in the management/cropping history (e.g., development of tillage induced compaction, erosion, fertility, and/or herbicide/pesticide carryover). Thus, soil conditions should be assessed when considering restoration techniques.

## **Ecological Site Interpretations**

### **Animal Community – Wildlife Interpretations**

#### Landscape

The MLRA 56A landscape is characterized by a nearly level glacial lake plain bordered on the east and west by outwash plains, till plains, gravelly beaches, and dunes. MLRA 56A is considered to have a continental climate with cold winters and hot summers, low humidity, light rainfall, and much sunshine. Extremes in temperature are common and characteristic of the MLRA. This area supports natural tall-grass prairie vegetation with bur oak, green ash, and willow growing in drainageways. This area is formed in silty and clayey lacustrine sediments from the former Glacial Lake Agassiz. Complex intermingled ecological sites create diverse grass/shrub land habitats interspersed with varying densities of linear, slope, depressional, and in-stream wetlands associated with headwater streams and tributaries to the Red River of the North. MLRA 56A is located within the boundaries of the Prairie Pothole Region and is an ecotone between the humid east and the sub-humid west regions. The primary land use is annual cropland (~80%). The Red River Valley is known for its exceptional fertility with major crops including corn, soybeans, small grains, and sugar beets.

By the mid-19<sup>th</sup> century, the majority of the Red River Valley had been converted from tall-grass prairie to annual crop production. To alleviate crop production loss from wetlands and overland flow, a system of shallow surface ditches, judicial ditches, and road ditches removes surface water in spring and during high rainfall events. The major soils are poorly drained with extensive areas of saline soils. Tile drainage systems have been or are being extensively installed throughout MLRA 56A for sub-surface field drainage to enhance annual crop production.

The east and west side of the Red River Valley formed in a complex pattern of sandy beach material, stratified inter-beach material, lacustrine silts, and lake washed glacial till. The soils vary from excessively drained on ridges to very poorly drained organic basins. Surface ditches serve to drain some of the area, although much of the area lacks adequate drainage for maximum crop production. Calcareous fens and saline seeps can occur at the base of beach ridges and result in rare plant communities. Native vegetation was mixed- and tall-grass prairie with scattered woodland and brush.

#### Historic Communities/Conditions within MLRA 56A:

The northern tall- and mixed-grass prairie was a disturbance-driven ecosystem with fire, herbivory, and climate functioning as the primary ecological drivers - either singly or often in combination. Frequent and expansive flooding along the Red River and its tributaries provided abundant opportunities for Native Americans to harvest wild rice. American bison roamed MLRA 56A wintering along the Red River and migrating west into MLRA 55A and 55B for parts of the season. Many species of grassland birds, small mammals, insects, reptiles, amphibians, and large herds of roaming American bison, elk, and pronghorn were historically among the inhabitants adapted to this region. Roaming herbivores, as well as several small mammal and insect species, were the primary consumers linking the grassland resources to large predators (such as the wolf and American black bear) and smaller carnivores (such as the coyote, bobcat, red fox, and raptors). Extirpated species include free-ranging American bison and gray wolf (breeding). Extinct from the region is the Rocky Mountain locust.

#### Present Communities/Conditions within MLRA 56A:

MLRA 56A has the most conversion to cropland of any MLRA within Region F-Northern Great Plains. European influence has impacted remaining grassland and shrubland by domestic livestock grazing, elimination of fire, removal of surface and subsurface hydrology via artificial drainage, and other anthropogenic factors influencing plant community composition and abundance.

Extensive drainage has taken place. Streams have been straightened, removing sinuosity, and riparian zones have been converted to annual crop production. These anthropogenic impacts have reduced flood water detention and retention on the landscape, increasing storm water runoff, sediment, and nutrient loading to the Red River and its tributaries. The installation of instream structures has reduced aquatic species movement within the MLRA.

Annual cropping is the main factor contributing to habitat fragmentation, reducing habitat quality for area-sensitive species. These influences fragmented the landscape, reduced, or eliminated ecological drivers (fire), and introduced exotic species including smooth brome, Kentucky bluegrass, and leafy spurge which further impacted plant and animal communities. The loss of the bison and fire as primary ecological drivers greatly influenced the character of the remaining native plant communities and the associated wildlife, moving towards a less diverse and more homogeneous landscape.

Included in this MLRA are approximately 70,000 acres of the United States Forest Service, Sheyenne National Grassland (southern portion of MLRA) with an additional 65,000 acres of intermingled privately owned land of sandy soils providing a large tract of intact tall grass prairie within the MLRA. United Fish and Wildlife Service refuges and waterfowl production areas, along with state wildlife management areas cover approximately 67,000 acres within the MLRA. Two of three largest cities in North Dakota are located within the MLRA.

USDA conservation programs have seeded thousands of cropland acres in riparian zones to native herbaceous vegetation. Natural succession is replacing the planted native herbaceous vegetation to native woody vegetation re-establishing native wooded riparian areas on previously cropland. Most of the plantings have been along the Red River and its tributaries in the northern portions of the MLRA within the United States. These areas are privately owned and protected from annual agricultural production with perpetual conservation easements.

Some characteristic wildlife species in this area are:

Birds: Mallard, blue-winged teal, red-tailed hawk, American kestrel, ring-necked pheasant, western meadowlark, killdeer, eastern and western kingbird, American crow, common yellowthroat, downy and hairy woodpecker, clay-colored sparrow, vesper sparrow, Savannah sparrow, and brown-headed cowbird.

Mammals: Northern short-tailed shrew, white-tailed jackrabbit, snowshoe hare, Franklin's ground squirrel, thirteen-lined ground squirrel, northern pocket gopher, plains pocket gopher, western harvest mouse, deer mouse, meadow vole, meadow jumping mouse, western jumping mouse, coyote, red fox, raccoon, American badger, striped skunk, white-tailed deer, North American beaver, and moose.

Reptiles/Amphibians: American toad, Great Plains toad, northern leopard frog, chorus frog, tree frog, tiger salamander, plains garter snake, and common garter snake.

Presence of wildlife species is often determined by ecological site characteristics including grass and forb species, hydrology, aspect, and other associated ecological sites. The home ranges of most species are usually larger than one ecological site or are dependent upon more than one ecological site for annual life requisites. Ecological sites offer different habitat elements as the annual life requisites change. Habitat improvement and creation must be conducted within the mobility limits of a known population for the species.

Insects play an important role providing ecological services for plant community development. Insects that are scavengers or aid in decomposition provide the food chain baseline sustaining the carnivorous insects feeding upon them. Many insects provide the ecological services necessary for pollination, keeping plant communities healthy and productive. Insects provide a protein food source for numerous species including grassland-nesting birds and their young. Extensive use of insecticides for specialty crops (such as potatoes, sugar beets, and other crops) has greatly reduced insects within this MLRA.

#### Species of Concern within MLRA 56A:

The following is a list of species considered "species of conservation priority" in the North Dakota State Wildlife Action Plan (2015); "species of greatest conservation need" in the Minnesota State Wildlife Action Plan, Conservation Focus Areas, Target Species (2015) and the South Dakota State

Wildlife Action Plan (2014); and “species listed as threatened, endangered, or petitioned” under the Endangered Species Act within MLRA 56A at the time this section was developed:

Invertebrates: Arogos skipper, Assiniboia skipper, Dakota skipper, dusted skipper, Leonard’s skipper, monarch butterfly, Poweshiek skippering, red-tailed leafhopper, regal fritillary, and Uhler’s Arctic.

Birds: American kestrel, American bittern, bobolink, American white pelican, bald eagle, black-billed cuckoo, chestnut-collared longspur, Dickcissel, grasshopper sparrow, greater prairie-chicken, Henslow’s sparrow, LeConte’s sparrow, loggerhead shrike, marbled godwit, Nelson’s sparrow, northern harrier, northern pintail, red-headed woodpecker, sharp-tailed grouse, short-eared owl, Swainson’s hawk, upland sandpiper, western meadowlark, willet, Wilson’s phalarope, and yellow rail.

Mammals: Arctic shrew, big brown bat, eastern spotted skunk, gray fox, little brown bat, northern grasshopper mouse, plains pocket mouse, prairie vole, pygmy shrew, Richardson’s ground squirrel, and river otter.

Amphibians/Reptiles: Canadian toad, common snapping turtle, northern prairie skink, and plains gognose snake.

Fish: Blacknose shiner, blue sucker, burbot, chestnut lamprey, finescale dace, hornyhead chub, largescale stoneroller, logperch, northern pearl dace, northern redbelly dace, pearl dace, shortnose gar, sickle-fin chub, silver chub, silver lamprey, trout-perch, and yellow bullhead.

Mussels: Black sandshell, creek heelsplitter, creeper, mapleleaf, pink heelsplitter, pink papershell, threeridge, and Wabash pigtoe.

### Grassland Management for Wildlife in MLRA 56A:

Management activities within the community phase pathways have both short and long term positive and negative impacts on wildlife. Community phase, transitional, and restoration pathways are keys to long-term management within each State and between States. Significant inputs must occur to cross the threshold between States (e.g., State 3.0 to 2.0) requiring substantial economic inputs and management (grazing intensity, reseeding, prescribed fire, woody vegetation removal, etc.). Timing, intensity, and frequency of these inputs can have dramatic positive or negative effects on vegetative structure impacting local wildlife species’ habitats. Ranchers and other land managers must always consider the long-term beneficial effects of management on the habitat in comparison to potential short-term negative effects to individual species.

Ecological sites occur as intermingled complexes on the landscape with gradual or sometimes abrupt transitions. Rarely do ecological sites exist in large enough acreage to manage independently. Ecological sites supporting a dominance of herbaceous vegetation (Wet Meadow, Subirrigated Sands) can be located adjacent to ecological sites that support trees (Choppy Sands and Loamy Overflow).

Management of these complex ecological sites can provide a heterogeneous or a homogenous landscape. Grassland bird use reduces as the plant community transitions to a homogenous state or increases in woody vegetation. Managers need to recognize ecological sites and the complexes they occur in to properly manage the landscape. A management regime for one ecological site may negatively impact an adjacent site (e.g., alteration of a grazing regime within a Choppy Sands

ecological site to encourage understory growth may encourage exotic cool-season grasses to increase or dominate an adjacent ecological site).

Life requisites and habitat deficiencies are determined for targeted species. Deficiencies need to be addressed along community phase, transitional, and restoration pathways as presented in specific state-and-transition models. Ecological sites should be managed and restored within the site's capabilities to provide sustainable habitat. Managers also must consider habitat provided by adjacent/intermingled ecological sites for species with home ranges or life requisites that cannot be provided by one ecological site.

With populations of many grassland-nesting birds in decline, it is important to maintain these ecological sites in a 1.0 Reference State or the 2.0 Native/Invaded State. Plant communities optimal for a guild of grassland species serve as a population source where the birth rate exceeds mortality. Species may use marginal plant communities; however, these sites may function as a population sink where mortality exceeds the birth rate.

Understanding preferred vegetative stature and sensitivity to woody encroachment is necessary to manage for the specific grassland species. Various grass heights may be used for breeding, nesting, or foraging habitat. While most species use varying heights, many have a preferred vegetative stature height. The following chart provides preferred vegetative stature heights and sensitivity to woody vegetation encroachment.

Grassland-nesting Bird Species	Preferred Vegetative Stature			Avoids woody vegetation*
	Short < 6 inches	Medium 6 - 12 inches	Tall >12 inches	
Baird's sparrow	x	x		x
Bobolink		x	x	x
Brewer's sparrow	x	x		
Burrowing owl	x			x
Chestnut-collared longspur	x	x		x
Common yellowthroat			x	
Dickcissel		x	x	
Ferruginous hawk	x	x		
Grasshopper sparrow	x	x		x
Horned lark	x			x
Killdeer	x			x
Lark bunting	x	x		
Lark sparrow	x			
Le Conte's sparrow			x	x
Long-bill curlew	x			x
Marbled godwit	x	x		x
McCown's longspur	x	x		x
Mountain plover	x			x
Nelson's sparrow			x	x
Nesting waterfowl		x	x	

Northern harrier		x	x	x
Savannah sparrow		x	x	x
Short-eared owl		x	x	x
Sprague's pipit	x	x		x
Upland sandpiper	x	x		x
Western meadowlark	x	x		
Willet	x	x		x
*Many of the listed species avoid nesting in grassland areas with large amounts of woody vegetation within a grassland or avoid nesting near woody vegetation in adjacent habitats. Although these species avoid areas with woody vegetation, most can tolerate a small amount of woody vegetation within areas dominated by grassland habitat, including short-statured shrubs (e.g., western snowberry) in this MLRA.				

### Sands Wildlife Habitat Interpretation:

Sands ecological sites are very deep, well to excessively drained, coarse textured (non-gravelly) soils usually found on sandy lake plains and deltas, as well as on beaches and beach ridges. These soils are highly susceptible to water and wind erosion. No significant water table or surface run-on influences vegetation production on this site. Associated ecological sites include Limy Subirrigated, Sandy, Subirrigated, Subirrigated Sands, and Choppy Sands. This complex of ecological sites provides habitat for many edge-sensitive grassland bird species which prefer medium- to tall-statured vegetation.

Insects rely on associated forbs and grasses for survival and serve as food sources for birds and their young, and as forage for small and large herbivores. Monarch butterfly and Regal fritillary use Sands ecological sites in MLRA 56A.

Sands ecological sites may be found in five plant community states (1.0 Reference State, 2.0 Native/Invaded State, 3.0 Invaded State, 4.0 Wooded State, and 5.0 Go-Back State) within a local landscape. Multiple plant community phases exist within States 1.0, and 2.0. Today, these states occur primarily in response to drought, fire, grazing, non-use, and other anthropogenic disturbances.

Because there is no known restoration pathway from State 2.0 to State 1.0, it is important to intensively manage using tools in State 1.0 and State 2.0 Community Phase Pathways to prevent further plant community degradation along the T2A Transitional Pathways to 4.0 Wooded State or T2B Transitional Pathway to 3.0 Invaded State. Native wildlife generally benefits from the heterogeneous grasslands found in States 1.0 and 2.0 that includes diverse grass and forb species of varying stature and density. As plant communities degrade within State 2.0, Kentucky bluegrass increases while native forbs are reduced. When Kentucky bluegrass exceeds 30%, the site transitions to 3.0 Invaded State. This transition results in reduced stature and increased plant community homogeneity. When adjacent and/or intermingled ecological sites undergo the same transition, the result can be an expansive, homogenous landscape.

Success along Restoration Pathway R3A from State 3.0 to State 2.0 is very difficult and is dependent upon presence of a remnant native grass population. This concept also applies to wildlife, as the target species must either be present on adjacent State 1.0 or State 2.0 plant communities or other ecological sites within the mobility limits of the species. Species with

limited mobility, such as Dakota skippers, must exist near the plant community to utilize restored sites. Mobile species (such as grassland-nesting birds) can easily locate isolated, restored plant communities.

Plant Community Phase 3.1 shows dramatic increased homogeneity of exotic cool-season grasses and further reduction in native forbs. Reduced forb diversity limits insect populations, negatively affecting foraging opportunities for grassland-nesting birds. Increased exotic grass litter can limit access to bare ground by nesting insects and can limit mobility by small chicks. A homogenous grassland landscape does not provide quality escape or winter cover. As a result, many species are not able to meet life requisites, especially non-migrating bird species.

Management along community phase, transition, or restoration pathways should focus upon attainable changes. Short- and long-term monetary costs must be evaluated against short- and long-term ecological services in creating and maintaining habitat of sufficient quality to support a sustainable population density.

## 1.0 Reference State

Community Phase 1.1 Prairie Sandreed-Porcupinegrass-Bluestems: This plant community offers quality vegetative cover for wildlife; every effort should be made to maintain this ecological site within this community phase. This phase retains high functionality through continued grassland management including prescribed grazing with adequate recovery period, as well as prescribed fire. Predominance of grass species in this community favors grazers and mixed-feeders (animals selecting grasses as well as forbs and shrubs). The structural diversity provides habitat for a wide array of migratory and resident birds.

**Invertebrates:** Insects play a role in maintaining the forb community and provide a forage base for grassland birds, reptiles, and rodents. Ecological services, historically provided by bison, are simulated by domestic livestock. These services include putting plant material and dung in contact with mineral soil to be used by lower trophic level consumers (such as invertebrate decomposers, scavengers, shredders, predators, herbivores, dung beetles, and fungal-feeders).

**Arogos skippers** may use these sites since it contains host plants, such as big bluestem and little bluestem, and can use nectar from various forbs including thistles. **Violet** species are limited but, when available, **regal fritillary** can make use of this site. **Monarch butterfly** may use flowering forbs on this site; however, few milkweed species are found on this site to support caterpillar food. **Bumblebees** and other native bees utilize forbs as a nectar source; however, bare ground and nesting sites are limited due to the dominance of sod-forming grasses.

**Birds:** This plant community provides quality nesting, foraging, and escape habitats favored by mid- to tallgrass-nesting birds. Prescribed fire maintains a grass-dominated plant community providing habitat for bird species sensitive to woody vegetation. In years with reduced precipitation or drought, nesting recruitment may be compromised. This plant community provides suitable lek sites for sharp-tailed grouse or greater prairie chicken but do not provide nesting, brood-rearing, and escape habitat. This site provides good hunting opportunities for grassland raptors.

**Mammals:** The diversity of grasses and forbs provide high nutrition levels for small and large herbivores including voles, mice, jackrabbits, white-tailed deer, and elk. Tall- to mid-statured vegetation provides suitable food, thermal, protective, and escape cover for small herbivores.

**Amphibians and Reptiles:** This ecological site is not typically found adjacent to Wet Meadow ecological sites. Habitat for the northern leopard frog and Canadian toad is dependent on distance to the Wet Meadow ecological site. This site provides habitat for the northern prairie skink and plains hog-nosed snake.

**Fish and Mussels:** This ecological site is not typically adjacent to streams, rivers, or water bodies. This site receives limited run-on hydrology from adjacent ecological sites (Choppy Sands) and provides hydrology to Limy Subirrigated, Subirrigated Sands, and Subirrigated ecological sites. Management on Sands sites, in conjunction with neighboring run-on sites, will have an indirect effect on aquatic species in streams and/or tributaries receiving water from Sands and adjacent sites. Optimum hydrological function and nutrient cycling limit potential for sediment yield and nutrient loading to the adjacent aquatic ecosystems from Community Phase 1.1.

**Community Phase 1.2 Needle and Thread-Prairie Sandreed-Blue Grama:** Short- to mid-statured grasses dominate this plant community phase because of repeated heavy grazing via Community Phase Pathway 1.1A. Facilitated by periods of below average precipitation, the tall warm-season grass component has decreased with increases in green sagewort, goldenrod, Cuman ragweed, and common yarrow. The forb component provides pollinator habitat and a food source for grassland-nesting birds.

**Invertebrates:** Provides similar life requisites as Community Phase 1.1. However, an increase in bare ground may provide additional habitat for ground nesting species.

**Birds:** Provides similar life requisites as Community Phase 1.1. However, an increase in short- to mid-grass species favors grassland-nesting birds that prefer short- to mid-statured vegetation.

**Mammals:** Provides similar life requisites as Community Phase 1.1, however cover for large ungulates maybe somewhat reduced.

**Amphibians and Reptiles:** Provides similar life requisites as Community Phase 1.1.

**Fish and Mussels:** Provides similar life requisites as Community Phase 1.1.

## **2.0 Native Invaded State**

**Community Phase 2.1 Prairie Sandreed-Porcupinegrass-Bluestems:** This plant community develops through Transition Pathway T1A due to changes in management (reduction in fire frequency) and the presence of exotic, cool-season grasses. The threshold between States 1.0 and 2.0 is crossed when Kentucky bluegrass or smooth brome become established. This plant community phase has a very similar appearance and function to the Plant Community 1.1, except that it has minor amounts of cool-season exotic grasses and forbs with declines in cool- and warm-season bunch grasses. This phase functions at a high level for native wildlife; therefore, managers should consider management within the State 2.0 Community Phase Pathways to avoid transitioning to State 3.0.

**Invertebrates:** Provides similar life requisites as Community Phase 1.1.

**Birds:** Provides similar life requisites as Community Phase 1.1.

**Mammals:** Provides similar life requisites as Community Phase 1.1.

**Amphibians and Reptiles:** Provides similar life requisites as Community Phase 1.1.

Fish and Mussels: Provides similar life requisites as Community Phase 1.1.

Community Phase 2.2 Needle and Thread-Prairie Sandreed-Blue Grama/Sun Sedge/Forbs: Heavy grazing increases grazing-tolerant needle and thread while decreasing porcupinegrass. Short, sod/turf forming grasses and sedges begin to dominate the site. The forb component is still diverse; however, leguminous native forbs are decreasing while less desirable, native pollinator forb species (such as common yarrow, tarragon, and white sagebrush) are increasing. Every effort should be made by managers to avoid implementing a grazing system that may favor Kentucky bluegrass (via Community Phase Pathway 2.2B or Transitional Pathway T2B). Continuous, heavy season-long grazing or complete rest could cause this plant community to reach the 30 percent threshold of Kentucky bluegrass, crossing over to State 3.0. Once in State 3.0, prescribed grazing with prescribed burning may help restore this plant community to State 2.0.

Invertebrates: Provides similar life requisites as Community Phase 1.1; however, the forb community is shifting toward less desirable species, such as common yarrow and white sagebrush, which are less desirable for native bees.

Birds: Provides similar life requisites as Community Phase 1.2.

Mammals: Provides similar life requisites as Community Phase 1.2.

Amphibians and Reptiles: Provides similar life requisites as Community Phase 1.1.

Fish and Mussels: Provides similar life requisites as Community Phase 1.1.

Community Phase 2.3 Blue Grama/Sun Sedge/Exotic Cool-Season Grasses/Forbs: Continuous heavy grazing without adequate recovery periods (along Community Phase Pathway 2.2B) allows short-statured grasses (blue grama) and sedges to be more competitive and dominate this community phase. Grazing tolerant forbs (such as goldenrods, tarragon, yellow salsify, and white heath aster) increase, thus reducing pollen and nectar sources. This is an “at risk” plant community with high amounts of Kentucky bluegrass out-competing native grasses. Every effort should be made by managers to avoid implementing a grazing system that may favor reaching the 30 percent Kentucky bluegrass threshold via Transitional Pathways T2B. When this plant community transitions over to State 3.0. Once in State 3.0, prescribed grazing with prescribed burning may help restore this plant community to State 2.0.

Invertebrates: Reduced litter amounts with slight increase in bare ground will favor ground nesting insects. Abundant forbs still provide for season-long pollen and nectar source for insects and increase bare ground for ground-nesting insects.

Birds: Grassland-nesting birds that favor short-statured vegetative cover will use this site.

Mammals: This site does not provide thermal or escape cover but provides limited forage or browse for large herbivores. Due to heavy continuous grazing and/or heavy continuous season-long grazing reducing litter depth and residual vegetation, thermal, protective, escape, and winter cover is limited for small mammals.

Amphibians and Reptiles: Provides similar life requisites as Community Phase 1.1.

Fish and Mussels: Provides similar life requisites as Community Phase 1.1.

### **3.0 Invaded State**

Community Phase 3.1 Exotic Cool-Season Grasses/Forbs: Heavy grazing without adequate recovery periods, extended periods of non-use, or cropland allowed to “go-back” can cause Community Phase 3.1 to establish. Kentucky bluegrass will dominate the site. Potential use by wildlife will be dependent upon management. Non-use will result on extensive areas of Kentucky bluegrass litter accumulation and have the effect of residual cover flopping. Heavy grazing without adequate recovery periods will result in little, if any, residual cover.

Community Phase 3.1 can also result via Restoration Pathway R4b from Community Phase 4.1 when shrubs are removed, resulting in a Kentucky bluegrass dominated plant community. Management following shrub removal will determine the quality of wildlife habitat. Extensive management using prescribed grazing, herbicide treatments, range seedings, and prescribed fire will be needed to move this plant community to State 2.0 (via Restoration Pathway R3A).

**Invertebrates:** Litter amounts will vary in this community phase. Non-native forbs dominate with goldenrods and Cuman ragweed also present. A reliable amount of season-long pollen and nectar sources will not be present. Intense grazing will reduce litter amounts and provide some bare ground for ground nesting insects. Non-use will increase litter, reduce sites for ground nesting pollinators, and limit pollen and nectar availability. Overall, this community phase will not favor pollinating species in either management scenario (heavy use or non-use). This plant community will not provide life requisites for Dakota skippers or Regal fritillary, but it may support monarch butterfly's depending on availability of milkweed.

**Birds:** Heavy grazing will favor short-grass bird species. Short- to mid-grass bird species will also be favored with the non-use management scenario. Lack of pollinator forb species limits invertebrate use, reducing protein sources for food. This plant community does not provide suitable areas for lek sites or for brood or escape cover for sharp-tailed grouse or greater prairie chicken. This site provides hunting opportunities for grassland raptors.

**Mammals:** Lacking residual cover under the over-grazing management scenario, this short-statured community provides limited food and thermal, protective, and escape cover for small herbivores. Under the non-use management scenario, exotic cool-season grasses will provide residual cover for small mammals (such as mice, ground squirrels, and voles). Use by large mammals will be limited under either management scenario.

**Amphibians and Reptiles:** Kentucky bluegrass litter accumulations may become too heavy for northern prairie skink and plains hog-nose snake under the non-use scenario. Under heavy grazing, cover for amphibians and reptiles can be eliminated.

**Fish and Mussels:** When grazed heavily, runoff increases and is the highest of any plant community phase on this ecological site, increasing run-off to adjacent ecological sites and into neighboring water bodies. Nutrient cycling is severely limited to the rooting depth of the Kentucky bluegrass, increasing nutrient loading to neighboring water bodies.

### **4.0 Invaded Wooded State**

Community Phase 4.1 Shrub/Exotic Cool-Season Grasses/Forbs: Long-term non-use or very light grazing and no fire, enhanced by above average precipitation (via Transitional Pathways T2A) causes this plant community to be dominated by shrubs. Plant litter accumulation favors shade-tolerant grasses and western snowberry. Forb number and diversity is low with western

snowberry and other flowering shrubs beginning to dominate or dominating the site. Pathway R4A will require intensive management practices and extensive time to bring forb and grass diversity back to State 2.0.

**Invertebrates:** The invasion of western snowberry reduces or eliminates habitats for all pollinating species of concern within MLRA 56A. Season-long pollen and nectar availability becomes limited on this site. The woody shrub component (western snowberry) provides an early season bloom period. Non-use will increase litter, reducing sites for ground nesting pollinators. Overall, pollinator plant diversity is low, limiting season-long nectar and pollen production. Shrubs, mainly western snowberry, provide early-season pollen sources. Season-long pollen and nectar is typically not available. This site no longer provides life requisites for butterfly and bee species. Overall, this community phase will not favor pollinating species in either management scenario (heavy use or no use).

**Birds:** This site may still provide habitat needs for many grassland nesting birds, dependent upon the density and height the shrubs. Many grassland nesting birds are tolerant or positively correlated with western snowberry. Northern harrier, lark bunting, and Dickcissel use western snowberry for nesting sites. This plant community does not provide suitable areas for sharp-tailed grouse or greater prairie chicken lek sites but does provide brood rearing and escape habitat. Winter cover is mainly provided by the western snowberry. This site provides hunting opportunities for grassland raptors. Management along Restoration Pathway R3A will require intensive mechanical and management practices and extensive time to bring forb and grass diversity back to this plant community phase.

**Mammals:** Western snowberry will provide increased year-round cover and browse for white-tailed deer. The short stature and density of western snowberry will not provide habitat for porcupines or other mammals that prefer woody habitat.

**Amphibians and Reptiles:** The increase in western snowberry will not provide life requisites for amphibians or reptiles.

**Fish and Mussels:** Provides similar life requisites as Community Phase 1.1.

## 5.0 Go-Back State

**Community Phase 5.1 Annual/Pioneer Perennial/Exotics:** These plant communities are the result of severe soil disturbance (such as cropping, recreational activity, or concentrated livestock activity for a prolonged time). Following cessation of disturbances, the resulting plant community is dominated by early pioneer annual and perennial plant species. Plant species composition and production are highly variable. Weedy plants can provide pollinator habitat along with spring and summer cover for many mammals and birds, and their young. Dense weed cover can keep soils moist, increasing the presence of insects. Milkweed can be an early pioneering pollinator species and host plant for monarch butterflies. Tall stature provided by some annual weeds offers thermal cover and seeds throughout winter for deer, small mammals, and over-wintering birds. The response by wildlife species will be dependent upon plant community composition, vegetative stature, patch size, and management activities (such as prescribed grazing, burning, inter-seeding, haying, or noxious weed control).

Successful restoration of native species along Transition Pathway R5A can result in a native grass and forb community in Native/Invaded State 2.0. Over time (with no management), the exotic cool-season perennial grasses (Kentucky bluegrass, smooth brome, and/or quackgrass) generally become

established and dominate the community. Failed range planting, via Transition Pathway R5B, can result in an invaded plant community (Invaded State 3.0).

## **Animal Community – Grazing Interpretations**

This site is well adapted to managed grazing by domestic livestock. The predominance of herbaceous plants across all plant community phases best lends these sites to grazing by cattle, but other domestic grazers with differing diet preferences may also be a consideration depending upon management objectives. Often, the current plant community does not entirely match any particular plant community (as described in the ecological site description). Because of this, a resource inventory is necessary to document plant composition and production. Proper interpretation of this inventory data will permit the establishment of a safe, initial stocking rate for the type and class of animals and level of grazing management. More accurate stocking rate estimates should eventually be calculated using actual stocking rate information and monitoring data.

NRCS defines prescribed grazing as “managing the harvest of vegetation with grazing and/or browsing animals with the intent to achieve specific ecological, economic, and management objectives”. As used in this site description, the term ‘prescribed grazing’ is intended to include multiple grazing management systems (e.g., rotational grazing, twice-over grazing, conservation grazing, targeted grazing, etc.) provided that, whatever management system is implemented, it meets the intent of prescribed grazing definition.

The basic grazing prescription addresses balancing forage demand (quality and quantity) with available forage, varying grazing and deferment periods from year-to-year, matching recovery/deferment periods to growing conditions when pastures are grazed more than once in a growing season, implementation of a contingency (e.g., drought) plan, and a monitoring plan. When the management goal is to facilitate change from one plant community phase or state to another, then the prescription needs to be designed to shift the competitive advantage to favor the native grass and forb species.

Grazing levels are noted within the plant community narratives and pathways in reference to grazing prescribed grazing management. “Degree of utilization” is defined as the proportion of the current year’s production that is consumed and/or destroyed by grazing animals (may refer to a single plant species or a portion or all the vegetation). “Grazing utilization” is classified as slight, moderate, full, close, and severe (see the following table for description of each grazing use category). The following utilization levels are also described in the Ranchers Guide to Grassland Management IV. Utilization levels are determined by using the landscape appearance method as outlined in the Interagency Technical Reference “Utilization Studies and Residual Measurements” 1734-3.

Utilization Level	%	Use Description
Slight (Light)	0-20	Appears practically undisturbed when viewed obliquely. Only choice areas and forage utilized.
Moderate	20-40	Almost all of accessible range shows grazing. Little or no use of poor forage. Little evidence of trailing to grazing.
Full	40-60	All fully accessible areas are grazed. The major sites have key forage species properly utilized (about half taken, half left). Points of concentration with overuse limited to 5 to 10 percent of accessible area.
Close (Heavy)	60-80	All accessible range plainly shows use and major sections closely cropped. Livestock forced to use less desirable forage, considering seasonal preference.
Severe	> 80	Key forage species completely used. Low-value forages are dominant.

## Hydrology Functions

Available water is the principal factor limiting herbage production on this site. The site is dominated by soils in hydrologic group A, but some soils in group B are included. Infiltration varies from moderately rapid to rapid; runoff potential varies from negligible to medium for this site depending on soil hydrologic group, surface texture, slope percent, and ground cover. In many cases, areas with greater than 75% ground cover have the greatest potential for high infiltration and lower runoff. An exception would be where short-grasses form a dense sod and dominate the site. Areas where ground cover is less than 50% have the greatest potential to have reduced infiltration and higher runoff (refer to Section 4, NRCS National Engineering Handbook for runoff quantities and hydrologic curves).

## Recreational Uses

Hunting and Bird Watching: The United States Fish and Wildlife Service manages approximately 4,000 acres of National Wildlife Refuges for hiking and bird watching and approximately 24,000 acres of Waterfowl Production Areas for public hunting, hiking, and bird watching. States within MLRA 56A manage approximately 39,000 acres of wildlife management areas for multiple use including hunting, fishing, hiking, birdwatching, berry picking, and other non-motorized uses. Of the 39,000 acres, approximately 21,400 are in Minnesota with approximately 16,000 acres in North Dakota and approximately 1,700 acres in South Dakota.

In North Dakota, the United States Forest Service manages 70,000 acres on the Sheyenne National Grassland for multiple uses including camping, hunting, photography, backpacking, birdwatching, biking, horseback riding, and other non-motorized recreation. The Sheyenne National Grassland is also managed for livestock grazing. The Choppy Sands and Sands ecological sites dominate the Grassland. It

is the only National Grassland in the tallgrass prairie region of the United States. The grassland provides habitat for greater prairie chickens as well as several other sensitive species, such as the Dakota skipper and regal fritillary. It also contains one of largest populations of the western prairie fringed orchid which is listed as a threatened species by the U.S. Fish and Wildlife Service.

Fishing: Approximately 20 lakes are managed for public fishing MLRA 56A. Most of these lakes offer boat docks and ramps. These lakes contain various sport fish including walleye, northern pike, yellow perch, crappie, and bluegill. The Red River runs from south to north through the center of the MLRA. The Red River is best known for channel catfish but also has walleye, sauger, northern pike, and smallmouth bass. The Red River is 550 miles long from its source in the southern end of the MLRA near Breckenridge, Minnesota to Lake Winnipeg in Manitoba, Canada. Between North Dakota and Minnesota, there are 32 public access points along the Red River with 18 having boat ramps.

Camping: Four state parks or recreation areas provide of modern and primitive camping facilities. Minnesota hosts the Buffalo River State Park and Red River State Park. North Dakota hosts the Icelandic State Park and Turtle River State Park. These Parks provide hiking, biking, birding, canoeing, and wildlife viewing opportunities. Many local parks and private parks provide modern and primitive camping opportunities. Limited primitive camping is also available on North Dakota Game and Fish Department Wildlife Management Areas.

Hiking/Biking/Horseback Riding: Hiking is permitted on most state and federally owned lands. Developed hiking and biking trails can be found the four state parks. The Grand Forks Greenway has over 22 miles of trials while municipalities along the Red River have extensive walking and hiking trails. A 30-mile segment of the North Country National Scenic Trail leads hikers through the Sheyenne National Grassland's unique landforms and plant communities. This trail has three trailheads along its route; it is a graveled, marked trail. The entire North Country National Scenic Trail stretches from Crown Point, New York to Lake Sakakawea near Garrison, North Dakota.

Canoeing/Kayaking: The Red River has six designated canoe/kayaking trials. Public access, with limited rentals, is available at these segments. Sheyenne River Water Trail has a segment within the MLRA Sheyenne National Grasslands. Canoe/kayak rentals are available at Icelandic State Park.

## Wood Products

No appreciable wood products are present on the site.

## Other Products

Seed harvest of native plant species can provide additional income on this site.

## Site Development and Testing Plan

- Review of water table depths for Hecla and Radium components is needed. Currently, it is populated in NASIS with a range of 76 to 152 cm and an RV of 114 cm. This range overlaps ESD criteria for Subirrigated Sands and Sands. MLRAs 55A and 55B have linked Hecla to Subirrigated Sands while MLRA 56A has linked Hecla to Sands. It is not known if the populated water table depths reflect natural hydrology or hydrology affected by drainage. If affected by drainage, components should be relinked to Subirrigated Sands.
- Further investigation may be needed on beach and beach ridge soils which have higher amounts of gravel-sized rock fragments than typical for this ESD. In particular, Sandberg soils can overlap the concepts of Sands and Shallow Gravel ESDs due to the variability of depth to gravel and the amount of gravel within a depth of 20 inches. MLRA map units needing investigation are:
  - Radium-Sandberg-Garborg complex, 0 to 2 percent slopes (map unit vx9x)

- Sandberg loamy sand, 15 to 25 percent slopes (map unit 2sw8y)
- Sandberg loamy sand, 6 to 15 percent slopes (map unit 2sw8z)
- Sandberg-Radium complex, 0 to 6 percent slopes (map unit prpd)
- In Pembina County, some areas of forested Maddock and Corliss soils with slopes <15% occur; these appear to be long-term forest conditions rather than an Invaded Wooded State. Field investigation may be needed.
- Further evaluation and refinement of the State-and-Transition model is needed to identify disturbance driven dynamics. Additional states and/or phases may be required to address grazing response.
- Further documentation may be needed for plant communities in all states. Plant data has been collected in previous range-site investigations, including clipping data; however, this data needs review. If geo-referenced sites meeting Tier 3 standards for either vegetative or soil data are not available, representative sites will be selected for further investigation.
- Site concepts will be refined as the above noted investigations are completed.
- The long-term goal is to complete an approved, correlated Ecological Site Description as defined by the National Ecological Site Handbook.
- NASIS revisions needed:
  - One minor component of Banks should be relinked from Sandy to Sands.

This ESD is the best available knowledge. The site concept and species composition table have been used in the field and tested for more than five years. It is expected that as additional information becomes available revisions may be required.

## Supporting Information

### Associated Sites

Ecological Site Name	Site ID	Narrative
Limy Subirrigated	R056AY087ND	This site is lower on the landscape. The soils are highly calcareous in the upper part of the subsoil (within a depth of 16 inches) and have redoximorphic features at a depth of 18 to 30 inches. All textures are included in the site.
Sandy	R056AY091ND	This site occurs on similar landscape positions. It has fine sandy loam or sandy loam textures (forms a ribbon <1 inch long) to a depth >10 inches.
Subirrigated	R056AY095ND	This site occurs in swales and blow-outs. It is non-calcareous to a depth >16 inches and has redoximorphic features at a depth of 18 to 30 inches.
Subirrigated Sands	R056AY096ND	This site occurs lower on the landscape. It has redoximorphic features at a depth of 30 to 40 inches. The subsoil does not form a ribbon.
Choppy Sands	R056AY104ND	This site occurs on dunes and ridges with slopes >15 percent. The surface and subsoil layers do not form a ribbon.

### Similar Sites

Ecological Site Name	Site ID	Narrative

Sandy	R056AY091ND	This site occurs on similar landscape positions. It has fine sandy loam or sandy loam textures (forms a ribbon <1 inch long) to a depth >10 inches.
Subirrigated Sands	R056AY096ND	This site occurs lower on the landscape. It has redoximorphic features at a depth of 30 to 40 inches. The subsoil does not form a ribbon.
Choppy Sands	R056AY104ND	This site occurs on dunes and ridges with slopes >15 percent. The surface and subsoil layers do not form a ribbon.

## Supporting Information

### Associated Sites

Ecological Site Name	Site ID	Narrative
Limy Subirrigated	R056AY087ND	This site is lower on the landscape. The soils are highly calcareous in the upper part of the subsoil (within a depth of 16 inches) and have redoximorphic features at a depth of 18 to 30 inches. All textures are included in the site.
Sandy	R056AY091ND	This site occurs on similar landscape positions. It has fine sandy loam or sandy loam textures (forms a ribbon <1 inch long) to a depth >10 inches.
Subirrigated	R056AY095ND	This site occurs in swales and blow-outs. It is non-calcareous to a depth >16 inches and has redoximorphic features at a depth of 18 to 30 inches.
Subirrigated Sands	R056AY096ND	This site occurs lower on the landscape. It has redoximorphic features at a depth of 30 to 40 inches. The subsoil does not form a ribbon.
Choppy Sands	R056AY104ND	This site occurs on dunes and ridges with slopes >15 percent. The surface and subsoil layers do not form a ribbon.

### Similar Sites

Ecological Site Name	Site ID	Narrative
Sandy	R056AY091ND	This site occurs on similar landscape positions. It has fine sandy loam or sandy loam textures (forms a ribbon <1 inch long) to a depth >10 inches.
Subirrigated Sands	R056AY096ND	This site occurs lower on the landscape. It has redoximorphic features at a depth of 30 to 40 inches. The subsoil does not form a ribbon.
Choppy Sands	R056AY104ND	This site occurs on dunes and ridges with slopes >15 percent. The surface and subsoil layers do not form a ribbon.

### Acknowledgements

We gratefully acknowledge Bernadette Braun, Dr. Mark Gonzales, Jamie Kienzle, Stacy Swenson, USFS; and Rob Self, The Nature Conservancy; for their helpful comments and suggestions.

### Developers

ND NRCS: Keith Anderson, Fred Aziz, Stan Boltz, David Dewald, Jonathan Fettig, Alan Gulsvig, Mark Hayek, Chuck Lura, Jeff Printz, Steve Sieler, Lee Voigt, and Hal Weiser.

**Non-discrimination Statement:** In accordance with Federal civil rights law and U.S. Department of Agriculture (USDA) civil rights regulations and policies, the USDA, its Agencies, offices, and employees,

and institutions participating in or administering USDA programs are prohibited from discriminating based on race, color, national origin, religion, sex, gender identity (including gender expression), sexual orientation, disability, age, marital status, family/parental status, income derived from a public assistance program, political beliefs, or reprisal or retaliation for prior civil rights activity, in any program or activity conducted or funded by USDA (not all bases apply to all programs). Remedies and complaint filing deadlines vary by program or incident. Persons with disabilities who require alternative means of communication for program information (e.g., Braille, large print, audiotape, American Sign Language, etc.) should contact the responsible Agency or USDA's TARGET Center at (202) 720-2600 (voice and TTY) or contact USDA through the Federal Relay Service at (800) 877-8339. Additionally, program information may be made available in languages other than English. To file a program discrimination complaint, complete the USDA Program Discrimination Complaint Form, AD-3027, found online at How to File a Program Discrimination Complaint and at any USDA office or write a letter addressed to USDA and provide in the letter all of the information requested in the form. To request a copy of the complaint form, call (866) 632-9992. Submit your completed form or letter to USDA by:(1) mail: U.S. Department of Agriculture Office of the Assistant Secretary for Civil Rights 1400 Independence Avenue, SW Washington, D.C. 20250-9410;(2) fax: (202) 690-7442; or(3) email: [program.intake@usda.gov](mailto:program.intake@usda.gov)USDA is an equal opportunity provider, employer, and lender.

## **Inventory Data References**

Information presented here has been derived from NRCS and other federal/state agency clipping and inventory data. Also, field knowledge of range-trained personnel was used. All descriptions were peer reviewed and/or field-tested by various private, state and federal agency specialists.

## **State Correlation**

This site has been correlated with Minnesota, North Dakota, and South Dakota.

## **Relationship to Other Established Classifications**

Level IV Ecoregions of the Conterminous United States: 48a Glacial Lake Agassiz Basin; 48b Beach Ridges and Sand Deltas; and 48d Lake Agassiz Plains.

## **Other References**

Bakker, K.K. 2003. The effect of woody vegetation on grassland nesting birds: an annotated bibliography. The Proceedings of the South Dakota Academy of Science 82:119-141.

Barker, W.T. and W. C. Whitman. 1988. Vegetation of the northern great plains. Rangelands 10(6): 266-272.

Bluemle, J.P. 2016. North Dakota's Geologic Legacy. North Dakota State University Press. 382 pages.

Briske, D.D. (editor). 2017. Rangeland Systems – Processes, Management, and Challenges. Springer Series on Environmental Management. 661 pages.

Burgess, R.L. 1965. A study of plant succession in the sandhills of southeastern North Dakota. Proceedings ND Academy of Science 19:62-80

DeKeyser, E.S., G. Clambey, K. Krabbenhoft, and J. Ostendorf. 2009. Are changes in species composition on central North Dakota rangelands due to non-use management? Rangelands 31:16-19

Dix, R.L. and F.E. Smeins. 1967. The prairie, meadow, and marsh vegetation of Nelson County, North Dakota. Canadian Journal of Botany 45:21-57.

Dornbusch, M.J., R.F. Limb, and C.K. Gasch. 2018. Facilitation of an exotic grass through nitrogen enrichment by an exotic legume. Rangeland Ecology & Management 71:691-694.

Dyke, S.R., S.K. Johnson, and P.T. Isakson. 2015. North Dakota State Wildlife Action Plan. North Dakota Game and Fish Department, Bismarck, ND. 468 pages.

Ehrenfeld, Joan G. 2002. Effects of exotic plant invasions on soil nutrient cycling processes. *Ecosystems* 6:503-523.

Endangered and threatened wildlife and plants; designation of critical habitat for the Dakota skipper and Poweshiek skipperling; Vol. 79 No. Final Rule October 1, 2015, 50 CFR Part 17.

Erath, C., J. Hendrickson, D. Kirby, E. DeKeyser, K. Sedevec, and M. West. Controlling Kentucky bluegrass with herbicide and burning is influenced by invasion level. *Invasive Plant Science and Management* 10: 80-89.

Ewing, J. 1924. Plant Succession on the Brush Prairie in Northwestern Minnesota. *Journal of Ecology* 12:228-266.

Gilgert, W. and S. Zack. 2010. Integrating multiple ecosystem services introduction ecological site descriptions. *Rangelands*: 32:49-54.

Grant, T.A. and R.K. Murphy. 2005. Changes on woodland cover on prairie refuges in North Dakota, USA. *Natural Areas Journal* 25:359-368.

Heitschmidt, R. K., K. D. Klement, and M. R. Haferkamp. 2005. Interactive effects of drought and grazing on northern great plains rangelands. *Rangeland Ecology and Management* 58:11-19.

Hendrickson, J.R., P. S. Johnson, M. A. Liebig, K. K. Sedivec, and G. A. Halvorson. 2016. Use of ecological sites in managing wildlife and livestock: an example with prairie dogs. *Rangelands*

Hendrickson, J.R., S.L. Kronberg, and E.J. Scholljegeredes. 2020. Can targeted grazing reduce abundance of invasive perennial grass (Kentucky Bluegrass) on native mixed-grass prairie? *Rangeland Ecology and Management*, 73:547-551.

Higgins, K.F. 1984. Lightning fires in grasslands in North Dakota and in pine-savanna lands in nearby South Dakota and Montana. *J. Range Manage.* 37:100-103.

Higgins, K.F. 1986. Interpretation and compendium of historical fire accounts in the northern great plains. United States Department of Interior, Fish and Wildlife Service. Resource Publication 161. 39 pages.

Higgins, K.F., A.D. Kruse, and J.L. Piehl. 1989. Effects of fire in the Northern Great Plains. U.S. Fish and Wildlife Service and Cooperative Extension Service South, Dakota State University. Extension Circular 761. 48 pages.

High Plains Regional Climate Center, University of Nebraska, 830728 Chase Hall, Lincoln, NE 68583-0728. (<http://hprcc.unl.edu>)

Johnson, Sandra. 2015. Reptiles and Amphibians of North Dakota. North Dakota Game and Fish Department. 64 pages.

Jordan, N. R., D.L. Larson, and S.C. Huerd. 2008. Soil modification by invasive plants: effects on native and invasive species of mixed-grass prairies. *Biological Invasions* 10:177-190.

Mader, E., M. Shepherd, M. Vaughan, and S.H. Black. 2011. [Attracting native pollinators: protecting North America's bees and butterflies](https://xerces.org). Accessed at <https://xerces.org>, May 1, 2017.

Minnesota Department of Natural Resources. 2005. Field guide to the native plant communities of Minnesota – the prairie parkland and tallgrass aspen parklands provinces. Minnesota DNR.

Nelson, W.T. 1986. Grassland habitat type classification of the Sheyenne National Grasslands of southeastern North Dakota. M.S. thesis. North Dakota State University. 139 pages.

North Dakota Division of Tourism, Accessed on February 25, 2019. Available at  
<https://www.ndtourism.com/sports-recreation>

North Dakota Parks and Recreation Department, Accessed on February 25, 2019. Available at  
<http://www.parkrec.nd.gov/recreationareas/recreationareas.html>

Palit, R., G. and E.S. DeKeyser. 2022. Impacts and drivers of smooth brome (*Bromus inermis* Leyes.) invasion in native ecosystems. *Plants*: 10,3390. <https://www.mdpi.com/2223-7747/11/10/1340>

Palit, R., G. Gramig, and E.S. DeKeyser. 2021. Kentucky bluegrass invasion in the Northern Great Plains and prospective management approaches to mitigate its spread. *Plants*: 10,817. <https://doi.org/10.3390/plants10040817>

Peterson, K. 2013. Remediation of Sand Dune Blowouts Along Pipeline Rights of Ways. M.S. thesis. University of New Mexico, Albuquerque.  
[https://digitalrepository.unm.edu/cgi/viewcontent.cgi?article=1015&context=geog\\_etds](https://digitalrepository.unm.edu/cgi/viewcontent.cgi?article=1015&context=geog_etds)

Printz, J.L. and J.R. Hendrickson. 2015. Impacts of Kentucky bluegrass Invasion (*Poa pratensis*) on Ecological Processes in the Northern Great Plains. *Rangelands* 37(6):226-232.

Reeves, J.L., J.D. Derner, M.A. Sanderson, J.R. Hendrickson, S.L. Kronberg, M.K. Petersen, and L.T. Vermeire. 2014. Seasonal weather influences on yearling beef steer production in C<sub>3</sub>-dominated Northern Great Plains rangeland. *Agriculture, Ecosystems and Environment* 183:110-117.

Ralston, R.D., and R.L. Dix. 1966. Green herbage production of native grasslands in the Red River Valley-1965. *Proceedings ND Academy of Science* 20:57-66.

Royer, R. A., 2003. Butterflies of North Dakota: An Atlas and Guide. Minot State University, Minot, ND.

Seabloom, R. 2020. Mammals of North Dakota. North Dakota Institute for Regional Studies, Fargo, ND. 470 pages.

Sedivec, K.D., J.L. Printz. 2014. Ranchers Guide to Grassland Management IV. NDSU Extension Service publication R1707.

Severson, K. E. and C. Hull Sieg. 2006. The Nature of Eastern North Dakota: Pre-1880 Historical Ecology. North Dakota Institute for Regional Studies.

South Dakota Dept. of Game, Fish and Parks. 2014. South Dakota Wildlife Action Plan. Wildlife Division Report 2014-03.

Spaeth, K.E., Hayek, M.A., Toledo, D., and Hendrickson, J. 2019. Cool Season Grass Impacts on Native Mixedgrass Prairie Species in the Northern Great Plains. America's Grassland Conference: Working Across Boundaries. The Fifth Biennial Conference on the Conservation of America's Grasslands. Bismarck, ND. 20-22 August.

Swingen, M., R. M. Walker, R. Baker, G. Nordquist, T. Catton, K. Kirschbaum, B. Dirks, and N. Dietz. 2018. Northern Long-eared Bat Roost Tree Characteristics 2015-2017. Natural Research Institute, University of Minnesota Duluth, Technical Report NRRI/TR-2018/41, 88p.

Tidwell, D., D.T. Fogarty, and J.R. Weir. 2021. Woody encroachment in grasslands, a guide for understanding risk and vulnerability. Oklahoma State University, Oklahoma Cooperative Extension Service publication E-1054. 32 pages.

Toledo, D., M. Sanderson, K. Spaeth, J. Hendrickson, and J. Printz. 2014. Extent of Kentucky bluegrass and its effect on native plant species diversity and ecosystem services in the northern great plains of the United States. *Invasive Plant Science and Management* 7(4): 543-552.

USDA, NRCS. 2021. National Range and Pasture Handbook, <https://www.nrcs.usda.gov/wps/portal/nrcs/detailfull/national/landuse/rangepasture/?cid=stelprdb1043084>

Site Type: Rangeland  
MLRA: 56A – Red River Valley of the North

**Sands**  
**R056AY090ND**

USDA, NRCS. 2006. Land Resource Regions and Major Land Resource Areas of the United States, the Caribbean, and the Pacific Basin. U.S. Department of Agriculture Handbook 296.

USDA, NRCS. National Soil Information System, 100 Centennial Mall North, Room 152, Lincoln, NE 68508-3866.

([https://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/survey/tools/?cid=nrcs142p2\\_053552](https://www.nrcs.usda.gov/wps/portal/nrcs/detail/soils/survey/tools/?cid=nrcs142p2_053552))

USDA, NRCS. National Water & Climate Center, 1201 NE Lloyd Blvd, Suite 802, Portland, OR 97232-1274. (<https://www.wcc.nrcs.usda.gov/>)

USDA, NRCS. 2001. The PLANTS Database, Version 3.1 (<http://plants.usda.gov>). National Plant Data Center, Baton Rouge, LA 70874-4490 USA.

USDA, NRCS, Various Published Soil Surveys.

USDI BLM.1999. Utilization Studies and Residual Measurements. Interagency Technical Reference 1734-3.

Vinton, M.A. and E.M. Goergen. 2006. Plant-soil feedbacks contribute to the persistence of *Bromus inermis* in tallgrass prairie. *Ecosystems* 9: 967-976.

Whitman, W.H., H. Hanson, and R. Peterson. 1943. Relation of drought and grazing to North Dakota range lands. *North Dakota Agricultural Experimentation Bulletin* 340.

## Site Description Approval

---

ND, State Range Management Specialist      Date

---

MN, State Range Management Specialist      Date

---

SD, State Range Management Specialist      Date

**INTERPRETING INDICATORS OF RANGELAND HEALTH, Version 5, REFERENCE SHEET**Ecological site name: Sands Ecological site code: RO56AY090NDAuthor(s)/participant(s): USAD-NRCS North DakotaContact for lead author: NRCS State Rangeland Management SpecialistDate: Dec. 2021 MLRA: 56 LRU: \_\_\_\_\_Composition based on (check one):  Cover  Annual Production

**Indicators.** For each indicator, describe the potential for the site using the reference sheet checklist. Where possible, (1) use quantitative measurements; (2) include expected range of values for above- and below-average years and natural disturbance regimes for each community phase within the reference state, when appropriate; and (3) cite data sources used. Continue descriptions on separate sheet.

**1. Rills:** Rills are not expected on this site.

**2. Water flow patterns:** Water flow patterns are not visible.

**3. Pedestals and/or terracettes:** Neither pedestals nor terracettes are expected.

**4. Bare ground:** Bare ground is 5% or less. Bare ground patches should be small (less than 4 inches in diameter) and not connected. Animal activity (burrows and ant mounds) may occasionally result in isolated bare patches of up to 24 inches in diameter.

**5. Gullies:** Active gullies are not expected on this site.

**6. Wind-scoured and/or depositional areas:** Active blowouts are not expected. Small areas of exposed/wind-blown sand less than 6 feet in diameter may be associated with localized animal disturbances and may be compounded by drought. If present, previously active blowouts and depositional areas are well vegetated with no active erosion/deposition visible.

**7. Litter movement:** Plant litter movement not expected on this site.

**8. Soil surface resistance to erosion:** Stability class anticipated to average 5 or greater.

**9. Soil surface loss and degradation:** Use soil series description for depth, color, and structure of A-horizon.

**10. Effects of plant community composition and distribution on infiltration:** Tall-statured rhizomatous grass, and mid- and short-statured bunch grasses are dominant and well distributed across the state. A diverse forb component, mid- and short-statured rhizomatous grasses and grass-likes are subdominant.

**11. Compaction layer:** No compaction layers occur naturally on this site.

**12. Functional/structural groups:** Due to differences in phenology, root morphology, soil biology relationships, and nutrient cycling Kentucky bluegrass, smooth brome, and crested wheatgrass are included in a new Functional/structural group, mid- and short-statured early cool-season grasses (MSeC3), **not expected for this site.**

Dominance Category <sup>1</sup>	Relative dominance of F/S groups for community phases in the <i>Reference State</i> <i>Minimum expected number of species for dominant and subdominant groups is included in parentheses.</i>		
	Dominance based on <sup>1</sup> : Annual Production <u>X</u> or Foliar Cover <u> </u>		
	Phase 1.1_	Phase 1. __	Phase 1. __
Dominant	Tall C4 rhizomatous grasses (3); Mid & short C3 bunch grasses (4); Mid & short C4 bunch grasses (3)		
Subdominant	Forbs (22); Mid & short C4 rhizomatous grasses (1); Grass-likes (2)		
Minor	Shrub; Mid & short C3 rhizomatous grasses		
Trace			
<sup>1</sup> Biological soil crust dominance is determined based on cover, rather than production. If biological soil crusts are an expected dominant or subdominant group, the number of expected life forms (e.g., lichen, moss) is listed, rather than number of individual species.			
<b>13. Dead or dying plants or plant parts:</b> Rare or not occurring on this site.			
<b>14. Litter cover and depth:</b> Plant litter cover is 60 to 70% with a depth of 0.25 to 0.50 inches. Litter is in contact with the soil surface.			
<b>15. Annual production:</b> Annual air-dry production is 2600 lbs./ac (reference value) with normal precipitation and temperatures. Low and high production years should yield 1800 lbs./ac to 3200 lbs./ac, respectively.			
<b>16. Invasive plants:</b> State and local noxious species, Kentucky bluegrass, smooth bromegrass, crested wheatgrass, quackgrass, Siberian elm, and Eastern red cedar/juniper.			
<b>17. Vigor with an emphasis on reproductive capability of perennial plants:</b> Noninvasive species in all functional/structural groups are vigorous and capable of reproducing annually under normal weather conditions.			

# Functional/Structural Groups Sheet

State \_\_\_\_\_ Office \_\_\_\_\_ Ecological site \_\_\_\_\_ Ecol. site code \_\_\_\_\_

Observers \_\_\_\_\_ Date \_\_\_\_\_

Evaluation site ID and/or name: \_\_\_\_\_

Dominance in ESD based on: Foliar Cover **Annual Production** Biomass

Species list of functional/structural groups in the <b>Reference State</b>												
Functional/Structural Group	Species List											
Biological soil crust <sup>1</sup>												
<b>Reference State</b> - Relative dominance of functional/structural groups for each community phase												
<i>Relative dominance annotations: Use the following annotations in the narrow columns to describe the relative dominance of the listed functional/structural groups: = "equal"; &gt; "greater than"; &gt;&gt; "much greater than"</i>												
Phase	Dominant **	>>	Subdominant **	>>	Minor **	>>						
		>		>		=						
		=		=								

\* Indicates species that may or may not be present on the site. Absence of these species may not constitute a departure.

\*\* See IIRH Version 5 page 70.

Circle the community phase that most closely matches the evaluation area. \*Revise functional/structural groups relative dominance for the community phase circled to represent changes in dominance given the time since disturbance(s) (see page 1 of site evaluation sheet).

Species list of functional/structural groups in the <b>Evaluation Area</b>						
Functional/Structural Group		Species List				
Tall C4 rhizomatous grasses						
Mid & short C3 bunch grasses						
Mid & short C4 bunch grasses						
Forbs						
Mid & short C4 rhizomatous grasses						
Grass-likes						
Shrub						
Mid & short C3 rhizomatous grasses						
<u>Groups not expected:</u>						
Mid & short early C3 grasses						
Biological soil crust <sup>1</sup>						
<b>Evaluation Area - Relative dominance of functional/structural groups</b>						
Dominant **	>>	Subdominant **	>>	Minor **	>>	Trace **
	>		>		>	
	=		=		=	

**Biological soil crust**<sup>1</sup> - dominance is evaluated solely on cover, not composition by weight

\*\* See IIRH Version 5 page 70.