

## FORAGE SUITABILITY GROUP


### Overflow

**FSG No.:** G053AY500MT

**Major Land Resource Area:** 053A - Northern Dark Brown Glaciated Plains

### Physiographic Features

The soils in this group are generally found in nearly level to gently sloping positions on stream terraces and flood plains, and in swales and drainage ways on uplands. They receive beneficial additional moisture as run-on from up slope, or from flooding.



	<u>Minimum</u>	<u>Maximum</u>
<b>Elevation (feet):</b>	2000	3000
<b>Slope (percent):</b>	0	3
<b>Flooding:</b>		
<b>Frequency:</b>	None	Occasional
<b>Duration:</b>	None	Brief
<b>Ponding:</b>		
<b>Depth (inches):</b>		
<b>Frequency:</b>	None	None
<b>Duration:</b>	None	None
<b>Runoff Class:</b>	Low	Medium

### Climatic Features

This group occurs in a mid-continental climate characterized by wide seasonal temperature and precipitation fluctuations and extremes.

Annual precipitation varies widely from year to year in MLRA 53A. Average annual precipitation for all climate stations listed below is about 13 inches. About 80 percent of that occurs during the months of April through September. On average there are about 22 days with greater than .1 inches of precipitation during the same time frame. Precipitation is less than needed for optimum forage production and is the single largest factor limiting production from this group on non-irrigated lands.

Average annual snowfall ranges from 9 inches at Redstone, MT to 42 inches at Ophiem 16 SE, MT. Snow cover at depths greater than 1 inch range from 16 days at Ophiem 10 N, MT to 97 days at Bredette, MT.

Average July temperatures are about 69 degrees F., and average January temperatures are about 9 degrees F. Recorded temperature extremes in the MLRA during the years 1961 to 1990 are a low of -52 at Redstone, MT, and a high of 110 at 3 Montana locations. The average dates of last and first frost (32 deg) for the listed stations are May 18 and September 14 for an average growing season length of 118 days. The MLRA lies in USDA Plant Hardiness Zones 3b and 4a.

At Williston, ND the average annual wind speeds are about 10 MPH. The highest wind speeds occur during March through June, but average monthly wind speeds do not vary significantly throughout the year. It is cloudy about 160 days a year with the lowest incidence of cloudiness occurring during the summer months. Average morning relative humidity in June is about 81 percent and average afternoon humidity is 54 percent.

The climate data listed in the tables below represent high and low ranges and averages for the climate stations and dates listed. For additional climate data access the National Water and Climate Center at <http://www.wcc.nrcs.usda.gov>.

	<b>From</b>	<b>To</b>
<b>Freeze-free period (28 deg)(days):</b> (9 years in 10 at least)	89	139
<b>Last Killing Freeze in Spring (28 deg):</b> (1 year in 10 later than)	Jun 06	May 13
<b>Last Frost in Spring (32 deg):</b> (1 year in 10 later than)	Jun 28	May 27
<b>First Frost in Fall (32 deg):</b> (1 year in 10 earlier than)	Aug 09	Sep 07
<b>First Killing Freeze in Fall (28 deg):</b> (1 year in 10 earlier than)	Aug 23	Sep 15
<b>Length of Growing Season (32 deg)(days):</b> (9 years in 10 at least)	52	126
<b>Growing Degree Days (40 deg):</b>	3216	4334
<b>Growing Degree Days (50 deg):</b>		
<b>Annual Minimum Temperature:</b>	-40	-25
<b>Mean annual precipitation (inches):</b>	12	15

**Monthly precipitation (inches) and temperature (F):**

<b>2 years in 10:</b>	<b><u>Jan</u></b>	<b><u>Feb</u></b>	<b><u>Mar</u></b>	<b><u>Apr</u></b>	<b><u>May</u></b>	<b><u>Jun</u></b>	<b><u>Jul</u></b>	<b><u>Aug</u></b>	<b><u>Sep</u></b>	<b><u>Oct</u></b>	<b><u>Nov</u></b>	<b><u>Dec</u></b>
<b>Precip. Less Than</b>	0.14	0.10	0.20	0.42	0.90	1.23	0.88	0.48	0.51	0.20	0.13	0.19
<b>Precip. More Than</b>	0.64	0.53	0.82	1.69	3.20	3.55	2.99	2.40	2.12	1.18	0.54	0.65
<b>Monthly Average:</b>	0.39	0.31	0.51	1.10	2.13	2.47	2.02	1.53	1.38	0.69	0.34	0.42
<b>Temp. Min.</b>	-1.8	5.0	16.5	29.5	40.8	49.7	53.9	51.1	40.6	30.2	15.4	3.8
<b>Temp. Max.</b>	19.2	26.5	39.3	55.9	68.5	77.8	84.2	82.3	69.5	56.8	36.8	24.7
<b>Temp. Avg.</b>	8.7	15.8	27.9	42.7	54.7	63.7	69.0	66.7	55.1	43.5	26.1	14.2

<b><u>Climate Station</u></b>	<b><u>Location</u></b>	<b><u>From</u></b>	<b><u>To</u></b>
MT1088	Bredette, MT	1961	1990
MT2122	Culbertson, MT	1961	1990
MT5285	Lustre, MT	1961	1990
MT5572	Medicine Lake 3 SE, MT	1961	1990
MT6236	Ophiem 10 N, MT	1961	1990
MT6238	Ophiem 16 SE, MT	1961	1990
MT6660	Poplar 2E, MT	1961	1990
MT6893	Raymond Border Stn, MT	1961	1990
MT6927	Redstone, MT	1961	1990
MT7424	Scobey, MT	1961	1990
MT8777	Westby, MT	1961	1990
ND1871	Crosby, ND	1961	1990
ND3196	Fortuna, ND	1964	1990
ND3736	Grenora, ND	1961	1990
ND8737	Tioga, ND	1961	1990
ND9400	Wildrose, ND	1961	1990
ND9425	Williston WSO AP, ND	1961	1990
ND9430	Williston Exp Farm, ND	1961	1990

**Soil Interpretations**

This group consists of very deep, moderately well and well drained, medium to moderately fine textured soils formed from glacial till and silty and loamy sediments and alluvium. Permeability is moderately slow to moderate, and available water capacity is high. Some have a watertable usually below 36 inches which may benefit deep rooted plants.

**Drainage Class:** Moderately well drained To Well drained  
**Permeability Class:** Moderately slow To Moderate  
 (0 - 40 inches)  
**Frost Action Class:** Moderate To Moderate

	<u>Minimum</u>	<u>Maximum</u>
<b>Depth (inches):</b>	72	
<b>Surface Fragments &gt;3" (% Cover):</b>		
<b>Organic Matter (percent):</b> (surface layer)	0.5	6.0
<b>Electrical Conductivity (mmhos/cm):</b> (0 - 24 inches)	0	3
<b>Sodium Absorption Ratio:</b> (0 - 12 inches)	0	0
<b>Soil Reaction (1:1) Water (pH):</b> (0 - 12 inches)	6.1	8.4
<b>Available Water Capacity (inches):</b> (0 - 60 inches)	10	11
<b>Calcium Carbonate Equivalent (percent):</b> (0 - 12 inches)	0	10

**Adapted Species List**

The following forage species are considered adapted to grow on the soils in this group. Additional information concerning plant characteristics of a number of the listed species as well as individual cultivars of many of those species can be accessed on the web at <http://plants.usda.gov/>.

	<u>Symbol</u>	<u>Dryland</u>	<u>Irrigated</u>
<b>Cool Season Grasses</b>			
Altai wildrye	LEAN3	G	NS
Basin wildrye	LECI4	F	NS
Bluebunch/Quackgrass Hybrid		G	G
Canada wildrye	ELCA4	G	NS
Creeping foxtail	ALAR	F	NS
Crested wheatgrass	AGCR	F	NS
Green needlegrass	NAVI4	G	NS
Intermediate wheatgrass	THIN6	G	G
Meadow bromegrass	BRBI2	G	G
Pubescent wheatgrass	THIN6	G	G
Reed canarygrass	PHAR3	F	G
Russian wildrye	PSJU3	G	NS
Slender wheatgrass	ELTR7	G	NS
Smooth bromegrass	BRINI2	G	G
Tall wheatgrass	THPO7	F	NS
Western wheatgrass	PASM	G	NS
<b>Warm Season Grasses</b>			
Big bluestem	ANGE	G	G
Indiangrass	SONU2	G	G
Little bluestem	SCSC	F	NS
Sideoats grama	BOCU	G	NS
Switchgrass	PAVIV	G	G
<b>Legumes</b>			
Alfalfa	MESA	G	G
Alsike clover	TRHY	F	NS
Canada milkvetch	ASCAC6	G	NS
Cicer milkvetch	ASCI4	G	F
Purple prairieclover	DAPUP	F	NS
White prairieclover	DACAC	F	NS

G - Good adaptation for forage production on this group of soils in this MLRA  
F - Fair adaptation but will not produce at its highest potential  
NS - Not suited

**Production Estimates**

Production estimates listed here should only be used for making general management recommendations. On site production information should always be used for making detailed planning and management recommendations.

The high forage production estimates listed below are based on dense, vigorous stands of climatically adapted, superior performing cultivars. They are properly fertilized for high yields, and pest infestations are kept below economic thresholds. Mechanical harvests are managed to maintain stand life by cutting at appropriate stages of maturity and harvest intervals. If grazed, optimum beginning and ending grazing heights are adhered to. Adequate time is allowed for plant recovery before entering winter dormancy under both uses.

The production estimates listed below represent total annual above ground plant production on an air-dry-matter basis. Estimates of hay and grazing yields can be calculated from these numbers by multiplying them by a harvest efficiency. A 70 percent harvest efficiency is commonly used when converting to hay yields. Pasture harvest efficiency is highly dependent on the grazing management system applied, ranging from 25 to 50 percent.

Forage Crop	<u>Dryland</u>		<u>Irrigated</u>	
	Management Intensity		Management Intensity	
	<u>Low</u> (lbs/ac)	<u>High</u> (lbs/ac)	<u>Low</u> (lbs/ac)	<u>High</u> (lbs/ac)
Alfalfa	4500	9800	8600	14300
Alfalfa/Intermediate wheatgrass	3600	7700	8600	14300
Alfalfa/Smooth bromegrass	3600	7700	8600	17100
Big bluestem	3500	7000	0	0
Indiangrass	3200	5100	0	0
Intermediate wheatgrass	2900	6000	0	0
Smooth bromegrass	2900	6000	6900	11400
Switchgrass	3700	7600	6900	11400

**Forage Growth Curves**

Growth curves estimate the seasonal distribution of growth of the various forage crops. They indicate when the forages may be available for grazing or mechanical harvest.

**Growth Curve Number:** MT0004

**Growth Curve Name:** Alfalfa 2 Cuttings Dry

**Growth Curve Description:** Dryland, 2 Cuttings For Hay Then Graze Aftermath

**Percent Production by Month**

<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	5	35	30	15	5	5	0	0	0

**Growth Curve Number:** MT0005

**Growth Curve Name:** 3 Cutting Irrigated Alfalfa

**Growth Curve Description:** Irrigated 3 Cuttings for Hay Graze Aftermath

**Percent Production by Month**

<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	5	25	25	20	20	10	0	0	0

**Growth Curve Number:** MT0001

**Growth Curve Name:** Cool Season Grass Irrigated

**Growth Curve Description:** Irrigated Cool Season Grass

**Percent Production by Month**

<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	5	35	30	20	10	10	0	0	0

**Growth Curve Number:** MT0002  
**Growth Curve Name:** Cool Season Grass Dryland  
**Growth Curve Description:** Dryland Cool Season Grass

**Percent Production by Month**

<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	5	40	40	10	0	5	0	0	0

**Growth Curve Number:** MT0003  
**Growth Curve Name:** Warm Season Grass Dryland  
**Growth Curve Description:** Dryland Warm Season Grass

**Percent Production by Month**

<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	0	10	30	40	20	0	0	0	0

**Soil Limitations**

These soils have few limitations to the production of climatically adapted forage crops. Production potential is high. Flooding is a potential hazard to some of these soils. Also, all of these soils receive additional moisture, so the potential exists for soil compaction from grazing or operating machinery on them when wet.

**Management Interpretations**

Soils in this group that are subject to flooding can have forage production adversely impacted if it occurs during the spring or growing season. Flooding duration or the time period plants are under water is more important than flooding frequency for the survival of forage crops. If these soils flood it is generally for only a brief time. Exclude livestock and machinery during extended periods of soil wetness to reduce soil compaction. When establishing new stands or renovating stands select species and varieties that can make best use of the additional soil moisture this group receives.

Pasture and hayland can include considerations for wildlife. Delaying grazing on portions of the pasture or rotating pastures will allow nest initiation of grassland nesting birds or species of concern. Nest initiation of most grassland nesting birds occurs from April 15 to June 1. Delaying haying until after July 15 allows for most species to fledge their young. Consider planting species with later maturity to allow for harvesting after nests have fledged. Avoid mowing around the field. Mow back and forth or from the inside to the outside of the field. Consider using flushing bars on swathers and mowers.

**FSG Documentation**

**Similar FSGs:**

<u>FSG ID</u>	<u>FSG Narrative</u>
G053AY100MT	Loamy soils do not receive the additional water and are less productive.
G053AY700MT	Subirrigated soils have elevated watertables between 18-48 of the surface during part of the growing season.

**Inventory Data References:**

- Agriculture Handbook 296-Land Resource Regions and Major Land Resource Areas
- Natural Resources Conservation Service (NRCS) National Water and Climate Center data
- USDA Plant Hardiness Zone Maps
- National Soil Survey Information System (NASIS) for soil surveys in Montana and North Dakota counties in MLRA 53A
- Montana and North Dakota NRCS Field Office Technical Guides
- NRCS National Range and Pasture Handbook
- Various Agricultural Research Service, Cooperative Extension Service, and NRCS research trials for plant adaptation and production.

**State Correlation:**

This site has been correlated with the following states: Montana and North Dakota

**Forage Suitability Group Approval:**

Original Author: Tim Nordquist  
Original Date: 10/1/200  
Approval by: Loretta J. Metz (MT) and Jeff Printz (ND)  
Approval Date: March 2005