

FORAGE SUITABILITY GROUP

Claypan


FSG No.: G053AY800MT

Major Land Resource 053A -Northern Dark Brown Glaciated Plains

Physiographic Features

These soils are found on till and outwash plains.

	<u>Minimum</u>	<u>Maximum</u>
Elevation (feet):	2000	3000
Slope (percent):	0	6
Flooding:		
Frequency:	None	None
Duration:	None	None
Ponding:		
Depth (inches):		
Frequency:	None	None
Duration:	None	None
Runoff Class:	Medium	Very high



Climatic Features

This group occurs in a mid-continental climate characterized by wide seasonal temperature and precipitation fluctuations and extremes.

Annual precipitation varies widely from year to year in MLRA 53A. Average annual precipitation for all climate stations listed below is about 13 inches. About 80 percent of that occurs during the months of April through September. On average there are about 22 days with greater than .1 inches of precipitation during the same time frame.

Average annual snowfall ranges from 9 inches at Redstone, MT to 42 inches at Ophiem 16 SE, MT. Snow cover at depths greater than 1 inch range from 16 days at Ophiem 10 N, MT to 97 days at Bredette, MT.

Average July temperatures are about 69 degrees F., and average January temperatures are about 9 degrees F. Recorded temperature extremes in the MLRA during the years 1961 to 1990 are a low of -52 at Redstone, MT, and a high of 110 at 3 Montana locations. The average dates of last and first frost (32 deg) for the listed stations are May 18 and September 14 for an average growing season length of 118 days. The MLRA lies in USDA Plant Hardiness Zones 3b and 4a.

At Williston, ND the average annual wind speeds are about 10 MPH. The highest wind speeds occur during March though June, but average monthly wind speeds do not vary significantly throughout the year. It is cloudy about 160 days a year with the lowest incidence of cloudiness occurring during the summer months. Average morning relative humidity in June is about 81 percent and average afternoon humidity is 54 percent.

The climate data listed in the tables below represent high and low ranges and averages for the climate stations and dates listed. For additional climate data access the National Water and Climate Center at <http://www.wcc.nrcs.usda.gov>.

	From	To
Freeze-free period (28 deg)(days): (9 years in 10 at least)	89	139
Last Killing Freeze in Spring (28 deg): (1 year in 10 later than)	Jun 06	May 13
Last Frost in Spring (32 deg): (1 year in 10 later than)	Jun 28	May 27
First Frost in Fall (32 deg): (1 year in 10 earlier than)	Aug 09	Sep 07
First Killing Freeze in Fall (28 deg): (1 year in 10 earlier than)	Aug 23	Sep 15
Length of Growing Season (32 deg)(days): (9 years in 10 at least)	52	126
Growing Degree Days (40 deg):	3216	4334
Growing Degree Days (50 deg):		
Annual Minimum Temperature:	-40	-25
Mean annual precipitation (inches):	12	15

Monthly precipitation (inches) and temperature (F):

2 years in 10:	<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
Precip. Less Than	0.14	0.10	0.20	0.42	0.90	1.23	0.88	0.48	0.51	0.20	0.13	0.19
Precip. More Than	0.64	0.53	0.82	1.69	3.20	3.55	2.99	2.40	2.12	1.18	0.54	0.65
Monthly Average:	0.39	0.31	0.51	1.10	2.13	2.47	2.02	1.53	1.38	0.69	0.34	0.42
Temp. Min.	-1.8	5.0	16.5	29.5	40.8	49.7	53.9	51.1	40.6	30.2	15.4	3.8
Temp. Max.	19.2	26.5	39.3	55.9	68.5	77.8	84.2	82.3	69.5	56.8	36.8	24.7
Temp. Avg.	8.7	15.8	27.9	42.7	54.7	63.7	69.0	66.7	55.1	43.5	26.1	14.2

<u>Climate Station</u>	<u>Location</u>	<u>From</u>	<u>To</u>
MT1088	Bredette, MT	1961	1990
MT2122	Culbertson, MT	1961	1990
MT5285	Lustre, MT	1961	1990
MT5572	Medicine Lake 3 SE, MT	1961	1990
MT6236	Ophiem 10 N, MT	1961	1990
MT6238	Ophiem 16 SE, MT	1961	1990
MT6660	Poplar 2E, MT	1961	1990
MT6893	Raymond Border Stn, MT	1961	1990
MT6927	Redstone, MT	1961	1990
MT7424	Scobey, MT	1961	1990
MT8777	Westby, MT	1961	1990
ND1871	Crosby, ND	1961	1990
ND3196	Fortuna, ND	1964	1990
ND3736	Grenora, ND	1961	1990
ND8737	Tioga, ND	1961	1990
ND9400	Wildrose, ND	1961	1990
ND9425	Williston WSO AP, ND	1961	1990
ND9430	Williston Exp Farm, ND	1961	1990

Soil Interpretations

This group consists of moderately well and well drained, medium to moderately coarse textured soils formed from glacial till and glacial outwash sediments. They have claypan subsoils with slow permeability. The underlying material and lower part of the subsoil typically have high amounts of soluble salts and are alkaline.

Drainage Class:	Moderately well drained	To	Well drained
Permeability Class: (0 - 40 inches)	Slow	To	Slow
Frost Action Class:	Moderate	To	Moderate

	<u>Minimum</u>	<u>Maximum</u>
Depth (inches):	72	
Surface Fragments >3" (% Cover):		
Organic Matter (percent): (surface layer)	1.0	5.0
Electrical Conductivity (mmhos/cm): (0 - 24 inches)	0	8
Sodium Absorption Ratio: (0 - 12 inches)	0	20
Soil Reaction (1:1) Water (pH): (0 - 12 inches)	0	0
Available Water Capacity (inches): (0 - 60 inches)	8	8
Calcium Carbonate Equivalent (percent): (0 - 12 inches)	0	20

Adapted Species List

The following forage species are considered adapted to grow on the soils in this group. Additional information concerning plant characteristics of a number of the listed species as well as individual cultivars of many of those species can be accessed on the web at <http://plants.usda.gov/>.

Cool Season Grasses	<u>Symbol</u>	<u>Dryland</u>
Bluebunch/Quackgrass Hybrid		F
Crested wheatgrass	AGCR	F
Green needlegrass	NAVI4	F
Slender wheatgrass	ELTR7	G
Tall wheatgrass	THPO7	F
Western wheatgrass	PASM	G
Legumes	<u>Symbol</u>	<u>Dryland</u>
Alfalfa	MESA	F

G - Good adaptation for forage production on this group of soils in this MLRA

F - Fair adaptation but will not produce at its highest potential

Production Estimates

Production estimates listed here should only be used for making general management recommendations. On site production information should always be used for making detailed planning and management recommendations.

The high forage production estimates listed below are based on dense, vigorous stands of climatically adapted, superior performing cultivars. They are properly fertilized for high yields, and pest infestations are kept below economic thresholds. Mechanical harvests are managed to maintain stand life by cutting at appropriate stages of maturity and harvest intervals. If grazed, optimum beginning and ending grazing heights are adhered to. Adequate time is allowed for plant recovery before entering winter dormancy under both uses.

The production estimates listed below represent total annual above ground plant production on an air-dry-matter basis. Estimates of hay and grazing yields can be calculated from these numbers by multiplying them by a harvest efficiency. A 70 percent harvest efficiency is commonly used when converting to hay yields. Pasture harvest efficiency is highly dependent on the grazing management system applied, ranging from 25 to 50 percent.

Dryland

Forage Crop Management Intensity

	<u>Low</u>	<u>High</u>
Alfalfa	2300	4600
Alfalfa/Crested wheatgrass	1800	3500
Alfalfa/Intermediate wheatgrass	1900	3700
Crested wheatgrass	1600	3400
Intermediate wheatgrass	1600	3600
Tall wheatgrass	1600	3600
Western wheatgrass	1100	2400

Forage Growth Curves

Growth curves estimate the seasonal distribution of growth of the various forage crops. They indicate when the forages may be available for grazing or mechanical harvest.

Growth Curve Number: MT0004

Growth Curve Name: Alfalfa 2 Cuttings Dry

Growth Curve Description: Dryland, 2 Cuttings For Hay Then Graze Aftermath

Percent Production by Month

<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	5	35	30	15	5	5	0	0	0

Growth Curve Number: MT0002

Growth Curve Name: Cool Season Grass Dryland

Growth Curve Description: Dryland Cool Season Grass

Percent Production by Month

<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	5	40	40	10	0	5	0	0	0

Growth Curve Number: MT0003

Growth Curve Name: Warm Season Grass Dryland

Growth Curve Description: Dryland Warm Season Grass

Percent Production by Month

<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	0	10	30	40	20	0	0	0	0

Soil Limitations

These soils have severe limitations to the production of climatically adapted forage species. The claypan and the soluble salts and sodicity in the subsoil produce an unfavorable rooting environment, limiting species selection and production potential.

Management Interpretations

The impact on yields can be reduced by selecting forage species that are tolerant of salinity and sodicity and can root in dense, clayey subsoils.

Pasture and hayland can include considerations for wildlife. Delaying grazing on portions of the pasture or rotating pastures will allow nest initiation of grassland nesting birds or species of concern. Nest initiation of most grassland nesting birds occurs from April 15 to June 1. Delaying haying until after July 15 allows for most species to fledge their young. Consider planting species with later maturity to allow for harvesting after nests have fledged. Avoid mowing around the field. Mow back and forth or from the inside to the outside of the field. Consider using flushing bars on swathers and mowers.

FSG Documentation

Similar FSGs:

FSG ID

G053AY210MT

FSG Narrative

Clayey subsoils are less saline and/or sodic and have a more favorable rooting zone.

Inventory Data References:

- Agriculture Handbook 296-Land Resource Regions and Major Land Resource Areas
- Natural Resources Conservation Service (NRCS) National Water and Climate Center data
- USDA Plant Hardiness Zone Maps
- National Soil Survey Information System (NASIS) for soil surveys in Montana and North Dakota counties in MLRA 53A
- Montana and North Dakota NRCS Field Office Technical Guides
- NRCS National Range and Pasture Handbook
- Various Agricultural Research Service, Cooperative Extension Service, and NRCS research trials for plant adaptation and production.

State Correlation:

This site has been correlated with the following states: Montana and North Dakota

Forage Suitability Group Approval:

Original Author: Tim Nordquist

Original Date: 10/1/200

Approval by: Loretta J. Metz (MT) and Jeff Printz (ND)

Approval Date: March 2005