

FORAGE SUITABILITY GROUP DROUGHTY LOAM

FSG No.: G055AY120ND

Major Land Resource Area - Northern Black Glaciated Plains

Physiographic Features

The soils in this group are mostly found on glacial lake, outwash, and delta plains and on terraces, deltas, and beaches.

	<u>Minimum</u>	<u>Maximum</u>
Elevation (feet):	980	2300
Slope (percent):	0	15
Flooding:		
Frequency:	None	None
Duration:	None	None
Ponding:		
Depth (inches):		
Frequency:	None	None
Duration:	None	None
Runoff Class:	Very low	Low



Climatic Features

This group occurs in a mid-continental climate characterized by wide seasonal temperature and precipitation fluctuations and extremes.

Annual precipitation varies widely from year to year in MLRA 55A. Average annual precipitation for all climate stations listed below is about 17.5 inches. About 79 percent of that occurs during the months of April through September. On average there are about 27 days with greater than .1 inches of precipitation during the same time frame. Precipitation is less than needed for optimum forage production and is the single largest factor limiting production from this group on non-irrigated lands.

Average annual snowfall ranges from 26 inches at Hannah, ND to 44 inches at Belcourt Keya, ND. Days with snow cover at depths greater than 1 inch range from 64 days at Velva, ND to 123 days at Hannah, ND.

Average July temperatures are about 68 degrees F., and average January temperatures are about 3 degrees F. Recorded temperature extremes in the MLRA during the years 1961 to 1990 are a low of -39 at both Petersburg and Oakes in ND, and a high of 108 recorded at both Granville and Velva. The MLRA lies in USDA Plant Hardiness Zones 3A and 3b.

The climate data listed in the tables below represent high and low ranges and averages for the climate stations and dates listed. For additional climate data access the National Water and Climate Center at <http://www.wcc.nrcs.usda.gov>.

	From	To
Freeze-free period (28 deg)(days): (9 years in 10 at least)	101	128
Last Killing Freeze in Spring (28 deg): (1 year in 10 later than)	Jun 03	May 19

	From	To
Last Frost in Spring (32 deg): (1 year in 10 later than)	Jun 22	May 29
First Frost in Fall (32 deg): (1 year in 10 earlier than)	Aug 21	Sep 11
First Killing Freeze in Fall (28 deg): (1 year in 10 earlier than)	Sep 04	Sep 18
Length of Growing Season (32 deg)(days): (9 years in 10 at least)	64	104
Growing Degree Days (40 deg):	3022	3776
Growing Degree Days (50 deg):	1541	2129
Annual Minimum Temperature:	-40	-30
Mean annual precipitation (inches):	16	19

Monthly precipitation (inches) and temperature (F):

2 years in 10:	<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
Precip. Less Than	0.23	0.18	0.19	0.48	0.93	1.30	0.96	0.82	0.69	0.38	0.15	0.15
Precip. More Than	0.86	0.81	0.99	2.85	3.64	4.14	4.81	4.11	3.18	2.03	0.78	0.78
Monthly Average:	0.49	0.44	0.68	1.42	2.21	3.00	2.75	2.35	1.95	1.14	0.49	0.50
Temp. Min.	-9.2	-2.5	9.9	25.3	37.5	47.2	52.4	49.7	39.2	29.6	14.2	-2.4
Temp. Max.	17.6	25.6	37.8	55.4	69.3	78.2	84.5	83.5	70.6	59.0	38.3	22.6
Temp. Avg.	3.0	9.3	22.5	39.6	53.1	62.6	67.7	65.7	54.4	43.2	25.1	8.9

<u>Climate Station</u>	<u>Location</u>	<u>From</u>	<u>To</u>
ND0626	Belcourt, ND	1961	1987
ND2158	Devils Lake, ND	1961	1990
ND2525	Edmore, ND	1961	1990
ND3686	Granville, ND	1961	1990
ND3936	Hannah, ND	1961	1986
ND4958	Langdon Exp. Farm, ND	1961	1990
ND6025	Mohall, ND	1961	1990
ND7664	Rolla, ND	1961	1990
ND8792	Towner, ND	1961	1990
ND8913	Upham, ND	1961	1990
ND8990	Velva, ND	1961	1986

Soil Interpretations

This group consists of moderately well to well drained, moderately coarse to medium textured soils formed in loamy and sandy materials deposited by wind or water. Available water capacity is in the moderate range due to moderately coarse soil textures or moderate depth to sand and gravel.

Drainage Class:	Moderately well drained	To	Well drained
Permeability Class: (0 - 40 inches)	Moderate	To	Moderately rapid
Frost Action Class:	Low	To	Moderate

	<u>Minimum</u>	<u>Maximum</u>
Depth:	72	
Surface Fragments >3" (% Cover):	0	3
Organic Matter (percent): (surface layer)	1.0	8.0
Electrical Conductivity (mmhos/cm): (0 - 24 inches)	0	2

	<u>Minimum</u>	<u>Maximum</u>
Sodium Absorption Ratio: (0 - 12 inches)	0	1
Soil Reaction (1:1) Water (pH): (0 - 12 inches)	5.6	8.4
Available Water Capacity (inches): (0 - 60 inches)	6	9
Calcium Carbonate Equivalent (percent): (0 - 12 inches)	0	5

Adapted Species List

The following forage species are considered adapted to grow on the soils in this group. Additional information concerning plant characteristics of a number of the listed species as well as individual cultivars of many those species can be accessed at <http://plants.usda.gov/>

Cool Season Grasses			Warm Season Grasses		
	<u>Symbol</u>			<u>Symbol</u>	
Altai wildrye	LEAN3	F	Big bluestem	ANGE	F
Bluebunch/Quackgrass Hybrid		G	Blue grama	BOGR2	G
Canada wildrye	ELCA4	G	Indiangrass	SONU2	F
Crested wheatgrass	AGCR	G	Little bluestem	SCSC	G
Dahurian wildrye	ELDA3	G	Prairie sandreed	CALO	F
Green needlegrass	NAVI4	G	Sand bluestem	ANHA	F
Intermediate wheatgrass	THIN6	G	Sideoats grama	BOCU	G
Meadow bromegrass	BRBI2	G	Switchgrass	PAVIV	F
Pubescent wheatgrass	THIN6	G	Legumes		
Russian wildrye	PSJU3	G	Alfalfa	MESA	G
Slender wheatgrass	ELTR7	G	American vetch	VIAM	G
Smooth bromegrass	BRINI2	G	Birdsfoot trefoil	LOCO6	F
Tall wheatgrass	THPO7	G	Canada milkvetch	ASCAC6	F
Western wheatgrass	PASM	G	Cicer milkvetch	ASCI4	G
			Hairy vetch	VIVI	F
			Purple prairieclover	DAPUP	G
			Red clover	TRPR2	F
			Sainfoin	ONVI	F
			Sweet clover	MELIL	G
			White prairieclover	DACAC	G

G - Good adaptation for forage production on this group of soils in this MLRA
F - Fair adaptation but will not produce at its highest potential

Production Estimates

Production estimates listed here should only be used for making general management recommendations. On site production information should always be used for making detailed planning and management recommendations.

The high forage production estimates listed below are based on dense, vigorous stands of climatically adapted, superior performing cultivars. They are properly fertilized for high yields, and pest infestations are kept below economic thresholds. Mechanical harvests are managed to maintain stand life by cutting at appropriate stages of maturity and harvest intervals. If grazed, optimum beginning and ending grazing heights are adhered to. Adequate time is allowed for plant recovery before entering winter dormancy under both uses.

The production estimates listed below represent total annual above ground plant production on an air-dry-matter basis. Estimates of hay and grazing yields can be calculated from these numbers by multiplying them by a harvest efficiency. A 70 percent harvest efficiency is commonly used when converting to hay yields. Pasture harvest efficiency is highly dependent on the grazing management system applied, ranging from 25 to 50 percent.

Forage Crop	<u>Dryland</u>	
	Management Intensity	
	<u>Low</u> (lbs/ac)	<u>High</u> (lbs/ac)
Alfalfa	3100	7100
Alfalfa/Crested wheatgrass	2400	5100
Alfalfa/Intermediate wheatgrass	2700	5400
Alfalfa/Smooth brome grass	2700	5400
Big bluestem	2500	5400
Crested wheatgrass	2400	4800
Green needlegrass	1500	3100
Intermediate wheatgrass	2500	5100
Smooth brome grass	2500	5100
Switchgrass	2500	6300
Western wheatgrass	1600	3400

Forage Growth Curves

Growth curves estimate the seasonal distribution of growth of the various forage crops. They indicate when the forages may be available for grazing or mechanical harvest.

Growth Curve Number: ND0001
Growth Curve Name: Alfalfa
Growth Curve Description: Alfalfa

<u>Percent Production by Month</u>											
<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	5	25	30	20	15	5	0	0	0

Growth Curve Number: ND0002
Growth Curve Name: Cool season grass
Growth Curve Description: Cool season grass

<u>Percent Production by Month</u>											
<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	5	40	35	10	5	5	0	0	0

Growth Curve Number: ND0003
Growth Curve Name: Warm season grass
Growth Curve Description: Warm season grass

<u>Percent Production by Month</u>											
<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	0	10	40	35	15	0	0	0	0

Soil Limitations

The primary limitation for these soils is their moderate available water capacity which limits plant growth during periods of moisture deficit. On steeper slopes water erosion is a potential problem during establishment. Livestock trail erosion is a potential problem on established stands. Also, wind erosion is a potential problem during stand establishment on moderately coarse textured soils.

Management Interpretations

The impact on yields of the moderate available water capacity of these soils can be reduced by selecting forage species that are tolerant to periods of drought and inadequate soil moisture. Including sod forming grass species in stands, especially on steeper slopes, will reduce the potential for sheet and rill erosion. Incorporate both wind and water erosion control practices during the establishment period. Properly locating facilitating practices such as fences, lanes, and water developments can help control livestock movement, reduce trailing perpendicular to steeper slopes, and evenly distribute grazing pressure.

Pasture and hayland can include considerations for wildlife. Delaying grazing on portions of the pasture or rotating pastures will allow nest initiation of grassland nesting birds or species of concern. Nest initiation of most grassland nesting birds occurs from April 15 to June 1. Delaying haying until after July 15 allows for most species to fledge their young. Consider planting species with later maturity to allow for harvesting after nests have fledged. Avoid mowing around the field. Mow back and forth or from the inside to the outside of the field. Consider using flushing bars on swathers and mowers.

FSG Documentation

Similar FSGs:

FSG ID

FSG Narrative

G055AY100ND

Loamy soils have greater available water capacity and greater production potential.

G055AY130ND

Very Droughty Loams have lower available water capacity and lower production potential.

Inventory Data References:

- Agriculture Handbook 296-Land Resource Regions and Major Land Resource Areas
- Natural Resources Conservation Service (NRCS) National Water and Climate Center data
- USDA Plant Hardiness Zone maps
- National Soil Survey Information System (NASIS) for soil surveys in North Dakota counties in MLRA 55A
- North Dakota NRCS Field Office Technical Guide
- NRCS National Range and Pasture Handbook
- Various Agricultural Research Service, Cooperative Extension Service, and NRCS research trials for plant adaptation and production

State Correlation:

This site has been correlated with the following states: North Dakota

Forage Suitability Group Approval:

Original Author: Tim Nordquist
Original Date: 9/3/2003
Approval by: Jeff Printz
Approval Date: March 2005