

FORAGE SUITABILITY GROUP Overflow

FSG No.: G060AY500SD

Major Land Resource Area: 60A - Pierre Shale Plains and Badlands

Physiographic Features

The soils in this group are generally found in nearly level to gently sloping positions on stream terraces and flood plains. They receive beneficial additional moisture as run-on from up slope, or from flooding.

	<u>Minimum</u>	<u>Maximum</u>
Elevation (feet):	2600	3300
Slope (percent):	0	3
Flooding:		
Frequency:	None	Frequent
Duration:	None	Brief
Ponding:		
Depth (inches):		
Frequency:	None	None
Duration:	None	None
Runoff Class:	Negligible	Medium



Climatic Features

This group occurs in a mid-continental climate characterized by wide seasonal temperature and precipitation fluctuations and extremes.

Annual precipitation varies widely from year to year in MLRA 60A. Average annual precipitation for all climate stations listed below is about 15 inches. About 77% of the annual precipitation occurs during the months of April through September. On average there are about 24 days with greater than .1 inches of precipitation during that same time period. Precipitation is less than needed for optimum forage production and is the single largest factor limiting production from this group on non-irrigated lands.

Average annual snowfall ranges from 25 inches at Newell, SD to 45 inches at Oelrichs, SD. Snow cover at depths greater than 1 inch range from 40 days at Newell, SD to 82 days at Colony, WY.

Average July temperatures across the MLRA are about 74 degrees F., and average January temperatures are about 20 degrees F. Recorded temperature extremes in the MLRA during the years 1961 to 1990 are a low of -47 at Redbird, WY, and a high of 114 recorded at Oelrichs, SD. The MLRA lies mostly in USDA Plant Hardiness Zones 4a and 4b.

At Rapid City, SD, the closest station with records, it is cloudy about 139 days a year. Average morning relative humidity in June is about 78% and average afternoon humidity is 49%.

The climate data listed in the tables below represent high and low ranges and averages for the climate stations and dates listed. For additional climate data access the National Water and Climate Center at

<http://www.wcc.nrcs.usda.gov>

PASTURE AND HAYLAND INTERPRETATIONS

Page 2

	From	To
Freeze-free period (28 deg)(days): (9 years in 10 at least)	118	137
Last Killing Freeze in Spring (28 deg): (1 year in 10 later than)	May 26	May 14
Last Frost in Spring (32 deg): (1 year in 10 later than)	Jun 07	May 26
First Frost in Fall (32 deg): (1 year in 10 earlier than)	Sep 02	Sep 13
First Killing Freeze in Fall (28 deg): (1 year in 10 earlier than)	Sep 11	Sep 21
Length of Growing Season (32 deg)(days): (9 years in 10 at least)	96	117
Growing Degree Days (40 deg):	4231	4913
Growing Degree Days (50 deg):	2400	2852
Annual Minimum Temperature:	-30	-20
Mean annual precipitation (inches):	14	17

Monthly precipitation (inches) and temperature (F):

2 years in 10:	<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
Precip. Less Than	0.11	0.09	0.30	0.65	1.05	1.04	1.06	0.52	0.37	0.38	0.20	0.15
Precip. More Than	0.49	0.74	1.27	2.50	4.02	4.63	2.98	2.22	1.68	1.62	0.89	0.66
Monthly Average:	0.33	0.42	0.83	1.71	2.69	2.78	1.99	1.47	1.24	1.03	0.53	0.41
Temp. Min.	5.3	10.9	20.0	30.6	40.5	49.8	56.3	53.2	41.7	29.9	18.2	6.5
Temp. Max.	34.3	40.5	49.5	61.5	71.8	82.5	91.2	89.8	79.0	65.6	48.3	36.3
Temp. Avg.	19.9	25.3	34.0	45.8	56.0	66.0	73.6	71.5	60.2	48.0	33.5	22.0

<u>Climate Station</u>	<u>Location</u>	<u>From</u>	<u>To</u>
SD0236	Ardmore, SD	1961	1990
SD6054	Newell, SD	1961	1990
SD6212	Oelrichs, SD	1961	1990
SD8911	Wasta, SD	1961	1990
SD9537	Zeona, SD	1961	1990
WY1905	Colony, WY	1961	1990
WY7555	Redbird, WY	1961	1990

Soil Interpretations

This group consists of very deep, mostly well drained, medium to moderately fine textured soils formed from alluvium. Permeability is slow to moderate, and available water capacity is moderate to high.

Drainage Class:	Moderately well drained	To	Well drained
Permeability Class: (0 - 40 inches)	Slow	To	Moderately rapid
Frost Action Class:	Low	To	High

	<u>Minimum</u>	<u>Maximum</u>
Depth:	72	
Surface Fragments >3" (% Cover):	0	3
Organic Matter (percent): (surface layer)	0.8	4.0
Electrical Conductivity (mmhos/cm): (0 - 24 inches)	0	4
Sodium Absorption Ratio: (0 - 12 inches)	0	5
Soil Reaction (1:1) Water (pH): (0 - 12 inches)	6.1	8.4

	<u>Minimum</u>	<u>Maximum</u>
Available Water Capacity (inches): (0 - 60 inches)	8	10
Calcium Carbonate Equivalent (percent): (0 - 12 inches)	0	10

Adapted Species List

The following forage species are considered adapted to grow on the soils in this group. Additional information concerning plant characteristics of a number of the listed species as well as individual cultivars of many of those species can be accessed on the web at <http://www.plants.usda.gov>

Cool Season Grasses	<u>Dryland</u>	<u>Irrigated</u>
Altai wildrye	G	NS
Basin wildrye	G	NS
Crested wheatgrass	G	NS
Green needlegrass	G	NS
Intermediate wheatgrass	G	G
Meadow brome	F	G
Orchardgrass	NS	G
Pubescent wheatgrass	G	G
Smooth brome	F	G
Streambank wheatgrass	F	NS
Thickspike wheatgrass	F	NS
Western wheatgrass	G	NS

Warm Season Grasses	<u>Dryland</u>	<u>Irrigated</u>
Big bluestem	F	G
Little bluestem	G	NS
Sideoats grama	F	NS
Switchgrass	F	G

Legumes	<u>Dryland</u>	<u>Irrigated</u>
Alfalfa	G	G
Birdsfoot trefoil	NS	G
Canada milkvetch	F	NS
Cicer milkvetch	G	F
Purple prairieclover	F	NS
Red clover	NS	G
Sainfoin	F	NS
White prairieclover	F	NS

G - Good adaptation for forage production on this group of soils in this MLRA

F - Fair adaptation but will not produce at its highest potential

NS - Species is not adapted to the site and should not be planted

Production Estimates

Production estimates listed here should only be used for making general management recommendations. On site production information should always be used for making detailed planning and management recommendations.

PASTURE AND HAYLAND INTERPRETATIONS

The high forage production estimates listed below are based on dense, vigorous stands of climatically adapted, superior performing cultivars. They are properly fertilized for high yields, and pest infestations are kept below economic thresholds. Mechanical harvests are managed to maintain stand life by cutting at appropriate stages of maturity and harvest intervals. If grazed, optimum beginning and ending grazing heights are adhered to. Adequate time is allowed for plant recovery before entering winter dormancy under both uses.

The production estimates listed below represent total annual above ground plant production on an air-dry-matter basis. Estimates of hay and grazing yields can be calculated from these numbers by multiplying them by a harvest efficiency. A 70 percent harvest efficiency is commonly used when converting to hay yields. Pasture harvest efficiency is highly dependent on the grazing management system applied, ranging from 25 to 50 percent.

Forage Crop	<u>Dryland</u>		<u>Irrigated</u>	
	Management Intensity		Management Intensity	
	<u>Low</u> (lbs/ac)	<u>High</u> (lbs/ac)	<u>Low</u> (lbs/ac)	<u>High</u> (lbs/ac)
Alfalfa	2600	6600		
Alfalfa/Crested wheatgrass	2300	5400		
Alfalfa/Intermediate wheatgrass	2600	6000	8600	14300
Alfalfa/Pubescent wheatgrass	2600	6000	8600	14300
Alfalfa/Smooth brome grass	2600	6000	8600	17100
Big bluestem	2600	4900		
Crested wheatgrass	1700	3700		
Intermediate wheatgrass	2000	4000	6900	11400
Pubescent wheatgrass	2000	4000	6900	11400
Smooth brome grass	2000	3700	6900	11400
Switchgrass	2600	4900		

Forage Growth Curves

Growth curves estimate the seasonal distribution of growth of the various forage crops. They indicate when the forages may be available for grazing or mechanical harvest.

Growth Curve Number: SD0003
Growth Curve Name: Irrigated Alfalfa
Growth Curve Description: Irrigated Alfalfa, state wide

<u>Percent Production by Month</u>											
<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	5	25	25	20	15	10	0	0	0

Growth Curve Number: SD0004
Growth Curve Name: Cool season grass
Growth Curve Description: Cool season grass, state wide

<u>Percent Production by Month</u>											
<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	10	40	30	10	5	5	0	0	0

Growth Curve Number: SD0005
Growth Curve Name: Warm season grass
Growth Curve Description: Warm season grass, state wide

<u>Percent Production by Month</u>											
<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	0	10	40	35	15	0	0	0	0

Growth Curve Number: SD0002
Growth Curve Name: Alfalfa
Growth Curve Description: MLRA 65, 64, 60A

<u>Percent Production by Month</u>											
<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	5	35	35	15	5	5	0	0	0

Soil Limitations

These soils have few limitations to the production of climatically adapted forage crops. Production potential is high. Flooding is a potential hazard to some of these soils. Also, all of these soils receive additional moisture, so the potential exists for soil compaction from grazing or operating machinery on them when wet.

Management Interpretations

Soils in this group that are subject to flooding can have forage production adversely impacted if it occurs during the spring or growing season. Flooding duration or the time period plants are under water is more important than flooding frequency for the survival of forage crops. If these soils flood it is generally for only a brief time. Exclude livestock and machinery during extended periods of soil wetness to reduce soil compaction. When establishing new stands or renovating stands select species and varieties that can make best use of the additional soil moisture this group receives.

FSG Documentation

Similar FSGs:

FSG ID

G060AY100SD

FSG Narrative

Loamy soils do not receive the additional water and are less productive.

Inventory Data References:

Agriculture Handbook 296-Land Resource Regions and Major Land Resource Areas
Natural Resources Conservation Service (NRCS) National Water and Climate Center data
USDA Plant Hardiness Zone Maps
National Soil Survey Information System (NASIS) for soil surveys in South Dakota, Nebraska, Wyoming, and Montana counties in MLRA 60A
South Dakota, Nebraska, Wyoming, and Montana NRCS Field Office Technical Guides
NRCS National Range and Pasture Handbook
Various Agricultural Research Service, Cooperative Extension Service, and NRCS research trials for plant adaptation and production.

State Correlation:

This site has been correlated with the following states:

MT
NE
SD
WY

Forage Suitability Group Approval:

Original Author: Tim Nordquist

Original Date: 4/17/02

Approval by: Dave Schmidt

Approval Date: