

FORAGE SUITABILITY GROUP Wet

FSG No.: G060AY900SD

Major Land Resource Area: 60A - Pierre Shale Plains and Badlands

Physiographic Features

The soils in this group are found in upland depressions.

	<u>Minimum</u>	<u>Maximum</u>
Elevation (feet):	2600	3300
Slope (percent):	0	1
Flooding:		
Frequency:	None	None
Duration:	None	None
Ponding:		
Depth (inches):	0	3
Frequency:		
Duration:	Long	Long
Runoff Class:	Negligible	Negligible



Climatic Features

This group occurs in a mid-continental climate characterized by wide seasonal temperature and precipitation fluctuations and extremes.

Annual precipitation varies widely from year to year in MLRA 60A. Average annual precipitation for all climate stations listed below is about 15 inches. About 77% of the annual precipitation occurs during the months of April through September. On average there are about 24 days with greater than .1 inches of precipitation during that same time period.

Average annual snowfall ranges from 25 inches at Newell, SD to 45 inches at Oelrichs, SD. Snow cover at depths greater than 1 inch range from 40 days at Newell, SD to 82 days at Colony, WY.

Average July temperatures across the MLRA are about 74 degrees F., and average January temperatures are about 20 degrees F. Recorded temperature extremes in the MLRA during the years 1961 to 1990 are a low of -47 at Redbird, WY, and a high of 114 recorded at Oelrichs, SD. The MLRA lies mostly in USDA Plant Hardiness Zones 4a and 4b.

At Rapid City, SD, the closest station with records, it is cloudy about 139 days a year. Average morning relative humidity in June is about 78% and average afternoon humidity is 49%.

The climate data listed in the tables below represent high and low ranges and averages for the climate stations and dates listed. For additional climate data access the National Water and Climate Center at

<http://www.wcc.nrcs.usda.gov>

	From	To
Freeze-free period (28 deg)(days): (9 years in 10 at least)	118	137
Last Killing Freeze in Spring (28 deg): (1 year in 10 later than)	May 26	May 14
Last Frost in Spring (32 deg): (1 year in 10 later than)	Jun 07	May 26

First Frost in Fall (32 deg): (1 year in 10 earlier than)	From Sep 02	To Sep 13
First Killing Freeze in Fall (28 deg): (1 year in 10 earlier than)	Sep 11	Sep 21
Length of Growing Season (32 deg)(days): (9 years in 10 at least)	96	117
Growing Degree Days (40 deg):	4231	4913
Growing Degree Days (50 deg):	2400	2852
Annual Minimum Temperature:	-30	-20
Mean annual precipitation (inches):	14	17

Monthly precipitation (inches) and temperature (F):

2 years in 10:	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Precip. Less Than	0.11	0.09	0.30	0.65	1.05	1.04	1.06	0.52	0.37	0.38	0.20	0.15
Precip. More Than	0.49	0.74	1.27	2.50	4.02	4.63	2.98	2.22	1.68	1.62	0.89	0.66
Monthly Average:	0.33	0.42	0.83	1.71	2.69	2.78	1.99	1.47	1.24	1.03	0.53	0.41
Temp. Min.	5.3	10.9	20.0	30.6	40.5	49.8	56.3	53.2	41.7	29.9	18.2	6.5
Temp. Max.	34.3	40.5	49.5	61.5	71.8	82.5	91.2	89.8	79.0	65.6	48.3	36.3
Temp. Avg.	19.9	25.3	34.0	45.8	56.0	66.0	73.6	71.5	60.2	48.0	33.5	22.0

Climate Station	Location	From	To
SD0236	Ardmore, SD	1961	1990
SD6054	Newell, SD	1961	1990
SD6212	Oelrichs, SD	1961	1990
SD8911	Wasta, SD	1961	1990
SD9537	Zeona, SD	1961	1990
WY1905	Colony, WY	1961	1990
WY7555	Redbird, WY	1961	1990

Soil Interpretations

This group consists of poorly drained, fine textured soils. Runoff from higher-lying land causes these soils to pond for several days or weeks following heavy rains or snow melt.

Drainage Class:	Poorly drained	To	Poorly drained
Permeability Class: (0 - 40 inches)	Very slow	To	Slow
Frost Action Class:	Low	To	Moderate

	Minimum	Maximum
Depth:	72	
Surface Fragments >3" (% Cover):	0	0
Organic Matter (percent): (surface layer)	2.0	6.0
Electrical Conductivity (mmhos/cm): (0 - 24 inches)	2	8
Sodium Absorption Ratio: (0 - 12 inches)	0	4
Soil Reaction (1:1) Water (pH): (0 - 12 inches)	6.6	9
Available Water Capacity (inches): (0 - 60 inches)	8	9
Calcium Carbonate Equivalent (percent): (0 - 12 inches)	0	9

Adapted Species List

The following forage species are considered adapted to grow on the soils in this group. Additional information concerning plant characteristics of a number of the listed species as well as individual cultivars of many of those species can be accessed on the web at <http://www.plants.usda.gov>

Cool Season Grasses

Creeping foxtail	G
Reed canarygrass	G
Tall wheatgrass	F
Western wheatgrass	F

Warm Season Grasses

Switchgrass	F
-------------	---

Legumes

Alsike clover	F
---------------	---

G - Good adaptation for forage production on this group of soils in this MLRA

F - Fair adaptation but will not produce at its highest potential

Production Estimates

Production estimates listed here should only be used for making general management recommendations. On site production information should always be used for making detailed planning and management recommendations.

The high forage production estimates listed below are based on dense, vigorous stands of climatically adapted, superior performing cultivars. They are properly fertilized for high yields, and pest infestations are kept below economic thresholds. Mechanical harvests are managed to maintain stand life by cutting at appropriate stages of maturity and harvest intervals. If grazed, optimum beginning and ending grazing heights are adhered to. Adequate time is allowed for plant recovery before entering winter dormancy under both uses.

The production estimates listed below represent total annual above ground plant production on an air-dry-matter basis. Estimates of hay and grazing yields can be calculated from these numbers by multiplying them by a harvest efficiency. A 70 percent harvest efficiency is commonly used when converting to hay yields. Pasture harvest efficiency is highly dependent on the grazing management system applied, ranging from 25 to 50 percent.

Forage Crop	Management Intensity	
	<u>Low</u> (lbs/ac)	<u>High</u> (lbs/ac)
Creeping foxtail	2300	4300
Reed canarygrass	3100	5100

Forage Growth Curves

Growth curves estimate the seasonal distribution of growth of the various forage crops. They indicate when the forages may be available for grazing or mechanical harvest.

Growth Curve Number: SD0004

Growth Curve Name: Cool season grass

Growth Curve Description: Cool season grass, state wide

<u>Percent Production by Month</u>											
<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	10	40	30	10	5	5	0	0	0

Growth Curve Number: SD0005
Growth Curve Name: Warm season grass
Growth Curve Description: Warm season grass, state wide

Percent Production by Month											
<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	0	10	40	35	15	0	0	0	0

Soil Limitations

The primary limitation for these soils is wetness, which may severely limit species selection, delay planting and harvesting of forage crops or result in wheeltrack ruts or livestock poach marks from hooves. The result can be soil compaction, injury to plants, poor soil aeration affecting plant growth, and problems with movement of livestock and machinery. The soils in this group are subject to ponding which will adversely impact forage production when it occurs during the growing season. The time period plants are under water and the soil temperature while it occurs are important for the survival of forage crops. Dormant forages are little affected by inundation unless the water turns to ice.

These soils may also be slightly saline.

Management Interpretations

When establishing new stands or renovating older stands select species that are tolerant of poorly drained soils and slight salinity. Exclude livestock and machinery during extended periods of soil wetness to reduce poaching, rutting, and soil compaction.

FSG Documentation

Similar FSGs:

FSG ID

G060AY895SD

FSG Narrative

Saline soils have elevated levels of salinity that are detrimental to plant

Inventory Data References:

Agriculture Handbook 296-Land Resource Regions and Major Land Resource Areas
Natural Resources Conservation Service (NRCS) National Water and Climate Center data
USDA Plant Hardiness Zone Maps
National Soil Survey Information System (NASIS) for soil surveys in South Dakota, Nebraska, Wyoming, and Montana counties in MLRA 60A
South Dakota, Nebraska, Wyoming, and Montana NRCS Field Office Technical Guides
NRCS National Range and Pasture Handbook
Various Agricultural Research Service, Cooperative Extension Service, and NRCS research trials for plant adaptation and production.

State Correlation:

This site has been correlated with the following states:

MT
NE
SD
WY

Forage Suitability Group Approval:

Original Author: Tim Nordquist
Original Date: 4/17/02
Approval by: Dave Schmidt
Approval Date: