


FORAGE SUITABILITY GROUP Clayey Subsoil

FSG No.: G063BY210SD
Major Land Resource Area: 63B - Southern Rolling Pierre Shale Plains



Physiographic Features

The soils in this group are found on upland slopes and terraces.

	<u>Minimum</u>	<u>Maximum</u>
Elevation (feet):	1300	2000
Slope (percent):	0	15
Flooding:		
Frequency:	None	None
Duration:	None	None
Ponding:		
Depth (inches):		
Frequency:	None	None
Duration:	None	None
Runoff Class:	Medium	Very high

Climatic Features

This group occurs in a mid-continental climate characterized by wide seasonal temperature and precipitation fluctuations and extremes.

Annual precipitation varies widely from year to year in MLRA 63B. Average annual precipitation for all climate stations listed below is about 22 inches. About 76 percent of the annual precipitation occurs during the months of April through September. On average there are about 29 days with greater than .1 inches of precipitation during that same time frame. Precipitation is less than needed for optimum forage production and is the single largest factor limiting production from this group on non-irrigated lands.

Average annual snowfall ranges from 19 inches at Creighton, NE to 44 inches at Winner, SD. Snow cover at depths greater than 1 inch range from 4 days at Stephan, SD to 57 days at Winner.

Average July temperatures across the MLRA are about 76 degrees F., and average January temperatures are about 17 degrees F. Recorded temperature extremes in the MLRA during the years 1961 to 1990 are a low of -37 and a high of 114 both recorded at Kennebec, SD. The MLRA lies in USDA Plant Hardiness Zones 4b and 5a.

The climate data listed in the tables below represent high and low ranges and averages for the climate stations and dates listed. For additional climate data access the National Water and Climate Center at <http://www.wcc.nrcs.usda.gov>

	From	To
Freeze-free period (28 deg)(days): (9 years in 10 at least)	128	152
Last Killing Freeze in Spring (28 deg): (1 year in 10 later than)	May 20	May 08
Last Frost in Spring (32 deg): (1 year in 10 later than)	Jun 09	May 17
First Frost in Fall (32 deg): (1 year in 10 earlier than)	Sep 01	Sep 21

First Killing Freeze in Fall (28 deg): (1 year in 10 earlier than)	From Sep 19	To Sep 30
Length of Growing Season (32 deg)(days): (9 years in 10 at least)	92	131
Growing Degree Days (40 deg):	4526	5505
Growing Degree Days (50 deg):	2652	3257
Annual Minimum Temperature:	-25	-15
Mean annual precipitation (inches):	18	25

Monthly precipitation (inches) and temperature (F):

2 years in 10:	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Precip. Less Than	0.06	0.09	0.27	0.66	1.18	1.80	1.24	0.73	0.65	0.55	0.12	0.13
Precip. More Than	0.54	1.24	2.70	3.97	5.70	5.65	4.96	3.94	4.34	2.64	1.49	0.85
Monthly Average:	0.41	0.55	1.56	2.36	3.34	3.54	3.08	2.45	2.13	1.45	0.77	0.56
Temp. Min.	1.3	7.5	18.2	31.1	42.2	52.3	58.2	55.5	44.9	32.8	18.9	6.1
Temp. Max.	32.4	38.6	48.5	62.8	74.0	84.0	91.1	88.9	78.7	66.0	47.7	35.2
Temp. Avg.	18.7	24.4	34.9	48.5	59.6	69.5	75.7	73.5	63.2	51.1	35.2	22.4

<u>Climate Station</u>	<u>Location</u>	<u>From</u>	<u>To</u>
NE1990	Creighton, NE	1961	1990
NE1365	Butte, NE	1961	1990
SD9367	Winner, SD	1961	1990
SD0778	Bonesteel, SD	1961	1990
SD3452	Gregory, SD	1961	1990
SD7992	Stephan, SD	1961	1990
SD4516	Kennebec, SD	1961	1990

Soil Interpretations

The clayey subsoil group consists mostly of well drained, moderately deep to very deep, medium to fine textured soils formed from or overlying materials weathered from shale. Available water capacity ranges from low to high, and permeability is mostly slow and very slow.

Some of these soils have medium surface textures, but all have clayey subsoils.

Drainage Class:	Moderately well drained	To	Well drained
Permeability Class: (0 - 40 inches)	Slow	To	Very slow
Frost Action Class:	Low	To	Moderate

	<u>Minimum</u>	<u>Maximum</u>
Depth:	20	
Surface Fragments >3" (% Cover):	0	3
Organic Matter (percent): (surface layer)	1.0	5.0
Electrical Conductivity (mmhos/cm): (0 - 24 inches)	0	4
Sodium Absorption Ratio: (0 - 12 inches)	0	6
Soil Reaction (1:1) Water (pH): (0 - 12 inches)	5.6	8.4
Available Water Capacity (inches): (0 - 60 inches)	3	11
Calcium Carbonate Equivalent (percent): (0 - 12 inches)	0	23

Soil Component List (Some phases of these soils may also occur in other FSGs)

Boro	Hilmoe	Millboro	Verdigre
Boyd	Labu	Opal	Witten
Dorna	Lakoma	Promise	
Fairlo	Mcclure	Verdel	

Adapted Species List

The following forage species are considered adapted to grow on the soils in this group. Additional information concerning plant characteristics of a number of the listed species as well as individual cultivars of many of those species can be accessed on the web at <http://plants.usda.gov/>

<u>Cool Season Grasses</u>	<u>Symbol</u>	<u>Dry</u>	<u>Irrig</u>	<u>Warm Season Grasses</u>	<u>Symbol</u>	<u>Dry</u>	<u>Irrig</u>
Crested wheatgrass	AGCR	G	NS	Big bluestem	ANGE	G	G
Green needlegrass	NAVI4	G	NS	Indiangrass	SONU2	G	G
Intermediate wheatgrass	THIN6	G	G	Little bluestem	SCSC	G	NS
Meadow brome	BRBI2	G	G	Sideoats grama	BOCU	G	NS
Orchardgrass	DAGL	F	G	Switchgrass	PAVIV	G	G
Pubescent wheatgrass	THIN6	G	G	<u>Legumes</u>			
Russian wildrye	PSJU3	G	NS	Alfalfa	MESA	G	G
Smooth brome	BRINI2	G	G	Birdsfoot trefoil	LOCO6	F	G
Tall wheatgrass	THPO7	F	NS	Canada milkvetch	ASCAC6	F	NS
Western wheatgrass	PASM	G	NS	Cicer milkvetch	ASCI4	G	F
				Illinois bundleflower	DEIL	F	NS
				Purple prairieclover	DAPUP	F	NS
				Red clover	TRPR2	F	G
				White prairieclover	DACAC	F	NS

G - Good adaptation for forage production on this group of soils in this MLRA
 F - Fair adaptation but will not produce at its highest potential
 NS - Species is not adapted to the site and should not be planted

Production Estimates

Production estimates listed here should only be used for making general management recommendations. On site production information should always be used for making detailed planning and management recommendations.

The high forage production estimates listed below are based on dense, vigorous stands of climatically adapted, superior performing cultivars. They are properly fertilized for high yields, and pest infestations are kept below economic thresholds. Mechanical harvests are managed to maintain stand life by cutting at appropriate stages of maturity and harvest intervals. If grazed, optimum beginning and ending grazing heights are adhered to. Adequate time is allowed for plant recovery before entering winter dormancy under both uses.

The production estimates listed below represent total annual above ground plant production on an air-dry-matter basis. Estimates of hay and grazing yields can be calculated from these numbers by multiplying them by a harvest efficiency. A 70 percent harvest efficiency is commonly used when converting to hay yields. Pasture harvest efficiency is highly dependent on the grazing management system applied, ranging from 25 to 50 percent.

Forage Crop	<u>Dryland</u>		<u>Irrigated</u>	
	Management Intensity		Management Intensity	
	<u>High</u> (lbs/ac)	<u>Low</u> (lbs/ac)	<u>High</u> (lbs/ac)	<u>Low</u> (lbs/ac)
Alfalfa	6900	3100		
Alfalfa/Intermediate wheatgrass	6300	2600	14300	8600
Alfalfa/Smooth brome grass	6300	2600	14300	8600
Big bluestem	6300	2900		
Intermediate wheatgrass	5700	2000	11400	6900
Smooth brome grass	5100	2000	11400	6900
Switchgrass	6300	2900		

Forage Growth Curves

Growth curves estimate the seasonal distribution of growth of the various forage crops. They indicate when the forages may be available for grazing or mechanical harvest.

Growth Curve Number: SD0001

Growth Curve Name: Alfalfa

Growth Curve Description: Alfalfa, MLRAs 107, 102B, 63B, 66, 65

<u>Percent Production by Month</u>											
<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	5	30	25	20	15	5	0	0	0

Growth Curve Number: SD0004

Growth Curve Name: Cool season grass

Growth Curve Description: Cool season grass, state wide

<u>Percent Production by Month</u>											
<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	10	40	30	10	5	5	0	0	0

Growth Curve Number: SD0005

Growth Curve Name: Warm season grass

Growth Curve Description: Warm season grass, state wide

<u>Percent Production by Month</u>											
<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	0	10	40	35	15	0	0	0	0

Growth Curve Number: SD0003

Growth Curve Name: Irrigated Alfalfa

Growth Curve Description: Irrigated Alfalfa, state wide

<u>Percent Production by Month</u>											
<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	5	25	25	20	15	10	0	0	0

Soil Limitations

The primary limiting factors to these soils are their tight, slowly permeable nature. Because of their slow water intake runoff is increased causing the soils to be droughty. This will be especially apparent on those soils with lower available water holding capacity. On steeper slopes, water erosion is a potential problem during establishment, when renovating stands, and in thin established stands. Livestock trail erosion is a potential problem in established stands.

Management Interpretations

The impact on yields due to the tight, slowly permeable nature and droughtiness of these soils can be reduced by selecting species adapted to those soil conditions when establishing new stands or renovating stands. Including sod forming grass species in stands, especially on steeper slopes will reduce the potential for sheet and rill erosion. Incorporate erosion control practices during the establishment period. Properly locating facilitating practices such as fences, lanes, and water developments can help control livestock movement, reduce trailing perpendicular to steeper slopes, and evenly distribute grazing pressure.

FSG Documentation

Similar FSGs:

FSG ID

G063BY800S

FSG Narrative

Claypan soils have elevated salinity, sodicity, and/or alkalinity and are less productive.

Inventory Data References:

Agriculture Handbook 296-Land Resource Regions and Major Land Resource Areas
Natural Resources Conservation Service (NRCS) National Water and Climate Center data
USDA Plant Hardiness Zone Maps
National Soil Survey Information System (NASIS) for soil surveys in South Dakota and Nebraska counties in MLRA 63B
South Dakota and Nebraska NRCS Field Office Technical Guides
NRCS National Range and Pasture Handbook
Various South Dakota and Nebraska Agricultural Research Service, Cooperative Extension Service, and NRCS research trials for plant adaptation and production

State Correlation:

This site has been correlated with the following states:

NE
SD

Forage Suitability Group Approval:

Original Author: Tim Nordquist

Original Date: 4/3/02

Approval by: Dave Schmidt

Approval Date: