

FORAGE SUITABILITY GROUP Overflow

FSG No.: G064XY500NE

Major Land Resource Area: 64 - Mixed Sandy and Silty Tableland

Physiographic Features

The soils in this group are generally found in nearly level to gently sloping positions on flood plains, footslopes, and upland swales. They receive beneficial additional moisture as run-on from up slope, or from flooding.



	<u>Minimum</u>	<u>Maximum</u>
Elevation (feet):	2950	3940
Slope (percent):	0	3
Flooding:		
Frequency:	None	Frequent
Duration:	None	Brief
Ponding:		
Depth (inches):		
Frequency:	None	None
Duration:	None	None
Runoff Class:	Negligible	Medium

Climatic Features

This group occurs in a mid-continental climate characterized by wide seasonal temperature and precipitation fluctuations and extremes.

Annual precipitation varies widely from year to year in MLRA 64. Average annual precipitation for all climate stations listed below is about 16 inches. About 79 percent of the annual precipitation occurs during the months of April through September. On average there are about 26 days with greater than .1 inches of precipitation during that same time period. Precipitation in this MLRA is less than needed for optimum forage production and is the single largest factor limiting production from this group on non-irrigated lands.

Average annual snowfall ranges from 20 inches at Interior, SD to 60 inches at Harrison, NE. Snow cover at depths greater than 1 inch range from 28 days at Interior, SD to 60 days at Long Valley, SD.

Average July temperatures across the MLRA are about 74 degrees F., and average January temperatures are about 22 degrees F. Recorded temperature extremes in the MLRA during the years 1961 to 1990 are a low of -45 and a high of 114 both recorded at Porcupine, SD. The MLRA lies mostly in USDA Plant Hardiness Zones 4a and 4b, with a small area of warmer 5a around Alliance, NE.

The climate data listed in the tables below represent high and low ranges and averages for the climate stations and dates listed. For additional climate data access the National Water and Climate Center at <http://www.wcc.nrcs.usda.gov>

	From	To
Freeze-free period (28 deg)(days): (9 years in 10 at least)	111	158
Last Killing Freeze in Spring (28 deg): (1 year in 10 later than)	Jun 01	May 05
Last Frost in Spring (32 deg): (1 year in 10 later than)	Jun 12	May 14
First Frost in Fall (32 deg): (1 year in 10 earlier than)	Sep 06	Sep 19

PASTURE AND HAYLAND INTERPRETATIONS

Page 2

First Killing Freeze in Fall (28 deg): (1 year in 10 earlier than)	From Sep 11	To Sep 28
Length of Growing Season (32 deg)(days): (9 years in 10 at least)	94	135
Growing Degree Days (40 deg):	3867	4974
Growing Degree Days (50 deg):	2128	2913
Annual Minimum Temperature:	-30	-15
Mean annual precipitation (inches):	15	18

Monthly precipitation (inches) and temperature (F):

2 years in 10:	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Precip. Less Than	0.07	0.10	0.32	0.71	1.22	1.19	1.43	0.66	0.34	0.47	0.12	0.15
Precip. More Than	0.37	0.61	1.87	3.10	4.14	5.14	3.84	2.61	2.02	1.57	0.81	0.50
Monthly Average:	0.33	0.38	0.97	1.88	2.89	2.90	2.32	1.50	1.33	0.97	0.47	0.38
Temp. Min.	8.5	13.4	19.6	28.9	38.8	48.6	55.1	52.7	42.0	31.1	19.8	11.0
Temp. Max.	35.1	40.2	49.7	63.0	73.1	83.6	92.0	91.0	79.9	67.9	48.9	37.1
Temp. Avg.	22.4	27.4	35.2	46.2	56.8	66.7	74.2	72.2	61.3	49.4	35.1	24.5

<u>Climate Station</u>	<u>Location</u>	<u>From</u>	<u>To</u>
SD4184	Interior, SD	1961	1990
SD4983	Long Valley, SD	1961	1990
SD6736	Porcupine, SD	1963	1990
NE1575	Chadron, NE	1961	1990
NE0130	Alliance, NE	1961	1990
NE3615	Harrison, NE	1961	1990

Soil Interpretations

This group consists of very deep, moderately well and well drained, moderately coarse to moderately fine textured soils formed from alluvium. Permeability is moderately slow to moderately rapid, and available water capacity is moderate to high.

Drainage Class:	Moderately well drained	To	Well drained
Permeability Class: (0 - 40 inches)	Moderately slow	To	Moderately rapid
Frost Action Class:	Low	To	High

	<u>Minimum</u>	<u>Maximum</u>
Depth:	72	
Surface Fragments >3" (% Cover):	0	3
Organic Matter (percent): (surface layer)	0.5	6.0
Electrical Conductivity (mmhos/cm): (0 - 24 inches)	0	4
Sodium Absorption Ratio: (0 - 12 inches)	0	10
Soil Reaction (1:1) Water (pH): (0 - 12 inches)	6.1	8.4
Available Water Capacity (inches): (0 - 60 inches)	7	12
Calcium Carbonate Equivalent (percent): (0 - 12 inches)	0	10

Soil Component List (Some phases of these soils may also occur in other FSGs)

Craft	Goshen	Mccook	Onita
Duroc	Haverson	Mobridge	
Glenberg	Lohmiller	Munjor	

Adapted Species List

The following forage species are considered adapted to grow on the soils in this group. Additional information concerning plant characteristics of a number of the listed species as well as individual cultivars of many of those species can be accessed on the web at <http://www.plants.usda.gov>

<u>Cool Season Grasses</u>	<u>Symbol</u>	<u>Dry</u>	<u>Irrig</u>	<u>Warm Season Grasses</u>	<u>Symbol</u>	<u>Dry</u>	<u>Irrig</u>
Altai wildrye	LEAN3	G	NS	Big bluestem	ANGE	F	G
Basin wildrye	LECI4	G	NS	Little bluestem	SCSC	G	NS
Crested wheatgrass	AGCR	G	NS	Sideoats grama	BOCU	F	NS
Green needlegrass	NAVI4	G	NS	Switchgrass	PAVIV	F	G
Intermediate wheatgrass	THIN6	G	G	<u>Legumes</u>	<u>Symbol</u>	<u>Dry</u>	<u>Irrig</u>
Meadow brome	BRBI2	F	G	Alfalfa	MESA	G	G
Orchardgrass	DAGL	NS	G	Birdsfoot trefoil	LOCO6	NS	G
Pubescent wheatgrass	THIN6	G	G	Canada milkvetch	ASCAC6	F	NS
Smooth brome	BRINI2	F	G	Cicer milkvetch	ASCI4	G	F
Streambank wheatgrass	ELLAL	F	NS	Purple prairieclover	DAPUP	F	NS
Thickspike wheatgrass	ELMA7	F	NS	Red clover	TRPR2	NS	G
Western wheatgrass	PASM	G	NS	Sainfoin	ONVI	F	NS
				White prairieclover	DACAC	F	NS

G - Good adaptation for forage production on this group of soils in this MLRA

F - Fair adaptation but will not produce at its highest potential

NS - Species is not adapted to the site and should not be planted

Production Estimates

Production estimates listed here should only be used for making general management recommendations. On site production information should always be used for making detailed planning and management recommendations.

The high forage production estimates listed below are based on dense, vigorous stands of climatically adapted, superior performing cultivars. They are properly fertilized for high yields, and pest infestations are kept below economic thresholds. Mechanical harvests are managed to maintain stand life by cutting at appropriate stages of maturity and harvest intervals. If grazed, optimum beginning and ending grazing heights are adhered to. Adequate time is allowed for plant recovery before entering winter dormancy under both uses.

The production estimates listed below represent total annual above ground plant production on an air-dry-matter basis. Estimates of hay and grazing yields can be calculated from these numbers by multiplying them by a harvest efficiency. A 70 percent harvest efficiency is commonly used when converting to hay yields. Pasture harvest efficiency is highly dependent on the grazing management system applied, ranging from 25 to 50 percent.

Forage Crop	<u>Dryland</u>		<u>Irrigated</u>	
	Management Intensity		Management Intensity	
	<u>High</u>	<u>Low</u>	<u>High</u>	<u>Low</u>
	(lbs/ac)	(lbs/ac)	(lbs/ac)	(lbs/ac)
Alfalfa	6600	2600		
Alfalfa/Crested wheatgrass	5400	2300		
Alfalfa/Intermediate wheatgrass	6000	2600	14300	8600
Alfalfa/Pubescent wheatgrass	6000	2600	14300	8600
Alfalfa/Smooth brome grass	6000	2600	17100	8600
Big bluestem	4900	2600		
Crested wheatgrass	3700	1700		
Intermediate wheatgrass	4000	2000	11400	6900
Pubescent wheatgrass	4000	2000	11400	6900
Smooth brome grass	3700	2000	11400	6900
Switchgrass	4900	2600		

Forage Growth Curves

Growth curves estimate the seasonal distribution of growth of the various forage crops. They indicate when the forages may be available for grazing or mechanical harvest.

Growth Curve Number: SD0002

Growth Curve Name: Alfalfa

Growth Curve Description: Alfalfa, MLRAs 65, 64, 60A

<u>Percent Production by Month</u>											
<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	5	35	35	15	5	5	0	0	0

Growth Curve Number: SD0004

Growth Curve Name: Cool season grass

Growth Curve Description: Cool season grass, state wide

<u>Percent Production by Month</u>											
<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	10	40	30	10	5	5	0	0	0

Growth Curve Number: SD0005

Growth Curve Name: Warm season grass

Growth Curve Description: Warm season grass, state wide

<u>Percent Production by Month</u>											
<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	0	10	40	35	15	0	0	0	0

Growth Curve Number: SD0003

Growth Curve Name: Irrigated Alfalfa

Growth Curve Description: Irrigated Alfalfa, state wide

<u>Percent Production by Month</u>											
<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	5	25	25	20	15	10	0	0	0

Soil Limitations

These soils have few limitations to the production of climatically adapted forage crops. Production potential is high relative to most other groups in this MLRA. Flooding is a potential hazard to some of these soils. Also, all of these soils receive additional moisture, so the potential exists for soil compaction from grazing or operating machinery on them when wet.

Management Interpretations

Soils in this group that are subject to flooding can have forage production adversely impacted if it occurs during the spring or growing season. Flooding duration or the time period plants are under water is more important than flooding frequency for the survival of forage crops. If these soils flood it is generally for only a brief time. Exclude livestock and machinery during extended periods of soil wetness to reduce soil compaction. When establishing new stands or renovating stands select species and varieties that can make best use of the additional soil moisture this group receives.

FSG Documentation

Similar FSGs:

FSG ID

G064XY100N

G064XY700N

FSG Narrative

Loamy soils do not receive the additional water and are less productive.

Subirrigated soils have elevated watertables that come closer to the surface during part of the growing season.

Inventory Data References:

Agriculture Handbook 296-Land Resource Regions and Major Land Resource Areas
Natural Resources Conservation Service (NRCS) National Water and Climate Center data
USDA Plant Hardiness Zone Maps
National Soil Survey Information System (NASIS) for soil surveys in South Dakota, Nebraska, and Wyoming counties in MLRA 64
South Dakota and Nebraska and Wyoming NRCS Field Office Technical Guides
NRCS National Range and Pasture Handbook
Various South Dakota and Nebraska Agricultural Research Service, Cooperative Extension Service, and NRCS research trials for plant adaptation and production.

State Correlation:

This site has been correlated with the following states:

NE
SD
WY

Forage Suitability Group Approval:

Original Author: Tim Nordquist
Original Date: 4/15/02
Approval by: Dana Larsen
Approval Date: