

FORAGE SUITABILITY GROUP

Limy Upland

FSG No.: G071XY400NE

Major Land Resource Area: 071X -Central Nebraska Loess Hills

Physiographic Features

These soils are found on upland positions.

	<u>Minimum</u>	<u>Maximum</u>
Elevation (feet):	1640	2600
Slope (percent):	3	30
Flooding:		
Frequency:	None	None
Duration:	None	None
Ponding:		
Depth (inches):		
Frequency:	None	None
Duration:	None	None
Runoff Class:	Low	High

Climatic Features

Average annual precipitation for all climate stations listed below is about 25 inches. About 76 percent of the precipitation received in MLRA 71 falls during the months of April through September. On average there are about 30 days during that period that receive greater than .1 inches. Precipitation is less than needed for optimum forage production and is the single largest factor limiting production from this group on non-irrigated lands.

Average annual snowfall ranges from 14 inches at Greeley, NE, to 36 inches at Loup City, NE. Snow cover at depths greater than 1 inch range from a just 2 days per year at Greeley to a high of 49 days per year at Central City, NE.

Average January temperatures for the listed climate stations during the years 1961 to 1990 are about 23 degree F., and average July temperatures are about 76 degrees. Recorded temperature extremes are a low of -36 at Broken Bow, NE, and a high of 106 at North Platte, NE, which lies just west of the MLRA. The MLRA lies in USDA Plant Hardiness Zones 4b and 5a.

It is cloudy an average of 140 days per year at Grand Island, and 141 days a year at North Platte. Average annual wind speeds are about 11.8 MPH at Grand Island and 10.1 at North Platte. Highest average wind speeds occur during the spring. At Grand Island average morning relative humidity in June is about 82 percent and average afternoon humidity in June is about 55 percent. At North Platte they are 84 and 57 percent respectively.

The climate data listed in the tables below represent high and low ranges and averages for the climate stations and dates listed. For additional climate data access the National Water and Climate Center at <http://www.wcc.nrcs.usda.gov>

	From	To
Freeze-free period (28 deg)(days): (9 years in 10 at least)	129	174
Last Killing Freeze in Spring (28 deg): (1 year in 10 later than)	May 18	Apr 27
Last Frost in Spring (32 deg): (1 year in 10 later than)	May 28	May 08

First Frost in Fall (32 deg): (1 year in 10 earlier than)	From Sep 10	To Sep 26
First Killing Freeze in Fall (28 deg): (1 year in 10 earlier than)	Sep 14	Oct 09
Length of Growing Season (32 deg)(days): (9 years in 10 at least)	113	145
Growing Degree Days (40 deg):	5020	5830
Growing Degree Days (50 deg):	2920	3590
Annual Minimum Temperature:	-25	-15
Mean annual precipitation (inches):	22	26

Monthly precipitation (inches) and temperature (F):

2 years in 10:	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Precip. Less Than	0.18	0.09	0.51	0.66	1.86	2.14	1.81	1.32	0.65	0.36	0.13	0.18
Precip. More Than	0.81	1.13	3.39	3.98	5.38	5.84	4.37	3.90	3.74	2.44	1.92	1.18
Monthly Average:	0.43	0.32	1.75	2.40	3.67	3.99	3.30	2.83	2.50	1.46	0.96	0.65
Temp. Min.	9.2	14.6	22.9	33.8	45.2	54.8	60.8	58.4	47.9	35.2	22.2	12.2
Temp. Max.	36.8	42.6	51.8	64.6	73.9	84.4	89.3	87.2	77.6	66.8	50.1	39.1
Temp. Avg.	22.5	27.9	37.4	50.2	60.5	70.3	75.5	73.1	63.5	52.1	37.1	25.5

<u>Climate Station</u>	<u>Location</u>	<u>From</u>	<u>To</u>
NE1200	Broken Bow, NE	1961	1990
NE6040	North Loup, NE	1961	1990
NE3425	Greeley, NE	1961	1990
NE4985	Loup City, NE	1961	1990
NE3365	Gothengurg, NE	1961	1990
NE1560	Central City, NE	1961	1990
NE4335	Kearney, NE	1961	1990

Soil Interpretations

Drainage Class:	Well drained	To	Well drained
Permeability Class: (0 - 40 inches)	Moderate	To	Moderate
Frost Action Class:	Low	To	Moderate

	<u>Minimum</u>	<u>Maximum</u>
Depth:	72	
Surface Fragments >3" (% Cover):	0	0
Organic Matter (percent): (surface layer)	0.5	2.0
Electrical Conductivity (mmhos/cm): (0 - 24 inches)	0	0
Sodium Absorption Ratio: (0 - 12 inches)	0	0
Soil Reaction (1:1) Water (pH): (0 - 12 inches)	7.4	8.4
Available Water Capacity (inches): (0 - 60 inches)	9	13
Calcium Carbonate Equivalent (percent): (0 - 12 inches)	0	15

Forage Growth Curves

Growth curves estimate the seasonal distribution of growth of the various forage crops. They indicate when the forages may be available for grazing or mechanical harvest.

Growth Curve Number: NE0003
Growth Curve Name: Warm-season grass
Growth Curve Description: Statewide

<u>Percent Production by Month</u>											
<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	0	15	35	30	15	5	0	0	0

Growth Curve Number: NE0005
Growth Curve Name: Alfalfa
Growth Curve Description: MLRAs 73, 72 dryland

<u>Percent Production by Month</u>											
<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	0	35	35	20	10	0	0	0	0

Growth Curve Number: NE0006
Growth Curve Name: Cool-season grass fertilized early
Growth Curve Description: MLRAs 73, 72 dryland

<u>Percent Production by Month</u>											
<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	10	40	35	0	5	10	0	0	0

Soil Limitations

Lime

- The primary limitation to the soils in this group is the high lime content close to the soil surface. This reduces species choices and yield potential.

Management Interpretations

Lime

- When establishing new stands select forage species that are tolerant to the high lime levels inherent to these soils.

Slope

- Safe equipment operation is needed on steeper slopes.

Water erosion

- Include sod forming grass species in new seedings on steeper slopes to reduce sheet and rill erosion. Incorporate erosion control practices during the establishment period. Locate fences, lanes, water developments, and mineral areas to reduce livestock trailing perpendicular to steeper slopes.

FSG Documentation

Similar FSGs:

FSG ID

G071XY100NE

FSG Narrative

Loamy soils do not have as high a lime content near the surface and are more productive.

Inventory Data References:

Agriculture Handbook 296-Land Resource Regions and Major Land Resource Areas
Natural Resources Conservation Service (NRCS) National Water and Climate Center data
USDA Plant Hardiness Zone maps
National Soil Survey Information System (NASIS) database for soil surveys in Nebraska counties in MLRA 71
Nebraska NRCS Field Office Technical Guide
NRCS National Range and Pasture Handbook
Various Agricultural Research Service, Cooperative Extension Service, and NRCS research trials for plant adaptation and production.

State Correlation:

This site has been correlated with the following states:
NE

Forage Suitability Group Approval:

Original Author: Tim Nordquist
Original Date: 7/10/200
Approval by: Dana Larsen
Approval Date: