

FORAGE SUITABILITY GROUP

Loam


FSG No.: G102CY100NE

Major Land Resource Area: 102C -Loess Uplands

Physiographic Features

The soils in this group are mostly located in upland positions, foot slopes, and stream terraces. A few also occur on floodplains.

	<u>Minimum</u>	<u>Maximum</u>
Elevation (feet):	980	1640
Slope (percent):	0	17
Flooding:		
Frequency:	None	Occasional
Duration:	None	Brief
Ponding:		
Depth (inches):	0	0
Frequency:	None	None
Duration:	None	None
Runoff Class:	Very low	High



Climatic Features

Annual precipitation varies widely from year to year in MLRA 102C. Average annual precipitation for all climate stations listed below is about 27 inches. About 73 percent of the annual precipitation occurs during the months of April through September. On average there are about 33 days with greater than .1 inches of precipitation during the same time period. Annual precipitation and temperature increase from the northwest to the southeast in the MLRA. Precipitation is less than needed for optimum forage production and is the single largest factor limiting production from this group on non-irrigated lands.

Average annual snowfall ranges from 19 inches at Creighton, to 36 inches at Wakefield. Days with snow cover at depths greater than 1 inch range from 9 days at Creighton to 55 days at Wakefield.

Average July temperatures are about 76 degrees F., and average January temperatures are about 20 degrees F. Recorded temperature extremes in the MLRA during the years 1961 to 1990 are a low of -36 at Creighton and a high of 108 recorded at both Columbus and Wakefield. The MLRA lies mostly in USDA Plant Hardiness Zone 4b with some small areas of warmer 5a.

At Norfolk, NE, the average annual wind speeds are about 11.2 MPH. The highest wind speeds occur during March and April. It is cloudy about 146 days a year. Average morning relative humidity in June is about 82 percent, and average afternoon humidity is 55 percent.

The climate data listed in the tables below represent high and low ranges and averages for the climate stations and dates listed. For additional climate data access the National Water and Climate Center at <http://www.wcc.nrcs.usda.gov>.

	From	To
Freeze-free period (28 deg)(days): (9 years in 10 at least)	138	168
Last Killing Freeze in Spring (28 deg): (1 year in 10 later than)	May 12	Apr 25
Last Frost in Spring (32 deg): (1 year in 10 later than)	May 20	May 10

PASTURE AND HAYLAND INTERPRETATIONS

Page 2

	From	To
First Frost in Fall (32 deg): (1 year in 10 earlier than)	Sep 12	Sep 23
First Killing Freeze in Fall (28 deg): (1 year in 10 earlier than)	Sep 20	Oct 04
Length of Growing Season (32 deg)(days): (9 years in 10 at least)	126	146
Growing Degree Days (40 deg):	4833	5730
Growing Degree Days (50 deg):	2815	3551
Annual Minimum Temperature:	-25	-15
Mean annual precipitation (inches):	23	30

Monthly precipitation (inches) and temperature (F):

2 years in 10:	<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
Precip. Less Than	0.15	0.17	0.51	0.83	1.92	1.98	1.71	1.35	1.07	0.44	0.15	0.30
Precip. More Than	1.13	1.20	3.39	3.93	5.93	6.63	4.33	5.39	5.53	3.85	2.16	1.47
Monthly Average:	0.55	0.76	2.04	2.53	4.03	4.20	3.09	3.03	3.00	2.02	1.18	0.86
Temp. Min.	5.3	10.6	23.0	35.4	46.5	56.8	61.9	58.7	48.4	35.6	23.8	10.7
Temp. Max.	32.4	38.2	50.2	65.2	75.6	84.8	88.7	86.1	77.7	66.5	49.7	35.3
Temp. Avg.	19.7	25.3	36.8	50.6	61.6	71.3	75.9	73.1	63.9	52.3	37.1	23.5

<u>Climate Station</u>	<u>Location</u>	<u>From</u>	<u>To</u>
NE1825	Columbus, NE	1961	1990
NE1990	Creighton, NE	1961	1990
NE3050	Fremont, NE	1961	1990
NE6018	NE Nebraska Experiment	1964	1990
NE8110	Stanton, NE	1961	1990
NE8480	Tekamah, NE	1961	1990
NE8915	Wakefield, NE	1961	1990
NE8935	Walthill, NE	1961	1990

Soil Interpretations

This group consists mostly of well drained, moderately coarse to moderately fine textured soils formed mostly from loess, alluvium, and colluvium. Available water capacity is high and permeability is moderately slow to moderate.

Drainage Class:	Moderately well drained	To	Well drained
Permeability Class: (0 - 40 inches)	Moderately slow	To	Moderately rapid
Frost Action Class:	Moderate	To	High

	<u>Minimum</u>	<u>Maximum</u>
Depth:	40	
Surface Fragments >3" (% Cover):		
Organic Matter (percent): (surface layer)	0.5	8.0
Electrical Conductivity (mmhos/cm): (0 - 24 inches)	0	4
Sodium Absorption Ratio: (0 - 12 inches)	0	0
Soil Reaction (1:1) Water (pH): (0 - 12 inches)	5.1	8.4
Available Water Capacity (inches): (0 - 60 inches)	9	
Calcium Carbonate Equivalent (percent): (0 - 12 inches)	0	10

Soil Component List (Some phases of these soils may also occur in other FSGs)

Alcester	Longford	Davis variant	Grovena
Belfore	Loretto	Dempster	Houdek
Clarno	Marshall	Dobalt	Ihlen
Crofton	Maskell	Enet	Mcpaul
Eltree	Merrick	Grable	Modale
Eudora	Monona	Henkin	Nora
Geary	Moody	Maddock	Omadi
Geary variant	Muir	Scroll	Salix
Grigston	Nora	Blake	Shindler
Hall	Nora variant	Blyburg	Splitrock
Haynie	Paka	Bonilla	Trent
Hersh	Shell	Davis	Wentworth
Hord	Steinauer	Egan	
Judson	Alwilda	Flandreau	
Leisy	Blendon	Graceville	

Adapted Species List

The following forage species are considered adapted to grow on the soils in this group. Additional information concerning plant characteristics of a number of the listed species as well as individual cultivars of many of those species can be accessed on the web at <http://plants.usda.gov>.

Cool Season Grasses	<u>Symbol</u>	<u>Dryland</u>	<u>Irrig</u>	Warm Season Grasses	<u>Symbol</u>	<u>Dryland</u>	<u>Irrig</u>
Canada wildrye	ELCA4	F	NS	Big bluestem	ANGE	G	G
Green needlegrass	NAVI4	G	NS	Indiangrass	SONU2	G	G
Intermediate wheatgrass	THIN6	G	G	Little bluestem	SCSC	F	NS
Meadow bromegrass	BRBI2	G	G	Prairie sandreed	CALO	F	NS
Orchardgrass	DAGL	G	G	Sand bluestem	ANHA	F	NS
Pubescent wheatgrass	THIN6	G	G	Sideoats grama	BOCU	F	NS
Smooth bromegrass	BRINI2	G	G	Switchgrass	PAVIV	G	G
Tall fescue	LOAR10	F	F	Legumes	<u>Symbol</u>	<u>Dryland</u>	<u>Irrig</u>
Tall wheatgrass	THPO7	G	NS	Alfalfa	MESA	G	G
Virginia wildrye	ELVI3	F	NS	Birdsfoot trefoil	LOCO6	G	G
Western wheatgrass	PASM	F	NS	Canada milkvetch	ASCAC6	F	NS
				Cicer milkvetch	ASCI4	G	F
				Purple prairieclover	DAPUP	F	NS
				Red clover	TRPR2	F	G
				White prairieclover	DACAC	F	NS

G - Good adaptation for forage production on this group of soils in this MLRA

F - Fair adaptation but will not produce at its highest potential

NS - Species is not adapted to the site and should not be planted

Production Estimates

Production estimates listed here should only be used for making general management recommendations. On site production information should always be used for making detailed planning and management recommendations.

The high forage production estimates listed below are based on dense, vigorous stands of climatically adapted, superior performing cultivars. They are properly fertilized for high yields, and pest infestations are kept below economic thresholds. Mechanical harvests are managed to maintain stand life by cutting at appropriate stages of maturity and harvest intervals. If grazed, optimum beginning and ending grazing heights are adhered to. Adequate time is allowed for plant recovery before entering winter dormancy under both uses.

PASTURE AND HAYLAND INTERPRETATIONS

Page 4

The production estimates listed below represent total annual above ground plant production on an air-dry-matter basis. Estimates of hay and grazing yields can be calculated from these numbers by multiplying them by a harvest efficiency. 70 percent harvest efficiency is commonly used when converting to hay yields. Pasture harvest efficiency is highly dependent on the grazing management system applied, ranging from 25 to 50 percent.

Forage Crop	<u>Dryland</u>		<u>Irrigated</u>	
	Management Intensity		Management Intensity	
	<u>Low</u> (lbs/ac)	<u>High</u> (lbs/ac)	<u>Low</u> (lbs/ac)	<u>High</u> (lbs/ac)
Alfalfa	4300	11400		
Alfalfa/Intermediate wheatgrass	4300	10800	10300	17100
Alfalfa/Smooth bromegrass	4300	10800	10300	17100
Big bluestem	4300	8600		
Intermediate wheatgrass	3700	7100	8600	14300
Smooth bromegrass	3700	6600	8600	14300
Switchgrass	4000	8600		

Forage Growth Curves

Growth curves estimate the seasonal distribution of growth of the various forage crops. They indicate when the forages may be available for grazing or mechanical harvest.

Growth Curve Number: SD0001

Growth Curve Name: Alfalfa

Growth Curve Description: Alfalfa, MLRAs 102B, 102C, 63B, 66, 65

<u>Percent Production by Month</u>											
<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	5	30	25	20	15	5	0	0	0

Growth Curve Number: SD0004

Growth Curve Name: Cool season grass

Growth Curve Description: Cool season grass, state wide

<u>Percent Production by Month</u>											
<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	10	40	30	10	5	5	0	0	0

Growth Curve Number: SD0005

Growth Curve Name: Warm season grass

Growth Curve Description: Warm season grass, state wide

<u>Percent Production by Month</u>											
<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	0	10	40	35	15	0	0	0	0

Growth Curve Number: SD0003

Growth Curve Name: Irrigated Alfalfa

Growth Curve Description: Irrigated Alfalfa, state wide

<u>Percent Production by Month</u>											
<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	5	25	25	20	15	10	0	0	0

Soil Limitations

These soils have few limitations to the production of climatically adapted forage crops. On steeper slopes, water erosion is a potential problem during establishment, when renovating stands, and in thin established stands. Livestock trail erosion is a potential problem in established stands.

Management Interpretations

Including sod forming grass species in stands, especially on steeper slopes, will reduce the potential for sheet and rill erosion. Incorporate erosion control practices during the establishment period. Properly locating facilitating practices such as fences, lanes, and water developments can help control livestock movement, reduce trailing perpendicular to steeper slopes, and evenly distribute grazing pressure.

FSG Documentation

Similar FSGs:

FSG ID

G102CY109NE

FSG Narrative

Steep Loam soils are more steeply sloping with greater runoff potential.

G102CY120NE

Droughty Loam soils are shallower or coarser textured resulting in lower available water capacity and lower production potential.

G102CY500NE

Overflow soils receive additional moisture due to a favorable landscape position resulting in a higher production potential.

Inventory Data References:

Agriculture Handbook 296-Land Resource Regions and Major Land Resource Areas
Natural Resources Conservation Service (NRCS) National Water and Climate Center data
USDA Plant Hardiness Zone maps
National Soil Survey Information System (NASIS) for soil surveys in Nebraska and South Dakota counties in MLRA 102C
Nebraska and South Dakota NRCS Field Office Technical Guide
NRCS National Range and Pasture Handbook
Various Agricultural Research Service, Cooperative Extension Service, and NRCS research trials for plant adaptation and production.

State Correlation:

This site has been correlated with the following states:
NE, SD

Forage Suitability Group Approval:

Original Author: Tim Nordquist

Original Date: 8/1/2001

Approval by: Dana Larsen

Approval Date: