

**Scenario Worksheet**

Practice and Scenario Description:	
Information Type	Data
Region	Mid Atlantic
State	New Jersey
Discipline Group	Agronomy
Practice Code/Name	329 - Residue and Tillage Management - No-Till/ Strip Till/ Direct Seed
Scenario ID	1
Scenario Name	No-Till/Strip-Till

Scenario Description	<p>This practice typically involves conversion from a clean-tilled (conventional tilled) system to no-till or strip-till (conservation tilled) system on 100 acres of cropland. This involves managing the amount, orientation and distribution of crop and other plant residue on the soil surface year round while limiting soil-disturbing activities used to grow and harvest crops in systems. The practice is used to reduce sheet and rill erosion, reduce wind erosion improve soil quality, reduce CO2 losses from the soil, reduce energy use, increase plant available moisture and provide food and escape cover for wildlife. The no-till/strip-till system includes chemical weed control (rather than cultivation) and may also include a period of chemical fallow. System is applicable in both irrigated and non-irrigated fields.</p> <p><b>Associated Practices:</b> Conservation Crop Rotation (328), Cover Crop (340), Residue Management Seasonal (344), and Nutrient Management (590).</p>	
Before Practice Situation	<p>Row crops or small grains are grown and harvested. Full width tillage is performed prior to planting and weed control during crop production is typically cultivation and chemical application. Fields are disked immediately following harvest, with additional operations in some fields to facilitate drainage or additional weed control. Residue amounts after tillage operations average 10% or less, resulting in bare soil being exposed to wind erosion and/or intense rainfall during the fall, winter, and early spring. Any crop residue that is present degrades and sediment/nutrient runoff from fields increases during rainfall events. Sheet and rill erosion occurs with visible rills by spring. Soil health (soil organic matter) declines over time as a result of tillage practices, low residue, and long periods of bare soil. This system will typically have a negative Soil Conditioning Index (SCI) and a high Soil Tillage Intensity Rating (STIR).</p>	
After Practice Situation	<p>Managing crop residue on the surface of a field (typical 100 acre) year around according to the 329 practice plan while limiting soil disturbing activities to those which place nutrients, and plant crops that meet the minimum criteria in the 329 practice standard. All crops are seeded/planted with a no-till drill or no-till/strip-till planter, which minimizes soil disturbance while establishing good seed-soil contact. All residues are to be maintained on the soil surface in a uniform distribution over the entire field and not burned or removed. Crop residues provide soil surface cover throughout the year. Runoff and erosion are reduced and no rills are visible on the soil surface. Wind erosion is reduced by standing residues and surface cover. Over time, soil health is improved due to the additional biomass (crop residues), ground cover, and soil infiltration. Crop residues and/or cover crop residues left on the soil surface may maximize weed control by increasing allelopathic and mulching effect, and provides cover for wildlife. The practice would require reducing soil disturbance and erosion and increasing biomass returned to the soil in sufficient amounts to achieve increased SCI and decreased STIR.</p>	
Scenario Feature Measure	Area planted	
Scenario Unit	Acre	
Scenario Typical Size	50	

Cost Summary:		
Cost Category	Scenario Cost	Scenario Cost/Unit
Materials	\$552.00	\$11.04
Equipment/Installation	\$1,070.50	\$21.41
Labor	\$0.00	\$0.00
Mobilization	\$0.00	\$0.00
Acquisition of Technical Knowledge	\$150.27	\$3.01
Foregone Income	\$0.00	\$0.00
<b>Total</b>	<b>\$1,772.77</b>	<b>\$35.46</b>



**Scenario Worksheet**

Practice and Scenario Description:		
Information Type	Data	
Region	Mid Atlantic	
State	New Jersey	
Discipline Group	Agronomy	
Practice Code/Name	329 - Residue and Tillage Management - No-Till/ Strip Till/ Direct Seed	
Scenario ID	2	
Scenario Name	Organic No-Till/Strip-Till	
Scenario Description	<p>This practice typically involves conversion from a clean or mulch tilled (conventional tilled) system to no-till or strip-till (conservation tilled) system on 20 acres of organic cropland. This involves managing the amount, orientation and distribution of crop and other plant residue on the soil surface year round while limiting soil disturbing activities used to grow and harvest crops in systems. The practice is used to reduce sheet and rill erosion, reduce wind erosion, improve soil quality, reduce CO2 losses from the soil, reduce energy use, increase plant available moisture and provide food and escape cover for wildlife. The organic no-till/strip-till system relies on mulching/residue management, organic-approved chemical weed control, or alternative methods of weed control such as hand weeding, flaming, etc. (rather than traditional cultivation). System is applicable in both irrigated and non-irrigated fields.</p> <p>Associated Practices: Conservation Crop Rotation (328), Cover Crop (340), Residue Management Seasonal (344), and Nutrient Management (590).</p>	
Before Practice Situation	<p>Organically grown crops such as various vegetable and fruit crops (along with organically produced row crops) are grown and harvested throughout the year. Full width tillage is performed prior to planting and weed control during crop production is typically cultivation and mulching. Fields are disked immediately following harvest, with additional operations in some fields to facilitate drainage or additional weed control. Residue amounts after tillage operations average 10% or less, resulting in bare soil being exposed to wind erosion and/or intense rainfall during the fall, winter, and early spring. Any crop residue that is present degrades and sediment/nutrient runoff from fields increases during rainfall events. Wind and/or water erosion occurs with visible rills by spring. Soil health (soil organic matter) declines over time as a result of tillage practices, low residue, and long periods of bare soil. This system will typically have a negative Soil Conditioning Index (SCI) and a high Soil Tillage Intensity Rating (STIR).</p>	
After Practice Situation	<p>329 is applied per the practice plan and all the appropriate criteria are followed for the planned purpose(s). Crop residue is managed on the surface of an organic crop field (typical 20 acre) year around while limiting soil disturbing activities to those which condition residue, place nutrients, and plant crops. All crops are seeded/planted with a no-till drill, no-till/strip-till transplanter, or no-till/strip-till planter, which minimize soil disturbance while establishing good seed-soil contact. All residues are to be maintained on the soil surface in a uniform distribution over the entire field and not burned or removed. Crop residues provide soil surface cover throughout the year. Runoff and erosion are reduced and no rills are visible on the soil surface. Wind erosion is reduced by standing residues and surface cover. Over time, soil health is improved due to the additional biomass (crop residues), ground cover, and soil infiltration. Crop residues and/or cover crop residues left on the soil surface may maximize weed control by increasing allelopathic and mulching effect and provide wildlife food and cover. The practice would require reducing soil disturbance and erosion and increasing biomass returned to the soil in sufficient amounts to achieve increased SCI and decreased STIR.</p>	
Scenario Feature Measure	Area planted	
Scenario Unit	Acre	
Scenario Typical Size	20	

Cost Summary:		
Cost Category	Scenario Cost	Scenario Cost/Unit
Materials	\$0.00	\$0.00
Equipment/Installation	\$646.80	\$32.34
Labor	\$0.00	\$0.00
Mobilization	\$0.00	\$0.00
Acquisition of Technical Knowledge	\$150.27	\$7.51
Foregone Income	\$0.00	\$0.00
Total	\$797.07	\$39.85

