
 CAPE MAY COUNTY, NEW JERSEY -- RAINFALL FACTOR: 200

 MINIMUM SLOPES AND SLOPE LENGTHS THAT CLASSIFY AS HIGHLY ERODIBLE
 FOR EACH SOIL PHASE USING FORMULA: $R \times K \times (LS) \geq 8T$

NO HIGHLY ERODIBLE SOILS FOUND IN THIS SOIL PHASE

BP	Berryland sand	0 - 2 %
CU	Coastal beach-Urban land complex	0 - 2 %
DOA	Downer loamy sand	0 - 3 %
DPA	Downer loamy sand watertable	0 - 2 %
DRA	Downer sandy loam	0 - 2 %
FL	Fill land sandy	0 - 2 %
FM	Fill land sandy organic substratum	0 - 2 %
HAA	Hammonton loamy sand	0 - 3 %
HBA	Hammonton sandy loam	0 - 3 %
KMA	Klej loamy sand	0 - 3 %
MU	Muck	0 - 1 %
PS	Pocomoke sandy loam	0 - 2 %
SAA	Sassafras sandy loam	0 - 2 %
SBA	Sassafras sandy loam water table	0 - 2 %
TD	Tidal marsh deep	0 - 2 %
TM	Tidal marsh moderately deep	0 - 2 %
TS	Tidal marsh shallow	0 - 2 %
WMA	Woodstown sandy loam	0 - 2 %

POTENTIALLY HIGHLY ERODIBLE SOILS FOUND IN THIS SOIL PHASE

ARB	Aura sandy loam	0 - 5 %
DRB	Downer sandy loam	2 - 5 %
DSB	Downer sandy loam gravelly substratum	0 - 5 %
EVB	Evesboro sand	0 - 5 %
FRB	Fort Mott sand	0 - 5 %
SAB	Sassafras sandy loam	2 - 5 %