

\*\*\*\*\*  
 CUMBERLAND COUNTY, NEW JERSEY -- RAINFALL FACTOR: 185

\*\*\*\*\*  
 MINIMUM SLOPES AND SLOPE LENGTHS THAT CLASSIFY AS HIGHLY ERODIBLE  
 FOR EACH SOIL PHASE USING FORMULA:  $R \times K \times (LS) \geq 8T$

-----  
 NO HIGHLY ERODIBLE SOILS FOUND IN THIS SOIL PHASE

AC	Atsion sand	0 - 2 %
AGA	Aura gravelly sandy loam	0 - 2 %
BP	Berryland sand	0 - 2 %
CHA	Chillum silt loam	0 - 2 %
DRA	Downer sandy loam	0 - 2 %
EVB	Evesboro sand	0 - 5 %
FD	Fallsington sandy loam	0 - 2 %
HBA	Hammonton sandy loam	0 - 2 %
KMA	Klej loamy sand	0-3 %
LAA	Lakehurst sand	0 - 3 %
LEB	Lakewood sand	0 - 5 %
MOA	Matapeake silt loam	0 - 2 %
MRA	Mattapex silt loam	0 - 2 %
MS	Muck	0 - 1 %
OT	Othello silt loam	0 - 2 %
PS	Pocomoke sandy loam	0 - 2 %
SGA	Sassafras gravelly sandy loam	0 - 2 %
SRA	Sassafras sandy loam	0 - 2 %
TM	Tidal Marsh	0 - 2 %
WMA	Woodstown sandy loam	0 - 2 %

POTENTIALLY HIGHLY ERODIBLE SOILS FOUND IN THIS SOIL PHASE

AGB	Aura gravelly sandy loam	2 - 5 %
AMA	Aura loamy sand	0 - 5 %
ARA	Aura sandy loam	0 - 2 %
ARB	Aura sandy loam	2 - 5 %
CHB	Chillum silt loam	2 - 5 %
DOB	Downer loamy sand	0 - 5 %
DOC	Downer loamy sand	5 - 10 %
DRB	Downer sandy loam	2 - 5 %
EVC	Evesboro sand	5 - 10 %
EVD	Evesboro sand	10 - 20 %
FRA	Fort Mott	0 - 5 %
HAA	Hammonton loamy sand	0 - 5 %
HBB	Hammonton sandy loam	2 - 5 %
MOB	Matapeake silt loam	2 - 5 %
MOC2	Matapeake silt loam eroded	5 - 10 %
MRB	Mattapex silt loam	2 - 5 %
SGB	Sassafras gravelly sandy loam	2 - 5 %
SGC2	Sassafras gravelly sandy loam eroded	5 - 10 %
SRB	Sassafras sandy loam	2 - 5 %
SRC2	Sassafras sandy loam eroded	5 - 10 %
WMB	Woodstown sandy loam	2 - 5 %