
 GLOUCESTER COUNTY, NEW JERSEY -- RAINFALL FACTOR: 185

 MINIMUM SLOPES AND SLOPE LENGTHS THAT CLASSIFY AS HIGHLY ERODIBLE
 FOR EACH SOIL PHASE USING FORMULA: $R \times K \times (LS) \geq 8T$

NO HIGHLY ERODIBLE SOILS FOUND IN THIS SOIL PHASE

AD	Alluvial land	0 - 2 %
BA	Bayboro loam	0 - 2 %
CNA	Collington sandy loam	0 - 2 %
DSA	Downer sandy loam	0 - 2 %
EK	Elkton loam	0 - 2 %
FA	Fallsington loam	0 - 2 %
FD	Fallsington sandy loam	0 - 2 %
FOA	Freehold sandy loam	0 - 2 %
FW	Fresh water marsh	0 - 0 %
LAA	Lakehurst sand	0 - 5 %
LEB	Lakewood sand	0 - 5 %
LO	Leon sand	0 - 2 %
MC	Made land coarse material	0 - 0 %
MF	Made land fine material	0 - 0 %
MU	Muck	0 - 2 %
PA	Pasquotank fine sandy loam	0 - 2 %
PG	Pits	0 - 0 %
PO	Pocomoke loam	0 - 2 %
PS	Pocomoke sandy loam	0 - 2 %
SA	St Johns sand	0 - 2 %
SRA	Sassafras sandy loam	0 - 2 %
TM	Tidal Marsh	0 - 2 %
WNA	WOODSTOWN and Dragston loams	0 - 2 %
WNA	Woodstown and DRAGSTON loams	0 - 2 %
WTB	Woodstown and KLEJ loamy sands	0 - 5 %

POTENTIALLY HIGHLY ERODIBLE SOILS FOUND IN THIS SOIL PHASE

AMB	Aura loamy sand	0 - 5 %
ARB	Aura sandy loam	0 - 5 %
ASB	AURA-Sassafras loamy sands	0 - 5 %
ASB	Aura-SASSAFRAS loamy sands	0 - 5 %
AUB	AURA-Sassafras sandy loams	0 - 5 %
AUB	Aura-SASSAFRAS sandy loams	0 - 5 %
AUC	Aura-SASSAFRAS sandy loams	5 - 10 %
AUC3	Aura-SASSAFRAS sandy loams	severely eroded 5 - 10 %
CK	Colemantown - Matlock loams	0 - 2 %
CMB	Collington loamy sand	0 - 5 %
CMC	Collington loamy sand	5 - 10 %
CNB	Collington sandy loam	2 - 5 %
CNC	Collington sandy loam	5 - 10 %
COB	Colts Neck soils	0 - 5 %
COC	Colts Neck soils	5 - 10 %

POTENTIALLY HIGHLY ERODIBLE SOILS FOUND IN THIS SOIL PHASE
(continued)

DOB	Downer loamy sand	0 - 5 %
DSB	Downer sandy loam	2 - 5 %
EK	Elkton loam	0 - 2 %
FHB	Freehold loamy sand	0 - 5 %
FHC	Freehold loamy sand	5 - 10 %
FNB	Freehold sand thick surface variant	0 - 10 %
FOB	Freehold sandy loam	2 - 5 %
FOC	Freehold sandy loam	5 - 10 %
FOC3	Freehold sandy loam severely eroded	5 - 10 %
KPB	Keyport sandy loam	0 - 5 %
KRB	Kresson sandy loam	0 - 5 %
LDB	Lakeland sand	0 - 10 %
LKA	LENOIR and Keyport loams	0 - 5 %
LKA	Lenoir and KEYPORT loams	0 - 5 %
MRB	Marlton sandy loam	0 - 5 %
NBB	NIXONTON and Barclay soils	0 - 5 %
NBB	Nixonton and BARCLAY soils	0 - 5 %
SFB	Sassafras loamy sand	0 - 5 %
SFC	Sassafras loamy sand	5 - 10 %
SRB	Sassafras sandy loam	2 - 5 %
SRC	Sassafras sandy loam	5 - 10 %
WHB	Westphalia soils	0 - 5 %
WOB	WOODSTOWN and Dragston loamy sands	0 - 5 %
WOB	Woodstown and DRAGSTON loamy sands	0 - 5 %
WSB	WOODSTOWN and Dragston sandy loams	0 - 5 %
WSB	Woodstown and DRAGSTON sandy loams	0 - 5 %
WTB	WOODSTOWN and Klej loamy sands	0 - 5 %

ALL SLOPES IN THIS SOIL PHASE ARE HIGHLY ERODIBLE

ASC	AURA-Sassafras loamy sands	5 - 10 %
ASC	Aura-SASSAFRAS loamy sands	5 - 10 %
AUC	AURA-Sassafras sandy loams	5 - 10 %
AUC3	AURA-Sassafras sandy loams severely eroded	5 - 10 %
FOD3	Freehold sandy loam severely eroded	10 - 15 %
FSD	Freehold soils	10 - 15 %
FTE	Freehold Colts Neck and Collington soils	15 - 25 %
FTF	Freehold Colts Neck and Collington soils	25 - 40 %
KPC3	Keyport sandy loam severely eroded	5 - 10 %
MRC	Marlton sandy loam	5 - 10 %
MRC3	Marlton sandy loam severely eroded	5 - 10 %
MRD	Marlton sandy loam	10 - 15 %
MRD3	Marlton sandy loam severely eroded	10 - 15 %
MRE	Marlton sandy loam	15 - 25 %
MRF	Marlton sandy loam	25 - 40 %
SRD3	Sassafras sandy loam severely eroded	10 - 15 %
SSD	Sassafras soils	10 - 15 %
SSE	Sassafras soils	15 - 40 %

ALL SLOPES IN THIS SOIL PHASE ARE HIGHLY ERODIBLE(continued)

WAD3	Westphalia fine sandy loam severely eroded	10 - 15 %
WHC	Westphalia soils	5 - 10 %
WHD	Westphalia soils	10 - 15 %
WHE	Westphalia soils	15 - 25 %