

# Hamilton

## NUMERICAL INDEX OF SOIL MAP UNITS

<u>Map Symbol</u>	<u>Mapping Unit Name</u>	<u>Page</u>
3 9	Ondawa fine sandy loam	41
3 10	Udifluvents-Fluvaquents complex, frequently flooded 0-3%	50
2 18B	Colton gravelly loamy sand, undulating	30
2 18C	Colton gravelly loamy sand, rolling	30
1 18DE	Colton gravelly loamy sand, hilly	31
3 22	Croghan loamy sand	32
3 23	Naumburg loamy fine sand	40
3 25A	Adams coarse sandy loam, 0 to 3 percent slopes	12
2 25B	Adams coarse sandy loam, 3 to 8 percent slopes	12
2 25C	Adams coarse sandy loam, 8 to 15 percent slopes	13
50	Dumps, landfills	33
2 58	Westbury very bouldery silt loam, 0 to 8 percent slopes	52
2 70B	Berkshire very bouldery sandy loam, 3 to 8 percent slopes	26
2 70C	Berkshire very bouldery sandy loam, 8 to 15 percent slopes	27
1 70DE	Berkshire very bouldery sandy loam, 15 to 35 percent slopes	27
2 75B	Potsdam very bouldery loam, 3 to 8 percent slopes	43
1 75C	Potsdam very bouldery loam, 8 to 15 percent slopes	44
1 75DE	Potsdam very bouldery loam, 15 to 35 percent slopes	45
2 78B	Crary very bouldery loam, 3 to 8 percent slopes	32
2 80B	Becket very bouldery loam, 3 to 8 percent slopes	18
2 80C	Becket very bouldery loam, 8 to 15 percent slopes	19
1 80DE	Becket very bouldery loam, 15 to 35 percent slopes	20
2 81B	Peru very bouldery very fine sandy loam, 3 to 8 percent slopes	41
1 81C	Peru very bouldery very fine sandy loam, 8 to 15 percent slopes	42
3 89	Burnham very bouldery loam	29
2 90BC	Becket very bouldery loam, slightly rocky, 3 to 15 percent slopes	20
1 90DE	Becket very bouldery loam, slightly rocky, 15 to 35 percent slopes	21

	<u>Map Symbol</u>	<u>Mapping Unit Name</u>	<u>Page</u>
2	91BC	Lyman-Becket-Berkshire Variant complex, 3 to 15 percent slopes	34
1	91DE	Lyman-Becket-Berkshire Variant complex, 15 to 35 percent slopes	35
1	92F	Lyman-Berkshire-Rock outcrop complex, 35 to 70 percent slopes	37
2	118BC	Adams-Colton complex, rolling	14
1	118DE	Adams-Colton complex, hilly	15
	178A	Aeric Haplaquods very bouldery fine sandy loam, 0 to 3 percent slopes	16
	178B	Aeric Haplaquods very bouldery fine sandy loam, 3 to 8 percent slopes	17
	185	Borosaprists, level	28
2	190BC	Tunbridge-Canaan loams, 3 to 15 percent slopes	49
1	190DE	Tunbridge-Canaan loams, 15 to 35 percent slopes	50
	193	Borosaprists-Fluvaquents complex, ponded	28
2	291BC	Becket-Lyman-Berkshire Variant complex, very rocky, sloping	24
1	291DE	Becket-Lyman-Berkshire Variant complex, very rocky, moderately steep	25
2	380BC	Becket-Peru complex, sloping <i>8-15</i>	22
1	380DE	Becket association, moderately steep <i>15-25</i>	23
2	391BC	Lyman-Berkshire Variant-Rock outcrop complex, sloping <i>3-15</i>	38
1	391DE	Lyman-Berkshire Variant-Rock outcrop complex, moderately steep <i>15-35</i>	39
	580AB	Aeric Haplaquods-Aquic Haplorthods complex, gently sloping	17
1	591F	Lyman-Berkshire complex, very steep <i>35+</i>	36
1	901BC	Potsdam very bouldery loam, slightly rocky, 3 to 15 percent slopes	45
1	901DE	Potsdam very bouldery loam, slightly rocky. 15 to 35 percent slopes	46
2	911BC	Potsdam-Lyman complex, 3 to 15 percent slopes	47
1	911DE	Potsdam-Lyman complex, 15 to 35 percent slopes	48
	GP	Pits, gravel and sand	43
	ML	Udorthents, smoothed	52