

**NATURAL RESOURCES CONSERVATION SERVICE  
CONSERVATION PRACTICE STANDARD**

**NUTRIENT MANAGEMENT**

(Ac.)

**CODE 590**

**DEFINITION**

Managing the amount (rate), source, placement (method of application), and timing of plant nutrients and soil amendments.

**PURPOSE**

- To budget, supply, and conserve nutrients for plant production.
- To minimize agricultural nonpoint source pollution of surface and groundwater resources.
- To properly utilize manure or organic by-products as a plant nutrient source.
- To protect air quality by reducing odors, nitrogen emissions (ammonia, oxides of nitrogen), and the formation of atmospheric particulates.
- To maintain or improve the physical, chemical, and biological condition of soil.

**CONDITIONS WHERE PRACTICE APPLIES**

This practice applies to all lands where plant nutrients and soil amendments are applied. This standard does not apply to one-time nutrient applications to establish perennial crops.

**CRITERIA**

**General Criteria Applicable to All Purposes**

A nutrient budget for nitrogen (N), phosphorus (P), and potassium (K) must be developed that considers all potential sources of nutrients including, but not limited to, green manures, legumes, crop residues, compost, animal manure, organic by-products, biosolids, waste water, organic matter, soil biological activity, commercial fertilizer, and irrigation water.

All persons (NRCS, non-NRCS personnel,

technical service providers) who review or approve plans for nutrient management or comprehensive nutrient management will be certified in accordance with the Technical Service Providers (TSP proficiency criteria available on the NRCS TechReg Web site, and with Title 180, General Manual (GM), Part 411, Technical Service Provider Assistance Policy.

Enhanced efficiency fertilizers must be defined by the Association of American Plant Food Control Officials (AAPFCO) and be accepted for use by the Oklahoma Department of Agriculture, Food and Forestry, Consumer Protection Services, with responsibility for verification of product guarantees, ingredients (by AAPFCO definition) and label claims.

Sheet, rill and wind erosion must be managed to protect soil and water quality. Concentrated flow erosion (ephemeral and classic gully) must be managed with appropriate conservation practices. When site erosion rates are greater than soil loss tolerance ("T"), a site assessment for nutrient and soil loss must be conducted to determine if mitigation practices are required to protect water quality.

The rate and placement of applied nitrogen and potassium in starter fertilizer shall avoid salt damage.

The NRCS-approved nutrient risk assessment for nitrogen must be completed on all sites. Refer to General Specification for guidance.

The NRCS-approved nutrient risk assessment for phosphorus must be completed when manure or organic by-product amendments are applied. Refer to General Specification for guidance.

On organic operations, the nutrient sources and management must be consistent with the USDA's National Organic Program.

Conservation practice standards are reviewed periodically and updated if needed. To obtain the current version of this standard, contact your Natural Resources Conservation Service [State Office](#) or visit the [Field Office Technical Guide](#).

**NRCS, OK  
August 2012**

Areas contained within minimum application setbacks (e.g., sinkholes, wellheads, gullies, ditches, or surface inlets) must receive nutrients consistent with the setback restrictions.

Applications of irrigation water must minimize the risk of nutrient loss to surface and groundwater.

Nutrient applications associated with irrigation systems shall be applied in accordance with the requirements of Oklahoma NRCS Irrigation Water Management (449) standard and according to the NRCS National Engineering Handbook, Part 652, Irrigation Guide.

Using effluent water for irrigating crops and grasses can increase salt concentrations in the soil creating a negative impact on plant growth. Oklahoma Technical Note Agronomy OK-17 contains guidance for irrigating with effluent water.

Soil pH must be maintained in a range that enhances an adequate level for crop nutrient availability and utilization. Refer to Table 4 in the General Specification for guidance.

#### **Soil, Manure, and Tissue Sampling and Laboratory Analyses (Testing).**

Nutrient planning must be based on current soil, manure, and (where used as supplemental information) tissue test results developed in accordance with Oklahoma State University guidance, or industry practice, if recognized by the university.

Current soil tests are those that are no older than 3 years, unless otherwise required by federal, state or local regulations.

Soil samples shall be taken at least once every three (3) years for analysis or more often if the crop rotation changes, or state law or regulation requires more frequent sampling.

If the area (field) represented by the soil test is extremely variable, the field shall be separated into smaller areas so that representative samples can be gathered across the field. In this way, some areas of the field will be treated differently from others to reduce variability so that the field can be sampled and treated as a unit in the future. Variability in a field can often be noted by differences in slope, soil texture, landscape position, previous crop, manure application history, surface soil color and crop growth or yield.

Where a conservation management unit (CMU) is used as the basis for a sampling unit, all acreage in the CMU must have similar soil type, cropping history, and management practice treatment.

The soil and tissue tests must include analyses pertinent to monitoring or amending the annual nutrient budget, e.g., pH, organic matter, nitrogen, phosphorus, potassium, or other nutrients where applicable. Where salts are a concern, electrical conductivity (EC) and sodicity analysis shall be taken.

Soil test analyses must be performed by laboratories successfully meeting the requirements and performance standards of the North American Proficiency Testing Program-Performance Assessment Program (NAPT-PAP) under the auspices of the Soil Science Society of America (SSSA), Agricultural Laboratory Proficiency Program or other NRCS-approved programs that consider laboratory performance and proficiency to assure accuracy of soil test results.

Nutrient values of manure, organic by-products and biosolids must be determined prior to land application.

Manure analyses must include, at minimum, total nitrogen (N), ammonium N, total phosphorus (P) or  $P_2O_5$ , total potassium (K) or  $K_2O$ , pH, EC and percent solids, or follow Oklahoma State University guidance regarding required analyses.

Manure, organic by-products, and biosolids samples must be collected and analyzed at least annually, or more frequently if needed to account for operational changes (feed management, animal type, manure handling strategy, etc.) impacting manure nutrient concentrations. If no operational changes occur, less frequent manure testing is allowable where operations can document a stable level of nutrient concentrations for the preceding three consecutive years, unless federal, state, or local regulations require more frequent testing.

Samples must be collected, prepared, stored, and shipped, following Oklahoma State University guidance or industry practice. Refer to OSU publication PSS-2248 "Sampling Animal Manure" for a additional guidance.

<http://pods.dasnr.okstate.edu/docushare/dsweb/Get/Document-1348/PSS-2248web.pdf>

When planning for new or modified livestock operations, acceptable “book values” recognized by the NRCS (e.g., NRCS Agricultural Waste Management Field Handbook) and the Oklahoma State University, or analyses from similar operations in the geographical area, may be used if they accurately estimate nutrient output from the proposed operation.

Manure testing analyses must be performed by laboratories successfully meeting the requirements and performance standards of the Manure Testing Laboratory Certification program (MTLCP) under the auspices of the Minnesota Department of Agriculture, or other NRCS-approved program that considers laboratory performance and proficiency to assure accurate manure test results.

### **Nutrient Application Rates.**

Planned nutrient application rates for nitrogen, phosphorus, and potassium shall follow Oklahoma State University guidelines, but cannot exceed the limits established in Table 2 and 3 of the General Specification.

At a minimum, determination of rate must be based on crop/cropping sequence, current soil test results, realistic yield goals, and NRCS-approved nutrient risk assessments.

Realistic yield goals shall be established. A realistic yield goal is generally the average yield over the last 5 years plus 20%. Rates of nutrient application established by the Oklahoma Cooperative Extension Service will be the basis for nutrient recommendations.

For new crops or varieties, industry yield recommendations may be used until documented yield information is available.

Estimates of yield response must consider factors such as poor soil quality, drainage, pH, salinity, etc., prior to assuming that nitrogen and/or phosphorus are deficient.

Lower-than-recommended nutrient application rates are permissible if the grower’s objectives are met.

Applications of biosolids, starter fertilizers, or pop-up fertilizers must be accounted for in the nutrient budget.

### **Nutrient Sources.**

Nutrient sources utilized must be compatible with the application timing, tillage and planting

system, soil properties, crop, crop rotation, soil organic content, and local climate to minimize risk to the environment.

### **Nutrient Application Timing and Placement.**

Timing and placement of all nutrients must correspond as closely as practical with plant nutrient uptake (utilization by crops), and consider nutrient source, cropping system limitations, soil properties, weather conditions, drainage system, soil biology, and nutrient risk assessment results.

Nutrients must not be surface-applied if nutrient losses offsite are likely. This precludes spreading on:

- frozen and/or snow-covered soils, and
- when the top 2 inches of soil are saturated from rainfall or snow melt.

### **Additional Criteria to Minimize Agricultural Nonpoint Source Pollution of Surface and Groundwater**

Planners must use the current NRCS-approved nitrogen, phosphorus, and soil erosion risk assessment tools to assess the risk of nutrient and soil loss. Identified resource concerns must be addressed to meet current planning criteria (quality criteria).

When there is a high risk of transport of nutrients, conservation practices must be coordinated to avoid, control, or trap manure and nutrients before they can leave the field by surface or subsurface drainage (e.g., tile). The number of applications and the application rates must also be considered to limit the transport of nutrients to tile.

Nutrients must be applied with the right placement, in the right amount, at the right time, and from the right source to minimize nutrient losses to surface and groundwater. The following nutrient use efficiency strategies or technologies must be considered:

- slow and controlled release fertilizers
- nitrification and urease inhibitors
- enhanced efficiency fertilizers
- incorporation or injection
- timing and number of applications
- soil nitrate and organic N testing

- coordinate nutrient applications with optimum crop nutrient uptake
- tissue testing, chlorophyll meters, and spectral analysis technologies
- Sensor based nitrogen management techniques
- Split nitrogen application
- other land-grant university recommended technologies that improve nutrient use efficiency and minimize surface or groundwater resource concerns.

**Additional Criteria Applicable to Properly Utilize Manure or Organic By-Products as a Plant Nutrient Source**

Soil salinity testing to monitor salt accumulation in the soil is recommended when large quantities of manure or organic by-products are being applied annually. This salinity analysis can be done in conjunction with routine soil testing every 3 years.

A soil salinity analysis shall be performed when waste water effluent used for irrigation has been applied on an area for 3 years or more. The area shall be monitored for salinity accumulation annually using a soil salinity analysis as long as effluent is being applied. This analysis should include results for Na, Ca, Mg, K, B, EC, TSS (total soluble salts), Sodium Adsorption Ratio (SAR), Exchangeable Sodium Percentage (ESP), and pH.

The total single application of liquid manure:

- must not exceed the soil's infiltration or water holding capacity
- be based on crop rooting depth
- must be adjusted to avoid runoff or loss to subsurface tile drains.

Crop production activities and nutrient use efficiency technologies must be coordinated to take advantage of mineralized plant-available nitrogen to minimize the potential for nitrogen losses due to denitrification or ammonia volatilization.

Nitrogen and phosphorus application rates must be planned based on risk assessment results as determined by NRCS-approved nitrogen and phosphorus risk assessment tools.

Manure or organic by-products may be applied on legumes at rates equal to the estimated

removal of nitrogen in harvested plant biomass, not to exceed recommendations found in Table 6 of the General Specification.

Manure may be applied at a rate equal to the recommended phosphorus application, or estimated phosphorus removal in harvested plant biomass for the crop rotation, or multiple years in the crop sequence at one time. When such applications are made, the application rate must not exceed the acceptable phosphorus risk assessment criteria, must not exceed the recommended nitrogen application rate during the year of application or harvest cycle, and no additional phosphorus must be applied in the current year and any additional years for which the single application of phosphorus is supplying nutrients.

**Additional Criteria to Protect Air Quality by Reducing Odors, Nitrogen Emissions and the Formation of Atmospheric Particulates**

To address air quality concerns caused by odor, nitrogen, sulfur, and/or particulate emissions; the source, timing, amount, and placement of nutrients must be adjusted to minimize the negative impact of these emissions on the environment and human health. One or more of the following may be used:

- slow or controlled release fertilizers
- nitrification inhibitors
- urease inhibitors
- nutrient enhancement technologies
- incorporation
- injection
- stabilized nitrogen fertilizers
- residue and tillage management
- no-till or strip-till
- windbreaks
- other technologies that minimize the impact of these emissions

Do not apply poultry litter, manure, or organic by-products of similar dryness/density when there is a high probability that wind will blow the material offsite.

When applying liquid forms of manure with irrigation equipment, select application conditions when there is high humidity, little/no

**NRCS, OK**

**August 2012**

wind, and/or other conditions that will minimize volatilization losses into the atmosphere.

**Additional Criteria to Improve or Maintain the Physical, Chemical, and Biological Condition of the Soil to Enhance Soil Quality for Crop Production and Environmental Protection**

Manure or organic by-products incorporated into the soil will improve soil structure.

Incorporate surface applied solid/semi-solid manure or other organic by-products, where tillage is possible, into the soil within 72 hours of application to minimize nutrient losses. Avoid applying materials that will not decompose in the soil.

Time the application of nutrients to avoid periods when field activities will result in soil compaction.

In areas where salinity is a concern, select nutrient sources that minimize the buildup of soil salts.

**CONSIDERATIONS**

Use no-till/strip-till in combination with cover crops to sequester nutrients, increase soil organic matter, increase aggregate stability, reduce compaction, improve infiltration, and enhance soil biological activity to improve nutrient use efficiency.

Use nutrient management strategies such as cover crops, crop rotations, and crop rotations with perennials to improve nutrient cycling and reduce energy inputs.

Use variable-rate nitrogen application based on expected crop yields, soil variability, soil nitrate or organic N supply levels, or chlorophyll concentration.

Use variable-rate nitrogen, phosphorus, and potassium application rates based on site-specific variability in crop yield, soil characteristics, soil test values, and other soil productivity factors.

Develop site-specific yield maps using a yield monitoring system. Use the data to further diagnose low- and high- yield areas, or zones, and make the necessary management changes. See Title 190, Agronomy Technical Note (TN) 190.AGR.3, Precision Nutrient Management Planning.

Use manure management conservation practices to manage manure nutrients to limit losses prior to nutrient utilization.

Apply manure at a rate that will result in an "improving" Soil Conditioning Index (SCI) without exceeding acceptable risk of nitrogen or phosphorus loss.

Use legume crops and cover crops to provide nitrogen through biological fixation and nutrient recycling.

Modify animal feed diets to reduce the nutrient content of manure following guidance contained in the Oklahoma NRCS Feed Management (592) standard.

Soil test information should be no older than 1 year when developing new plans.

Excessive levels of some nutrients can cause induced deficiencies of other nutrients, e.g., high soil test phosphorus levels can result in zinc deficiency in corn.

Use soil tests, plant tissue analyses, and field observations to check for secondary plant nutrient deficiencies or toxicity that may impact plant growth or availability of the primary nutrients.

Use the adaptive nutrient management learning process to improve nutrient use efficiency on farms as outlined in the NRCS' National Nutrient Policy in GM 190, Part 402, Nutrient Management.

Potassium should not be applied in situations where an excess (greater than soil test potassium recommendation) causes nutrient imbalances in crops or forages.

Workers should be protected from and avoid unnecessary contact with plant nutrient sources. Extra caution must be taken when handling anhydrous ammonia or when dealing with organic wastes stored in unventilated enclosures.

Material generated from cleaning nutrient application equipment should be utilized in an environmentally safe manner. Excess material should be collected and stored or field applied in an appropriate manner.

Nutrient containers should be recycled in compliance with State and local guidelines or regulations.

Considerations to Minimize Agricultural Nonpoint Source Pollution of Surface and Groundwater.

Use conservation practices that slow runoff, reduce erosion, and increase infiltration, e.g.,

filter strip, contour farming, or contour buffer strips. These practices can also reduce the loss of nitrates or soluble phosphorus.

Use application methods and timing strategies that reduce the risk of nutrient transport by ground and surface waters, such as:

- split applications of nitrogen to deliver nutrients during periods of maximum crop utilization,
- banded applications of nitrogen and/or phosphorus to improve nutrient availability,
- drainage water management to reduce nutrient discharge through drainage systems, and
- incorporation of surface-applied manures or organic by-products if precipitation capable of producing runoff or erosion is forecast within the time of planned application.

Use bioreactors and multistage drainage strategies when approved by the land-grant university.

#### **Considerations to Protect Air Quality by Reducing Nitrogen and/or Particulate Emissions to the Atmosphere.**

Avoid applying manure and other by-products upwind of inhabited areas.

Use high-efficiency irrigation technologies (e.g., reduced-pressure drop nozzles for center pivots) to reduce the potential for nutrient losses.

#### **PLANS AND SPECIFICATIONS**

The following components must be included in the nutrient management plan:

- aerial site photograph(s)/imagery or site map(s), and a soil survey map of the site,
- soil information including: soil type surface texture, pH, drainage class, permeability, available water capacity, depth to water table, restrictive features, and flooding and/or ponding frequency,
- location of designated sensitive areas and the associated nutrient application restrictions and setbacks,
- for manure applications, location of nearby residences, or other locations where humans may be present on a regular basis, and any identified meteorological (e.g., prevailing winds at different times of the

year), or topographical influences that may affect the transport of odors to those locations,

- results of approved risk assessment tools for nitrogen, phosphorus, and erosion losses,
- documentation establishing that the application site presents low risk for phosphorus transport to local water when phosphorus is applied in excess of crop requirement.
- current and/or planned plant production sequence or crop rotation,
- soil, water, compost, manure, organic by-product, and plant tissue sample analyses applicable to the plan,
- when soil phosphorus levels are increasing, include a discussion of the risk associated with phosphorus accumulation and a proposed phosphorus draw-down strategy,
- realistic yield goals for the crops,
- complete nutrient budget for nitrogen, phosphorus, and potassium for the plant production sequence or crop rotation,
- listing and quantification of all nutrient sources and form,
- all enhanced efficiency fertilizer products that are planned for use,
- in accordance with the nitrogen and phosphorus risk assessment tool(s), specify the recommended nutrient application source, timing, amount (except for precision/variable rate applications specify method used to determine rate), and placement of plant nutrients for each field or management unit, and
- guidance for implementation, operation and maintenance, and recordkeeping.

In addition, the following components must be included in a precision/variable rate nutrient management plan:

- Document the geo-referenced field boundary and data collected that was processed and analyzed as a GIS layer or layers to generate nutrient or soil amendment recommendations.
- Document the nutrient recommendation guidance and recommendation equations

**NRCS, OK**

**August 2012**

used to convert the GIS base data layer or layers to a nutrient source material recommendation GIS layer or layers.

- Document if a variable rate nutrient or soil amendment application was made.
- Provide application records per management zone or as applied map within individual field boundaries (or electronic records) documenting source, timing, method, and rate of all applications that resulted from use of the precision agriculture process for nutrient or soil amendment applications.
- Maintain the electronic records of the GIS data layers and nutrient applications for at least 5 years.

If increases in soil phosphorus levels are expected (i.e., when N-based rates are used), the nutrient management plan must document:

- the soil phosphorus levels at which it is desirable to convert to phosphorus based planning,
- the potential plan for soil test phosphorus drawdown from the production and harvesting of crops, and
- management activities or techniques used to reduce the potential for phosphorus transport and loss,
- for AFOs, a quantification of manure produced in excess of crop nutrient requirements, and
- a long-term strategy and proposed implementation timeline for reducing soil P to levels that protect water quality,

## **OPERATION AND MAINTENANCE**

Conduct periodic plan reviews to determine if adjustments or modifications to the plan are needed. At a minimum, plans must be reviewed and revised, as needed with each soil test cycle, changes in manure volume or analysis, crops, or crop management.

Fields receiving animal manures and/or biosolids must be monitored for the accumulation of heavy metals and phosphorus in accordance with Oklahoma State University guidance and State law.

Significant changes in animal numbers, management, and feed management will necessitate additional manure analyses to establish a revised average nutrient content.

Calibrate application equipment to ensure accurate distribution of material at planned rates.

Document the nutrient application rate. When the applied rate differs from the planned rate, provide appropriate documentation for the change.

Records must be maintained for at least 5 years to document plan implementation and maintenance. As applicable, records include:

- soil, plant tissue, water, manure, and organic by-product analyses resulting in recommendations for nutrient application,
- quantities, analyses and sources of nutrients applied,
- dates, and method(s) of nutrient applications, source of nutrients, and rates of application,
- weather conditions and soil moisture at the time of application; lapsed time to manure incorporation; rainfall or irrigation event,
- crops planted, planting and harvest dates, yields, nutrient analyses of harvested biomass, and crop residues removed,
- dates of plan review, name of reviewer, and recommended changes resulting from the review, and
- all enhanced efficiency fertilizer products used.

Additional records for precision/variable rate sites must include:

- maps identifying the variable application source, timing, amount, and placement of all plant nutrients applied, and
- GPS-based yield maps for crops where yields can be digitally collected.

## **REFERENCES**

Association of American Plant Food Control Officials (AAPFCO). 2011. AAPFCO Official Publication no. 64. AAPFCO Inc., Little Rock, AR.

- Follett, R.F. 2001. Nitrogen transformation and transport processes. *In* Nitrogen in the environment; sources, problems, and solutions, (eds.) R.F. Follett and J. Hatfield, pp. 17-44. Elsevier Science Publishers. The Netherlands. 520 pp.
- Schepers, J.S., and W.R. Ruan, (eds.) 2008. Nitrogen in agricultural systems. Agron. Monogr. no. 49, American Society of Agronomy (ASA), Crop Science Society of America (CSSA), Soil Science Society of America (SSSA). Madison, WI.
- Sims, J.T. (ed.) 2005. Phosphorus: Agriculture and the environment. Agron. Monogr. no. 46. ASA, CSSA, and SSSA, Madison, WI.
- Stevenson, F.J. (ed.) 1982. Nitrogen in agricultural soils. Agron. Series 22. ASA, CSSA, and SSSA, Madison, WI.
- U.S. Department of Agriculture, Natural Resources Conservation Service. 2010. Agronomy Technical Note, (TN) 190-AGR-3, Precision Nutrient Management Planning. Washington, DC.
- U.S. Department of Agriculture, Natural Resources Conservation Service. 2011. Title 190, General Manual, (GM), Part 402, Nutrient Management. Washington, DC.
- U.S. Department of Agriculture, Natural Resources Conservation Service. 2011, Title 190, National Instruction (NI), Part 302, Nutrient Management Policy Implementation. Washington, DC.
- OSU Cooperative Extension Service Publications: PSS-2207 "How to Get a Good Soil Sample", PSS-2225 "OSU Soil Interpretations,