

Woody Plants for Use in Pollinator-Friendly Plantings

(applies to the following NRCS practices: 327, 380, 386, 391, 422, 612, 650, ANM14, PLT01, PLT06, PLT08, PLT15)

TREES

| | | |
|-----------------|---|--|
| Dry site | Basswood / American Linden (<i>Tilia americana</i>) | |
| | Cherry, Black (<i>Prunus serotina</i>) | |
| | *Locust, Black (<i>Robinia psuedoacacia</i>) | |
| | Mulberry, Red (<i>Morus rubra</i>) | *may serve as host plants for stink bugs |
| | Poplar, Tulip / Yellow (<i>Liriodendron tulipifera</i>) | |
| | *Redbud, Eastern (<i>Cercis canadensis</i>) | |
| | Sassafras (<i>Sassafras albidum</i>) | |
| (wet or dry) | Maple, Red (<i>Acer rubrum</i>) | |
| Wet site | Willow, Black (<i>Salix nigra</i>) | |

SHRUBS

| | |
|-----------------|---|
| Dry site | Blueberry, Lowbush (<i>Vaccinium angustifolium</i>) |
| | Huckleberry, Black (<i>Gaylussacia baccata</i>) |
| | Chokeberry, Black (<i>Aronia melanocarpa</i>) |
| | Rose, Pasture (<i>Rosa carolina</i>) |
| | Serviceberry, Allegheny (<i>Amelanchier laevis</i>) |
| | Spicebush (<i>Lindera benzoin</i>) |
| | Viburnum, Mapleleaf (<i>Viburnum acerifolium</i>) |
| | Viburnum, Blackhaw (<i>Viburnum prunifolium</i>) |
| Wet site | Blueberry, Highbush (<i>Vaccinium corymbosum</i>) |
| | Buttonbush (<i>Cephalanthus occidentalis</i>) |
| | Chokeberry, Red (<i>Aronia arbutifolia</i>) |
| | Dogwood, Alternate (<i>Cornus alternifolia</i>) |
| | Dogwood, Redosier (<i>Cornus stolonifera</i>) |
| | Dogwood, Silky (<i>Cornus amomum</i>) |
| | Elderberry (<i>Sambucus canadensis</i>) |
| | Rose, Swamp (<i>Rosa palustris</i>) |
| | Viburnum, American Cranberrybush (<i>Viburnum opulus</i>) |
| | Viburnum, Arrowwood (<i>Viburnum dentatum</i>) |