

Field Office Technical Guide Soils Information (SC)

Marlboro County, South Carolina

| Mapunit Symbol and Soil Name | % of Unit | Slope % | Important Farmland Class | Highly Erodible Land* | Drainage Class | Hydric Rating | Hydrologic Soil Group | Land Capability Class | T factor* | Kf* |
|---|-----------|---------|----------------------------------|-------------------------------------|------------------------------|---------------|-----------------------|-----------------------|-----------|-----|
| <i>AaB Ailey sand, moderately wet, 0 to 6 percent slopes</i> | | | | 137 Carolina and Georgia Sand Hills | | | | | | |
| Ailey | 90 | 2-6 | Farmland of statewide importance | Not highly erodible land | Well drained | No | B | 3s | 4 | .10 |
| <i>AeC Ailey loamy sand, 6 to 10 percent slopes</i> | | | | 137 Carolina and Georgia Sand Hills | | | | | | |
| Ailey | 90 | 6-10 | Not prime farmland | Highly erodible land | Well drained | No | B | 6s | 4 | .10 |
| <i>AeD Ailey loamy sand, 10 to 15 percent slopes</i> | | | | 137 Carolina and Georgia Sand Hills | | | | | | |
| Ailey | 90 | 10-15 | Not prime farmland | Highly erodible land | Well drained | No | B | 6s | 4 | .10 |
| <i>AgD Ailey gravelly loamy sand, 6 to 15 percent slopes</i> | | | | 137 Carolina and Georgia Sand Hills | | | | | | |
| Ailey | 90 | 8-15 | Not prime farmland | Highly erodible land | Well drained | No | B | 6s | 4 | .15 |
| <i>AgE Ailey gravelly loamy sand, 15 to 25 percent slopes</i> | | | | 137 Carolina and Georgia Sand Hills | | | | | | |
| Ailey | 90 | 15-25 | Not prime farmland | Highly erodible land | Well drained | No | B | 7e | 4 | .15 |
| <i>AhB Alaga sand, 0 to 6 percent slopes</i> | | | | 133A Southern Coastal Plain | | | | | | |
| Alaga | 85 | 0-6 | Not prime farmland | Not highly erodible land | Somewhat excessively drained | No | A | 3s | 5 | .10 |
| <i>ApB Alpin sand, 0 to 6 percent slopes</i> | | | | 137 Carolina and Georgia Sand Hills | | | | | | |
| Alpin | 90 | 0-6 | Not prime farmland | Not highly erodible land | Excessively drained | No | A | 4s | 5 | .10 |
| <i>ApC Alpin sand, 6 to 10 percent slopes</i> | | | | 137 Carolina and Georgia Sand Hills | | | | | | |
| Alpin | 90 | 6-10 | Not prime farmland | Not highly erodible land | Excessively drained | No | A | 6s | 5 | .10 |
| <i>AuB Autryville sand, 0 to 6 percent slopes</i> | | | | 133A Southern Coastal Plain | | | | | | |
| Autryville | 90 | 0-6 | Farmland of statewide importance | Not highly erodible land | Well drained | No | A | 2s | 5 | .10 |
| <i>BaD Badin silt loam, 2 to 15 percent slopes</i> | | | | 136 Southern Piedmont | | | | | | |
| Badin | 95 | 2-15 | Not prime farmland | Highly erodible land | Well drained | No | B | 4e | 3 | .37 |
| <i>BaF Badin silt loam, 15 to 40 percent slopes</i> | | | | 136 Southern Piedmont | | | | | | |
| Badin | 95 | 15-40 | Not prime farmland | Highly erodible land | Well drained | No | B | 7e | 3 | .37 |
| <i>BnB Blanton sand, 0 to 6 percent slopes</i> | | | | 133A Southern Coastal Plain | | | | | | |
| Blanton | 90 | 0-6 | Not prime farmland | Not highly erodible land | Moderately well drained | No | A | 3s | 5 | .10 |

Field Office Technical Guide Soils Information (SC)

Marlboro County, South Carolina

| Mapunit Symbol and Soil Name | % of Unit | Slope % | Important Farmland Class | Highly Erodible Land* | Drainage Class | Hydric Rating | Hydrologic Soil Group | Land Capability Class | T factor* | Kf* |
|---|-----------|---------|----------------------------------|--|------------------------------|---------------|-----------------------|-----------------------|-----------|-----|
| <i>BnC Blanton sand, 6 to 10 percent slopes</i> | | | | <i>133A Southern Coastal Plain</i> | | | | | | |
| Blanton | 90 | 6-10 | Not prime farmland | Not highly erodible land | Moderately well drained | No | A | 4s | 5 | .10 |
| <i>BoB Bonneau sand, 0 to 4 percent slopes</i> | | | | <i>133A Southern Coastal Plain</i> | | | | | | |
| Bonneau | 90 | 0-4 | Farmland of statewide importance | Not highly erodible land | Somewhat excessively drained | No | A | 2s | 5 | .10 |
| <i>By Byars loam</i> | | | | <i>133A Southern Coastal Plain</i> | | | | | | |
| Byars | 95 | 0-2 | Farmland of statewide importance | Not highly erodible land | Very poorly drained | Yes | D | 3w | 5 | .28 |
| <i>CaB Candor sand, 0 to 6 percent slopes</i> | | | | <i>137 Carolina and Georgia Sand Hills</i> | | | | | | |
| Candor | 90 | 0-6 | Farmland of statewide importance | Not highly erodible land | Somewhat excessively drained | No | A | 3s | 5 | .10 |
| <i>CaC Candor sand, 6 to 10 percent slopes</i> | | | | <i>137 Carolina and Georgia Sand Hills</i> | | | | | | |
| Candor | 90 | 6-10 | Farmland of statewide importance | Not highly erodible land | Somewhat excessively drained | No | A | 4s | 5 | .10 |
| <i>CaD Candor sand, 10 to 15 percent slopes</i> | | | | <i>137 Carolina and Georgia Sand Hills</i> | | | | | | |
| Candor | 90 | 10-15 | Not prime farmland | Highly erodible land | Somewhat excessively drained | No | A | 4s | 5 | .10 |
| <i>CdB Candor sand, moderately wet, 0 to 6 percent slopes</i> | | | | <i>133A Southern Coastal Plain</i> | | | | | | |
| Candor | 95 | 0-6 | Farmland of statewide importance | Not highly erodible land | Somewhat excessively drained | No | A | 3s | 5 | .10 |
| <i>Ce Chastain-Chenneby complex, frequently flooded</i> | | | | <i>133A Southern Coastal Plain</i> | | | | | | |
| Chastain | 65 | 0-2 | Not prime farmland | Not highly erodible land | Poorly drained | Yes | D | 7w | 4 | .32 |
| Chenneby | 25 | 0-2 | Not prime farmland | Not highly erodible land | Somewhat poorly drained | Yes | C | 4w | 5 | .37 |
| <i>Ch Chenneby silt loam, occasionally flooded</i> | | | | <i>133A Southern Coastal Plain</i> | | | | | | |
| Chenneby | 85 | 0-2 | Not prime farmland | Not highly erodible land | Somewhat poorly drained | Yes | C | 2w | 5 | .37 |
| <i>CkA Chewacla loam, 0 to 2 percent slopes, frequently flooded</i> | | | | <i>133A Southern Coastal Plain</i> | | | | | | |
| Chewacla | 90 | 0-2 | Farmland of statewide importance | Not highly erodible land | Somewhat poorly drained | No | C | 4w | 5 | .28 |
| <i>CmA Clayham loam, 0 to 2 percent slopes</i> | | | | <i>133A Southern Coastal Plain</i> | | | | | | |
| Clayham | 95 | 0-2 | All areas are prime farmland | Not highly erodible land | Well drained | No | B | 1 | 5 | .32 |
| <i>CmB Clayham loam, 2 to 6 percent slopes</i> | | | | <i>133A Southern Coastal Plain</i> | | | | | | |
| Clayham | 95 | 2-6 | All areas are prime farmland | Highly erodible land | Well drained | No | B | 2e | 5 | .32 |

Field Office Technical Guide Soils Information (SC)

Marlboro County, South Carolina

| Mapunit Symbol and Soil Name | % of Unit | Slope % | Important Farmland Class | Highly Erodible Land* | Drainage Class | Hydric Rating | Hydrologic Soil Group | Land Capability Class | T factor* | Kf* |
|---|-----------|---------|----------------------------------|-------------------------------------|-------------------------|---------------|-----------------------|-----------------------|-----------|-----|
| <i>CnB2 Clayham clay loam, 2 to 6 percent slopes, eroded</i> | | | | 133A Southern Coastal Plain | | | | | | |
| Clayham | 95 | 2-6 | Farmland of statewide importance | Highly erodible land | Well drained | No | B | 2e | 5 | .32 |
| <i>CoC Cowarts loamy sand, 6 to 10 percent slopes</i> | | | | 137 Carolina and Georgia Sand Hills | | | | | | |
| Cowarts | 96 | 6-10 | Not prime farmland | Highly erodible land | Moderately well drained | No | C | 4e | 4 | .15 |
| <i>CoD Cowarts loamy sand, 10 to 15 percent slopes</i> | | | | 137 Carolina and Georgia Sand Hills | | | | | | |
| Cowarts | 90 | 10-15 | Not prime farmland | Highly erodible land | Moderately well drained | No | C | 4e | 4 | .15 |
| <i>CwC Cowarts gravelly loamy sand, 6 to 10 percent slopes</i> | | | | 137 Carolina and Georgia Sand Hills | | | | | | |
| Cowarts | 90 | 6-10 | Not prime farmland | Highly erodible land | Moderately well drained | No | C | 6e | 4 | .32 |
| <i>CwD Cowarts gravelly loamy sand, 10 to 15 percent slopes</i> | | | | 137 Carolina and Georgia Sand Hills | | | | | | |
| Cowarts | 90 | 10-15 | Not prime farmland | Highly erodible land | Moderately well drained | No | C | 6e | 4 | .32 |
| <i>CwE Cowarts gravelly loamy sand, 15 to 25 percent slopes</i> | | | | 137 Carolina and Georgia Sand Hills | | | | | | |
| Cowarts | 90 | 15-25 | Not prime farmland | Highly erodible land | Moderately well drained | No | C | 7e | 4 | .32 |
| <i>Cx Coxville loam</i> | | | | 133A Atlantic Coast Flatwoods | | | | | | |
| Coxville | 95 | 0-2 | Farmland of statewide importance | Not highly erodible land | Poorly drained | Yes | D | 3w | 5 | .24 |
| <i>EmA Emporia loamy sand, 0 to 2 percent slopes</i> | | | | 133A Southern Coastal Plain | | | | | | |
| Emporia | 95 | 0-2 | All areas are prime farmland | Not highly erodible land | Well drained | No | C | 1 | 5 | .15 |
| <i>EmB Emporia loamy sand, 2 to 6 percent slopes</i> | | | | 133A Southern Coastal Plain | | | | | | |
| Emporia | 90 | 2-6 | All areas are prime farmland | Not highly erodible land | Well drained | No | C | 2e | 5 | .15 |
| <i>EmB2 Emporia sandy loam, 2 to 6 percent slopes, eroded</i> | | | | 133A Southern Coastal Plain | | | | | | |
| Emporia | 90 | 2-6 | Farmland of statewide importance | Potentially highly erodible land | Well drained | No | C | 2e | 5 | .28 |
| <i>EuA Eunola loamy sand, 0 to 2 percent slopes</i> | | | | 133A Southern Coastal Plain | | | | | | |
| Eunola | 90 | 0-2 | All areas are prime farmland | Not highly erodible land | Moderately well drained | No | C | 2w | 5 | .15 |
| <i>FaA Faceville loamy sand, 0 to 2 percent slopes</i> | | | | 133A Southern Coastal Plain | | | | | | |
| Faceville | 95 | 0-2 | All areas are prime farmland | Not highly erodible land | Well drained | No | B | 1 | 5 | .17 |

Field Office Technical Guide Soils Information (SC)

Marlboro County, South Carolina

| Mapunit Symbol and Soil Name | % of Unit | Slope % | Important Farmland Class | Highly Erodible Land* | Drainage Class | Hydric Rating | Hydrologic Soil Group | Land Capability Class | T factor* | Kf* |
|---|-----------|---------|----------------------------------|----------------------------------|------------------------------|---------------|-----------------------|-----------------------|-----------|-----|
| <i>FaB Faceville loamy sand, 2 to 6 percent slopes</i> | | | | 133A Southern Coastal Plain | | | | | | |
| Faceville | 90 | 2-6 | All areas are prime farmland | Potentially highly erodible land | Well drained | No | B | 2e | 5 | .17 |
| <i>FaB2 Faceville sandy clay loam, 2 to 6 percent slopes, eroded</i> | | | | 133A Southern Coastal Plain | | | | | | |
| Faceville | 90 | 2-6 | Not prime farmland | Highly erodible land | Well drained | No | B | 3e | 5 | .32 |
| <i>FxB Foxworth sand, 0 to 6 percent slopes</i> | | | | 133A Southern Coastal Plain | | | | | | |
| Foxworth | 95 | 0-6 | Not prime farmland | Not highly erodible land | Somewhat excessively drained | No | A | 3s | 5 | .10 |
| <i>GoA Goldsboro loamy sand, 0 to 2 percent slopes</i> | | | | 133A Atlantic Coast Flatwoods | | | | | | |
| Goldsboro | 85 | 0-2 | All areas are prime farmland | Not highly erodible land | Moderately well drained | No | B | 2w | 5 | .17 |
| <i>HnA Hornsville loam, 0 to 2 percent slopes</i> | | | | 133A Southern Coastal Plain | | | | | | |
| Hornsville | 90 | 0-2 | All areas are prime farmland | Not highly erodible land | Moderately well drained | No | C | 2w | 5 | .20 |
| <i>HnB Hornsville loam, 2 to 6 percent slopes</i> | | | | 133A Southern Coastal Plain | | | | | | |
| Hornsville | 90 | 2-6 | All areas are prime farmland | Potentially highly erodible land | Moderately well drained | No | C | 2e | 5 | .24 |
| <i>HnB2 Hornsville sandy clay loam, 2 to 6 percent slopes, eroded</i> | | | | 133A Southern Coastal Plain | | | | | | |
| Hornsville | 90 | 2-6 | Farmland of statewide importance | Highly erodible land | Moderately well drained | No | C | 3e | 5 | .24 |
| <i>Jo Johnston mucky loam, frequently flooded</i> | | | | 133A Southern Coastal Plain | | | | | | |
| Johnston | 95 | 0-2 | Not prime farmland | Not highly erodible land | Very poorly drained | Yes | D | 7w | 5 | .20 |
| <i>Kn Kinston loam, frequently flooded</i> | | | | 133A Southern Coastal Plain | | | | | | |
| Kinston | 95 | 0-2 | Not prime farmland | Not highly erodible land | Poorly drained | Yes | B/D | 6w | 5 | .37 |
| <i>Le Leon sand</i> | | | | 133A Southern Coastal Plain | | | | | | |
| Leon | 100 | 0-2 | Not prime farmland | Not highly erodible land | Poorly drained | Yes | B/D | 4w | 5 | .10 |
| <i>LuB Lucy sand, 0 to 6 percent slopes</i> | | | | 133A Southern Coastal Plain | | | | | | |
| Lucy | 95 | 0-6 | Farmland of statewide importance | Not highly erodible land | Well drained | No | A | 2s | 5 | .10 |
| <i>Ly Lynchburg sandy loam</i> | | | | 133A Atlantic Coast Flatwoods | | | | | | |
| Lynchburg | 90 | 0-2 | Prime farmland if drained | Not highly erodible land | Somewhat poorly drained | No | C | 2w | 5 | .20 |

Field Office Technical Guide Soils Information (SC)

Marlboro County, South Carolina

| Mapunit Symbol and Soil Name | % of Unit | Slope % | Important Farmland Class | Highly Erodible Land* | Drainage Class | Hydric Rating | Hydrologic Soil Group | Land Capability Class | T factor* | Kf* |
|--|-----------|---------|----------------------------------|----------------------------------|----------------|---------------|-----------------------|-----------------------|-----------|-----|
| <i>MaA Marlboro fine sandy loam, moderately wet, 0 to 2 percent slopes</i> | | | | 133A Southern Coastal Plain | | | | | | |
| Marlboro | 90 | 0-2 | All areas are prime farmland | Not highly erodible land | Well drained | No | B | 1 | 5 | .20 |
| <i>MaB Marlboro fine sandy loam, moderately wet, 2 to 6 percent slopes</i> | | | | 133A Southern Coastal Plain | | | | | | |
| Marlboro | 90 | 2-6 | All areas are prime farmland | Potentially highly erodible land | Well drained | No | B | 2e | 5 | .20 |
| <i>Mc McColl loam</i> | | | | 133A Southern Coastal Plain | | | | | | |
| McColl | 95 | 0-1 | Farmland of statewide importance | Not highly erodible land | Poorly drained | Yes | D | 3w | 5 | .24 |
| <i>NaB Nankin loamy fine sand, 2 to 6 percent slopes</i> | | | | 133A Southern Coastal Plain | | | | | | |
| Nankin | 90 | 2-6 | Farmland of statewide importance | Not highly erodible land | Well drained | No | C | 2e | 3 | .17 |
| <i>NbB2 Nankin sandy clay loam, 2 to 6 percent slopes, eroded</i> | | | | 133A Southern Coastal Plain | | | | | | |
| Nankin | 90 | 2-6 | Farmland of statewide importance | Highly erodible land | Well drained | No | C | 3e | 3 | .17 |
| <i>NbC2 Nankin sandy clay loam, 6 to 10 percent slopes, eroded</i> | | | | 133A Southern Coastal Plain | | | | | | |
| Nankin | 90 | 6-10 | Not prime farmland | Highly erodible land | Well drained | No | C | 4e | 3 | .17 |
| <i>NcA Noboco loamy sand, 0 to 2 percent slopes</i> | | | | 133A Southern Coastal Plain | | | | | | |
| Noboco | 90 | 0-2 | All areas are prime farmland | Not highly erodible land | Well drained | No | B | 1 | 5 | .10 |
| <i>NcB Noboco loamy sand, 2 to 6 percent slopes</i> | | | | 133A Southern Coastal Plain | | | | | | |
| Noboco | 90 | 2-6 | All areas are prime farmland | Potentially highly erodible land | Well drained | No | B | 2e | 5 | .10 |
| <i>NcB2 Noboco sandy loam, 2 to 6 percent slopes, eroded</i> | | | | 133A Southern Coastal Plain | | | | | | |
| Noboco | 90 | 2-6 | Farmland of statewide importance | Highly erodible land | Well drained | No | B | 2e | 5 | .10 |
| <i>NoA Norfolk loamy sand, 0 to 2 percent slopes</i> | | | | 133A Southern Coastal Plain | | | | | | |
| Norfolk | 90 | 0-2 | All areas are prime farmland | Not highly erodible land | Well drained | No | B | 1 | 5 | .10 |
| <i>NoB Norfolk loamy sand, 2 to 6 percent slopes</i> | | | | 133A Southern Coastal Plain | | | | | | |
| Norfolk | 90 | 2-6 | All areas are prime farmland | Not highly erodible land | Well drained | No | B | 2e | 5 | .10 |
| <i>NrB2 Norfolk sandy clay loam, 2 to 6 percent slopes, eroded</i> | | | | 133A Southern Coastal Plain | | | | | | |
| Norfolk | 80 | 2-6 | Farmland of statewide importance | Highly erodible land | Well drained | No | B | 2e | 5 | .10 |

Field Office Technical Guide Soils Information (SC)

Marlboro County, South Carolina

| Mapunit Symbol and Soil Name | % of Unit | Slope % | Important Farmland Class | Highly Erodible Land* | Drainage Class | Hydric Rating | Hydrologic Soil Group | Land Capability Class | T factor* | Kf* |
|--|-----------|---------|-------------------------------------|--|------------------------------|---------------|-----------------------|-----------------------|-----------|-----|
| <i>OcB</i> <i>Ocilla sand, 0 to 4 percent slopes</i> | | | | <i>133A Southern Coastal Plain</i> | | | | | | |
| Ocilla | 90 | 0-2 | Farmland of statewide importance | Not highly erodible land | Somewhat poorly drained | No | C | 3w | 5 | .10 |
| <i>Og</i> <i>Ogeechee sandy loam</i> | | | | <i>133A Southern Coastal Plain</i> | | | | | | |
| Ogeechee | 95 | 0-2 | Farmland of statewide importance | Not highly erodible land | Poorly drained | Yes | B/D | 3w | 5 | .10 |
| <i>OrA</i> <i>Orangeburg loamy sand, 0 to 2 percent slopes</i> | | | | <i>133A Southern Coastal Plain</i> | | | | | | |
| Orangeburg | 95 | 0-2 | All areas are prime farmland | Not highly erodible land | Well drained | No | B | 1 | 5 | .10 |
| <i>Pa</i> <i>Pamlico muck, frequently flooded</i> | | | | <i>133A Southern Coastal Plain</i> | | | | | | |
| Pamlico | 95 | 0-1 | Not prime farmland | Not highly erodible land | Very poorly drained | Yes | D | 7w | 2 | |
| <i>Pe</i> <i>Paxville fine sandy loam</i> | | | | <i>133A Southern Coastal Plain</i> | | | | | | |
| Paxville | 95 | 0-2 | Farmland of statewide importance | Not highly erodible land | Very poorly drained | Yes | B/D | 6w | 4 | .20 |
| <i>PnA</i> <i>Pelion loamy sand, 0 to 2 percent slopes</i> | | | | <i>137 Carolina and Georgia Sand Hills</i> | | | | | | |
| Pelion | 95 | 0-2 | Farmland of statewide importance | Not highly erodible land | Moderately well drained | No | B/D | 2w | 3 | .15 |
| <i>PnB</i> <i>Pelion loamy sand, 2 to 6 percent slopes</i> | | | | <i>137 Carolina and Georgia Sand Hills</i> | | | | | | |
| Pelion | 90 | 2-6 | Farmland of statewide importance | Not highly erodible land | Moderately well drained | No | B/D | 2e | 3 | .15 |
| <i>PrA</i> <i>Persanti loam, 0 to 2 percent slopes</i> | | | | <i>133A Southern Coastal Plain</i> | | | | | | |
| Persanti | 95 | 0-2 | All areas are prime farmland | Not highly erodible land | Moderately well drained | No | C | 2w | 5 | .28 |
| <i>Qz</i> <i>Quartzipsammets, sloping</i> | | | | <i>133A Southern Coastal Plain</i> | | | | | | |
| Quartzipsamm | 95 | 0-10 | Not prime farmland | Not highly erodible land | Somewhat excessively drained | No | A | 6s | 5 | .15 |
| <i>Ra</i> <i>Rains sandy loam</i> | | | | <i>133A Atlantic Coast Flatwoods</i> | | | | | | |
| Rains | 95 | 0-2 | Farmland of statewide importance | Not highly erodible land | Poorly drained | Yes | B/D | 3w | 5 | .20 |
| <i>Rv</i> <i>Riverview fine sandy loam, occasionally flooded</i> | | | | <i>133A Southern Coastal Plain</i> | | | | | | |
| Riverview | 90 | 0-2 | Prime farmland if protected from fl | Not highly erodible land | Well drained | No | B | 4w | 5 | .24 |
| <i>Sm</i> <i>Smithboro silt loam</i> | | | | <i>133A Southern Coastal Plain</i> | | | | | | |
| Smithboro | 90 | 0-2 | Farmland of statewide importance | Not highly erodible land | Somewhat poorly drained | No | D | 3w | 5 | .37 |

Field Office Technical Guide Soils Information (SC)

Marlboro County, South Carolina

| Mapunit Symbol and Soil Name | % of Unit | Slope % | Important Farmland Class | Highly Erodible Land* | Drainage Class | Hydric Rating | Hydrologic Soil Group | Land Capability Class | T factor* | Kf* |
|---|-----------|---------|----------------------------------|------------------------------------|------------------------------|---------------|-----------------------|-----------------------|-----------|-----|
| <i>TbB Tarboro sand, 0 to 4 percent slopes</i> | | | | <i>133A Southern Coastal Plain</i> | | | | | | |
| Tarboro | 90 | 0-4 | Not prime farmland | Not highly erodible land | Somewhat excessively drained | No | A | 3s | 5 | .10 |
| <i>TrB Troup sand, 0 to 6 percent slopes</i> | | | | <i>133A Southern Coastal Plain</i> | | | | | | |
| Troup | 90 | 0-6 | Not prime farmland | Not highly erodible land | Somewhat excessively drained | No | A | 3s | 5 | .10 |
| <i>TrC Troup sand, 6 to 10 percent slopes</i> | | | | <i>133A Southern Coastal Plain</i> | | | | | | |
| Troup | 90 | 6-10 | Not prime farmland | Potentially highly erodible land | Somewhat excessively drained | No | A | 6s | 5 | .10 |
| <i>UgB Uchee sand, 0 to 6 percent slopes</i> | | | | <i>133A Southern Coastal Plain</i> | | | | | | |
| Uchee | 90 | 0-6 | Farmland of statewide importance | Not highly erodible land | Well drained | No | A | 2s | 5 | .15 |
| <i>UgC Uchee sand, 6 to 10 percent slopes</i> | | | | <i>133A Southern Coastal Plain</i> | | | | | | |
| Uchee | 90 | 6-10 | Not prime farmland | Highly erodible land | Well drained | No | A | 3s | 5 | .15 |
| <i>UhB Uchee gravelly sand, 0 to 6 percent slopes</i> | | | | <i>133A Southern Coastal Plain</i> | | | | | | |
| Uchee | 95 | 0-6 | Farmland of statewide importance | Not highly erodible land | Well drained | No | A | 2s | 5 | .10 |
| <i>UhC Uchee gravelly sand, 6 to 10 percent slopes</i> | | | | <i>133A Southern Coastal Plain</i> | | | | | | |
| Uchee | 98 | 6-10 | Farmland of statewide importance | Highly erodible land | Well drained | No | A | 3s | 5 | .10 |
| <i>Ur Udorthents</i> | | | | <i>133A Southern Coastal Plain</i> | | | | | | |
| Udorthents | 95 | 0-2 | Not prime farmland | Highly erodible land | Well drained | No | B | 4e | 5 | .28 |
| <i>WaB Wagram sand, 0 to 6 percent slopes</i> | | | | <i>133A Southern Coastal Plain</i> | | | | | | |
| Wagram | 95 | 0-6 | Farmland of statewide importance | Not highly erodible land | Well drained | No | A | 2s | 5 | .10 |
| <i>WkA Wickham fine sandy loam, 0 to 2 percent slopes</i> | | | | <i>133A Southern Coastal Plain</i> | | | | | | |
| Wickham | 95 | 0-2 | All areas are prime farmland | Not highly erodible land | Well drained | No | B | 1 | 5 | .24 |