


South Carolina Longleaf Initiative Sumter County




Updated February 2012




Legend

Tiers

-  **1** Highly Suitable
-  **2** Moderately Suitable
-  **3** Slightly Suitable
-  **4** Unsuitable
-  Open Water



1:400,000



Map Unit Symbol	Map Unit Name	Tier
AaD	Ailey-Troup-Alpin complex, 10 to 15 percent slopes	1
AgB	Alaga loamy coarse sand, 0 to 6 percent slopes	1
AnB	Alaga-Blanton-Johns complex, 0 to 2 percent slopes, rarely flooded	1
ApB	Alpin-Candor-Troup complex, 0 to 6 percent slopes	1
AuB	Autryville-Norfolk complex, 0 to 4 percent slopes	1
BaB	Barnwell-Fuquay complex, 2 to 6 percent slopes	1
BoB	Bonneau-Norfolk complex, 0 to 6 percent slopes	1
BuA	Butters-Blanton complex, 0 to 2 percent slopes	1
CxA	Coxville-Rains complex, 0 to 2 percent slopes	3
DoA	Dothan-Norfolk complex, 0 to 2 percent slopes	1
DoB	Dothan-Norfolk complex, 2 to 6 percent slopes	1
FaB2	Faceville sandy loam, 2 to 6 percent slopes, moderately eroded	1
FcB	Faceville-Lucy complex, 2 to 6 percent slopes	1
FuB	Fuquay-Dothan complex, 0 to 6 percent slopes	1
GoA	Goldsboro-Noboco complex, 0 to 2 percent slopes	2
JnA	Johnston mucky sandy loam, 0 to 2 percent slopes, frequently flooded	4
JoA	Johnston mucky sandy loam, 0 to 2 percent slopes, ponded	4
KaA	Kalmia-Johns complex, 0 to 2 percent slopes, rarely flooded	2
LaD	Lakeland sand, 6 to 15 percent slopes	1
LbA	Lumbee-Johns complex, 0 to 2 percent slopes, rarely flooded	3
LeA	Lumbee-Rutlege complex, 0 to 2 percent slopes	4
LfA	Lynchburg-Foreston-Butters complex, 0 to 2 percent slopes	3
LyA	Lynchburg-Rains complex, 0 to 2 percent slopes	3
MaA	Mantachie-Mimms complex, 0 to 2 percent slopes, frequently flooded	4
MdA	Masada-Hornsville complex, 0 to 2 percent slopes, very rarely flooded	2
MeA	Meggett-Lumbee complex, 0 to 2 percent slopes, rarely flooded	4
NbA	Noboco-Norfolk complex, 0 to 2 percent slopes	1
NfA	Norfolk-Butters complex, 0 to 2 percent slopes	1
NnB	Norfolk-Faceville-Noboco complex, 2 to 6 percent slopes	1
NoA	Norfolk-Noboco complex, 0 to 2 percent slopes	1
NoB	Norfolk-Noboco complex, 2 to 6 percent slopes	1
OkA	Okeetee-Yemassee complex, 0 to 2 percent slopes, rarely flooded	2
OrA	Orangeburg loamy sand, 0 to 2 percent slopes	1
OuB	Orangeburg-Lucy complex, 2 to 6 percent slopes	1
PuD	Pits-Udorthents, loamy substratum complex, 0 to 15 percent slopes	4
RaA	Rains sandy loam, 0 to 2 percent slopes	3
RcA	Rains-Coxville-Lynchburg complex, 0 to 2 percent slopes	3
RmB	Rimini sand, 0 to 6 percent slopes	1
ScA	Scapo-Mouzon complex, 0 to 2 percent slopes, frequently flooded	4
ShA	Shellbluff-Tawcaw complex, 0 to 2 percent slopes, frequently flooded	4
SmA	Smithboro-Persanti complex, 0 to 2 percent slopes	3
SpD	Springhill-Lucy-Nankin complex, 6 to 15 percent slopes	1
TaA	Tawcaw-Duckbottom-Mullers complex, 0 to 2 percent slopes, frequently flooded	4
TcA	Tawcaw-Shellbluff-Mullers complex, 0 to 2 percent slopes, frequently flooded	4
ThA	Thursa loamy sand, 0 to 2 percent slopes	1
ThB	Thursa loamy sand, 2 to 6 percent slopes	1
TpB	Troup-Lucy complex, 0 to 6 percent slopes	1
TrD	Troup-Lucy-Nankin complex, 10 to 15 percent slopes	1
UdC	Udorthents, reclaimed, 0 to 10 percent slopes	4
UpD	Udorthents, refuse substratum-Pits complex, 0 to 15 percent slopes	4
VaB	Vaocluse-Ailey complex, 2 to 6 percent slopes	1
VaD	Vaocluse-Ailey complex, 6 to 15 percent slopes	1
VcF	Vaocluse-Ailey-Lucy complex, 6 to 45 percent slopes	1
WaB	Wagram-Norfolk-Lucknow complex, 0 to 4 percent slopes	1
WuD	Water-Udorthents, gravelly substratum, 0 to 15 percent slopes	4
YeA	Yemassee-Johns complex, 0 to 2 percent slopes, rarely flooded	3

Tier values reflect undrained conditions; if site is drained move up 1 tier (e.g., Tier 4 to Tier 3)