

## FORAGE SUITABILITY GROUP (FSG)

### Loam


**FSG No.:** G053BY100ND

**Major Land Resource Area (MLRA):** 53B - Central Dark Brown Glaciated Plains

#### Physiographic Features

Soils in this group typically occur in upland positions of glacial till, outwash, and lake plains. They also occur on fans, terraces, and flood plains.

	<u>Minimum</u>	<u>Maximum</u>
<b>Elevation (feet):</b>	1640	1970
<b>Slope (percent):</b>	0	15
<b>Flooding:</b>		
<b>Frequency:</b>	None	Occasional
<b>Duration:</b>	None	Brief
<b>Ponding:</b>		
<b>Depth (inches):</b>		
<b>Frequency:</b>	None	None
<b>Duration:</b>	None	None
<b>Runoff Class:</b>	Very low	High



#### Climatic Features

This group occurs in a mid-continental climate characterized by wide seasonal temperature and precipitation fluctuations and extremes.

Annual precipitation varies widely from year to year in MLRA 53B. Average annual precipitation for all climate stations listed below is about 17 inches. About 79 percent of that occurs during the months of April through September. On average, there are about 27 days with greater than .1 inches of precipitation during the same time frame. Precipitation is less than needed for optimum forage production and is the single largest factor limiting production from this group on non-irrigated lands.

Average annual snowfall ranges from 21 inches at Turtle Lake, North Dakota (ND), to 38 inches at Eureka, South Dakota (SD). Days with snow cover at depths greater than 1 inch range from 22 at Garrison, ND, to 100 at Max, ND.

Average July temperatures are about 70°F and average January temperatures are about 8°F. Recorded temperature extremes in the MLRA during the years 1961 to 1990 are a low of -48°F at Powers Lake, ND, and a high of 111 recorded at Linton, ND. The MLRA lies in USDA Plant Hardiness Zones 3b and 4a.

At Bismarck, the average morning relative humidity in June is about 84 percent and average afternoon humidity is 55 percent. It is cloudy an average of 165 days a year.

The climate data listed in the tables below represent high and low ranges and averages for the climate stations and dates listed. For additional climate data, access the National Water and Climate Center at <http://www.wcc.nrcs.usda.gov>.

	<b>From</b>	<b>To</b>
<b>Freeze-free period (28 deg) (days):</b> (9 years in 10 at least)	100	134
<b>Last Killing Freeze in Spring (28 deg):</b> (1 year in 10 later than)	Jun 06	May 16
<b>Last Frost in Spring (32 deg):</b> (1 year in 10 later than)	Jun 18	May 26
<b>First Frost in Fall (32 deg):</b> (1 year in 10 earlier than)	Aug 23	Sep 12
<b>First Killing Freeze in Fall (28 deg):</b> (1 year in 10 earlier than)	Sep 02	Sep 19
<b>Length of Growing Season (32 deg) (days):</b> (9 years in 10 at least)	78	116
<b>Growing Degree Days (40 deg):</b>	3317	4367
<b>Growing Degree Days (50 deg):</b>	1793	2441
<b>Annual Minimum Temperature:</b>	-35	-25
<b>Mean annual precipitation (inches):</b>	15	19

**Monthly precipitation (inches) and temperature (F):**

<b>2 years in 10:</b>	<b>Jan</b>	<b>Feb</b>	<b>Mar</b>	<b>Apr</b>	<b>May</b>	<b>Jun</b>	<b>Jul</b>	<b>Aug</b>	<b>Sep</b>	<b>Oct</b>	<b>Nov</b>	<b>Dec</b>
<b>Precip. Less Than</b>	0.17	0.14	0.26	0.37	0.95	1.79	0.86	0.65	0.65	0.22	0.13	0.23
<b>Precip. More Than</b>	0.63	0.80	1.96	3.53	3.81	4.82	3.82	2.89	2.63	1.66	1.07	0.70
<b>Monthly Average:</b>	0.42	0.42	0.82	1.80	2.30	3.21	2.49	1.96	1.69	0.98	0.46	0.43
<b>Temp. Min.</b>	-1.5	4.9	18.8	31.6	43.3	53.4	58.8	55.4	44.1	32.5	18.7	4.1
<b>Temp. Max.</b>	30.6	36.4	47.0	62.4	73.4	83.0	90.4	88.6	78.2	65.5	46.7	33.4
<b>Temp. Avg.</b>	8.0	14.3	26.6	42.2	54.8	64.3	70.2	68.2	56.8	45.2	27.8	13.1

<b><u>Climate Station</u></b>	<b><u>Location</u></b>	<b><u>From</u></b>	<b><u>To</u></b>
ND0961	Bowbells, ND	1961	1990
ND7281	Powers Lake, ND	1961	1990
ND1225	Butte, ND	1961	1990
ND3376	Garrison, ND	1961	1990
ND5638	Max, ND	1961	1990
ND8804	Turtle Lake, ND	1961	1990
ND8872	Underwood, ND	1961	1990
ND0382	Ashley, ND	1961	1990
ND9515	Wishek, ND	1961	1990
ND5210	Linton, ND	1961	1990
SD2797	Eureka, SD	1961	1990
SD4891	Leola, SD	1961	1990
SD4206	Ipswich, SD	1961	1990

**Soil Interpretations**

This group consists of very deep, well drained, moderately coarse to moderately fine textured soils formed mostly from loess, glacial till, alluvium, and colluvium. Available water capacity is mostly high and permeability is moderately slow to moderately rapid.

<b>Drainage Class:</b>	Moderately well drained	To	Well drained
<b>Permeability Class:</b> (0 - 40 inches)	Moderately slow	To	Moderately rapid
<b>Frost Action Class:</b>	Moderate	To	Moderate

	<u>Minimum</u>	<u>Maximum</u>
<b>Depth:</b>	72	
<b>Surface Fragments &gt;3" (% Cover):</b>	0	3
<b>Organic Matter (percent):</b> (surface layer)	1.0	8.0
<b>Electrical Conductivity (mmhos/cm):</b> (0 - 24 inches)	0	4
<b>Sodium Absorption Ratio:</b> (0 - 12 inches)	0	1
<b>Soil Reaction (1:1) Water (pH):</b> (0 - 12 inches)	5.6	8.4
<b>Available Water Capacity (inches):</b> (0 - 60 inches)	6	13
<b>Calcium Carbonate Equivalent (percent):</b> (0 - 12 inches)	0	15

### Adapted Species List

The following forage species are considered adapted to grow on the soils in this group. Additional information concerning plant characteristics of a number of the listed species as well as individual cultivars of many of those species can be accessed at <http://plants.usda.gov/>.

<u>Cool Season Grasses</u>	<u>Symbol</u>	<u>Adapted</u>	<u>Warm Season Grasses</u>	<u>Symbol</u>	<u>Adapted</u>
Altai wildrye	LEAN3	G	Big bluestem	ANGE	G
Basin wildrye	LECI4	F	Blue grama	BOGR2	G
Canada wildrye	ELCA4	G	Indiangrass	SONU2	F
Crested wheatgrass	AGCR	G	Little bluestem	SCSC	G
Dahurian wildrye	ELDA3	G	Prairie sandreed	CALO	F
Green needlegrass	NAVI4	G	Sideoats grama	BOCU	G
Intermediate wheatgrass	THIN6	G	Switchgrass	PAVIV	G
Meadow brome grass	BRBI2	G			
Newhy hybrid wheatgrass		G	<u>Legumes</u>		
Pubescent wheatgrass	THIN6	G	Alfalfa	MESA	G
Russian wildrye	PSJU3	G	American vetch	VIAM	G
Slender wheatgrass	ELTR7	G	Canada milkvetch	ASCAC6	G
Smooth brome grass	BRINI2	G	Cicer milkvetch	ASCI4	G
Tall wheatgrass	THPO7	G	Hairy Vetch	VIVI	G
Western wheatgrass	PASM	G	Purple prairieclover	DAPUP	G
			Sainfoin	ONVI	F
			Sweetclover	MELIL	G
			White clover	TRRE3	F
			White prairie clover	DACAC	G

G - Good adaptation for forage production on this group of soils in this MLRA

F - Fair adaptation but will not produce at its highest potential

### Production Estimates

Production estimates listed here should only be used for making general management recommendations. Onsite production information should always be used for making detailed planning and management recommendations.

The high forage production estimates listed below are based on dense, vigorous stands of climatically adapted, superior performing cultivars. They are properly fertilized for high yields and pest infestations are kept below economic thresholds. Mechanical harvests are managed to maintain stand life by cutting at appropriate stages of maturity and harvest intervals. If grazed, optimum beginning and ending grazing heights are adhered to. Adequate time is allowed for plant recovery before entering winter dormancy under both uses.

The production estimates listed below represent total annual above ground plant production on an air-dry-matter basis. Estimates of hay and grazing yields can be calculated from these numbers by multiplying them by predicted harvest efficiency. Seventy percent harvest efficiency is commonly used when converting to hay yields. Pasture harvest efficiency is highly dependent on the grazing management system applied, ranging from 25 to 50 percent.

**Forage Crop****Management Intensity**

	<u>High</u> (lbs/ac)	<u>Low</u> (lbs/ac)
Alfalfa	8600	3700
Alfalfa/Intermediate wheatgrass	7100	3100
Alfalfa/smooth brome grass	7100	3100
Big bluestem	6300	2700
Crested wheatgrass	5500	2300
Green needlegrass	4300	1900
Intermediate wheatgrass	5900	2500
Smooth brome grass	5900	2500
Switchgrass	7000	2900
Western wheatgrass	4300	1900

**Forage Growth Curves**

Growth curves estimate the seasonal distribution of growth of the various forage crops. They indicate when the forages may be available for grazing or mechanical harvest.

**Growth Curve Number:** ND0001

**Growth Curve Name:** Alfalfa

**Growth Curve Description:** Alfalfa

**Percent Production by Month**

<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	5	25	30	20	15	5	0	0	0

**Growth Curve Number:** ND0002

**Growth Curve Name:** Cool season grass

**Growth Curve Description:** Cool season grass

**Percent Production by Month**

<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	5	40	35	10	5	5	0	0	0

**Growth Curve Number:** ND0003

**Growth Curve Name:** Warm season grass

**Growth Curve Description:** Warm season grass

**Percent Production by Month**

<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	0	10	40	35	15	0	0	0	0

**Soil Limitations**

These soils have few limitations to the production of climatically adapted forage crops. On steeper slopes, water erosion is a potential problem during establishment, when renovating stands, and in thin established stands. Livestock trail erosion is a potential problem in established stands.

**Management Interpretations**

Including sod forming grass species in stands, especially on steeper slopes, will reduce the potential for sheet and rill erosion. Incorporate erosion control practices during the establishment period. Properly locating facilitating practices such as fences, lanes, and water developments can help control livestock movement, reduce trailing perpendicular to steeper slopes, and evenly distribute grazing pressure.

Pasture and hayland can include considerations for wildlife. Delaying grazing on portions of the pasture or rotating pastures will allow nest initiation of grassland nesting birds or species of concern. Nest initiation of most grassland nesting birds occurs from April 15 to June 1. Delaying haying until after July 15 allows for most species to fledge their young. Consider planting species with later maturity to allow for harvesting after nests have fledged. Avoid mowing around the field. Mow back and forth or from the inside to the outside of the field. Consider using flushing bars on swathers and mowers.

**FSG Documentation**

**Similar FSG's:**

**FSG ID**

G053BY120ND

**FSG Narrative**

Droughty Loam soils are shallower or coarser textured resulting in lower available water capacity and lower production potential.

G053BY500ND

Overflow soils receive additional moisture due to a favorable landscape position resulting in a higher production potential.

**Inventory Data References:**

Agriculture Handbook 296-Land Resource Regions and Major Land Resource Areas  
Natural Resources Conservation Service (NRCS) National Water and Climate Center data  
USDA Plant Hardiness Zone Maps  
National Soil Survey Information System for soil surveys in North Dakota and South Dakota counties in MLRA 53B  
NRCS North Dakota Field Office Technical Guide and South Dakota Technical Guide  
NRCS National Range and Pasture Handbook  
Various Agricultural Research Service, Cooperative Extension Service, and NRCS research trials for plant adaptation and production.

**State Correlation:**

This site has been correlated with the following states: North and South Dakota

**Forage Suitability Group Approval:**

Original Author: Tim Nordquist

Original Date: 1/10/05

Approval by: Dave Schmidt

Approval Date: 1/10/05