

FORAGE SUITABILITY GROUP Overflow

FSG No.: G053CY500SD

Major Land Resource Area: 53C - Southern Dark Brown Glaciated Plains

Physiographic Features

The soils in this group are generally found in nearly level to gently sloping positions on stream terraces and flood plains, and in swales and drainage ways on uplands. They receive beneficial additional moisture as run-on from up slope, or from flooding.

	<u>Minimum</u>	<u>Maximum</u>
Elevation (feet):	1300	2300
Slope (percent):	0	3
Flooding:		
Frequency:	None	Frequent
Duration:	None	Brief
Ponding:		
Depth (inches):		
Frequency:	None	None
Duration:	None	None
Runoff Class:	Negligible	Medium

Climatic Features

This group occurs in a mid-continental climate characterized by wide seasonal temperature and precipitation fluctuations and extremes.

Annual precipitation varies widely from year to year in MLRA 53C. Average annual precipitation for all climate stations listed below is about 20 inches. About 75 percent of that occurs during the months of April through September. On average, there are about 28 days with greater than .1 inches of precipitation during the same timeframe. Annual precipitation and temperature increase from the north to the south in the MLRA. Precipitation is less than needed for optimum forage production and is the single largest factor limiting production from this group on non-irrigated lands.

Average annual snowfall ranges from 23 inches at Pickstown to 46 inches at Harrold. Snow cover at depths greater than 1 inch range from 38 days at Gettysburg to 65 days at Harrold.

Average July temperatures are about 75⁰F and average January temperatures are about 16⁰F. Recorded temperature extremes in the MLRA during the years 1961 to 1990 are a low of -40 and a high of 113 both recorded at Harrold. The MLRA lies mostly in USDA Plant Hardiness Zones 4a and 4b, with a small area of warmer 5a along the Missouri River.

At Huron, the closest station with records, the average annual wind speeds are about 11.5 mph. The highest wind speeds occur during March through May. It is cloudy about 154 days a year. Average morning relative humidity in June is about 86 percent and average afternoon humidity is 59 percent.

The climate data listed in the tables below represent high and low ranges and averages for the climate stations and dates listed. For additional climate data, access the National Water and Climate Center at <http://www.wcc.nrcs.usda.gov>.

	From	To
Freeze-free period (28 deg)(days): (9 years in 10 at least)	122	156
Last Killing Freeze in Spring (28 deg):	May 24	May 07
Last Frost in Spring (32 deg): (1 year in 10 later than)	Jun 14	May 17
First Frost in Fall (32 deg): (1 year in 10 earlier than)	Aug 27	Sep 23
First Killing Freeze in Fall (28 deg): (1 year in 10 earlier than)	Sep 16	Oct 04
Length of Growing Season (32 deg)(days): (9 years in 10 at least)	85	136
Growing Degree Days (40 deg):	4388	5543
Growing Degree Days (50 deg):	2532	3338
Annual Minimum Temperature:	-30	-15
Mean annual precipitation (inches):	17	25

Monthly precipitation (inches) and temperature (F)

2 years in 10:	<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
Precip. Less Than	0.10	0.13	0.55	1.00	1.21	1.40	1.26	1.02	0.46	0.47	0.15	0.19
Precip. More Than	0.62	0.83	2.36	3.48	4.20	5.55	3.95	3.29	3.60	2.43	1.49	0.94
Monthly Average:	0.40	0.55	1.38	2.23	2.88	3.25	2.74	2.21	1.78	1.37	0.70	0.55
Temp. Min.	1.3	7.5	18.2	31.1	42.2	52.3	58.2	55.5	44.9	32.8	18.9	6.1
Temp. Max.	30.1	35.8	47.6	63.8	75.1	85.3	91.1	88.7	78.0	65.5	47.0	32.9
Temp. Avg.	15.7	21.9	32.8	47.3	58.6	68.5	75.1	72.9	62.2	49.8	33.5	19.6

<u>Climate Station</u>	<u>Location</u>	<u>From</u>	<u>To</u>
SD6574	Pickstown, SD	1961	1990
SD8767	Wagner, SD	1961	1990
SD3294	Gettysburg, SD	1961	1990
SD3608	Harrold, SD	1963	1990
SD0043	Academy, SD	1961	1990
SD6292	Onida, SD	1961	1990
SD3832	Highmore, SD	1961	1990
SD7992	Stephan, SD	1961	1990

Soil Interpretations

This group consists of very deep, moderately well and well-drained, medium to moderately fine textured soils formed from glacial till and silty and loamy sediments and alluvium. Permeability is slow to moderate. Many of these soils have a seasonal water table within three feet of the surface, which benefits deep rooted plants.

Drainage Class:	Moderately well drained	To	Well drained
Permeability Class: (0 - 40 inches)	Slow	To	Moderate
Frost Action Class:	Moderate	To	High

	<u>Minimum</u>	<u>Maximum</u>
Depth:	72	
Surface Fragments >3" (% Cover):	0	3
Organic Matter (percent): (surface layer)	2.0	6.0
Electrical Conductivity (mmhos/cm): (0 - 24 inches)	0	4
Sodium Absorption Ratio: (0 - 12 inches)	0	1
Soil Reaction (1:1) Water (pH): (0 - 12 inches)	5.6	8.4
Available Water Capacity (inches): (0 - 60 inches)	9	

	<u>Minimum</u>	<u>Maximum</u>
Calcium Carbonate Equivalent (percent): (0 - 12 inches)	0	9

Adapted Species List

The following forage species are considered adapted to grow on the soils in this group. Additional information concerning plant characteristics of a number of the listed species as well as individual cultivars of many those species can be accessed at <http://plants.usda.gov/>

Cool Season Grasses

Altai wildrye	F
Canada wildrye	G
Creeping foxtail	F
Crested wheatgrass	G
Green needlegrass	G
Intermediate wheatgrass	G
Meadow bromegrass	G
Newhy hybrid wheatgrass	G
Pubescent wheatgrass	G
Reed canarygrass	F
Russian wildrye	G
Slender wheatgrass	G
Smooth bromegrass	G
Tall wheatgrass	G
Western wheatgrass	G

Warm Season Grasses

Big bluestem	G
Indiangrass	G
Little bluestem	G
Sand bluestem	F
Sideoats grama	G
Switchgrass	G

Legumes

Alfalfa	G
Birdsfoot trefoil	G
Canada milkvetch	G
Cicer milkvetch	G
Purple prairieclover	F
Sainfoin	F
Sweetclover	G
White prairieclover	F

G - Good adaptation for forage production on this group of soils in this MLRA

F - Fair adaptation but will not produce at its highest potential

Production Estimates

Production estimates listed here should only be used for making general management recommendations. Onsite production information should always be used for making detailed planning and management recommendations.

The high forage production estimates listed below are based on dense, vigorous stands of climatically adapted, superior performing cultivars. They are properly fertilized for high yields and pest infestations are kept below economic thresholds. Mechanical harvests are managed to maintain stand life by cutting at appropriate stages of maturity and harvest intervals. If grazed, optimum beginning and ending grazing heights are adhered to. Adequate time is allowed for plant recovery before entering winter dormancy under both uses.

The production estimates listed below represent total annual above ground plant production on an air-dry-matter basis. Estimates of hay and grazing yields can be calculated from these numbers by multiplying them by a harvest efficiency. A 70 percent harvest efficiency is commonly used when converting to hay yields. Pasture harvest efficiency is highly dependent on the grazing management system applied, ranging from 25 to 50 percent.

Forage Crop	Management Intensity	
	<u>High</u> (lbs/ac)	<u>Low</u> (lbs/ac)
Alfalfa	9800	4500
Alfalfa/Intermediate wheatgrass	7700	3600
Alfalfa/Smooth bromegrass	7700	3600
Big bluestem	7000	3500
Indiangrass	5100	3200
Intermediate wheatgrass	6000	2900
Smooth bromegrass	6000	2900
Switchgrass	7600	3700

Forage Growth Curves

Growth curves estimate the seasonal distribution of growth of the various forage crops. They indicate when the forages may be available for grazing or mechanical harvest.

Growth Curve Number: SD0001
Growth Curve Name: Alfalfa
Growth Curve Description: Alfalfa, MLRAs 107, 102B, 63B, 66, 65

Percent Production by Month

<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	5	30	25	20	15	5	0	0	0

Growth Curve Number: SD0004
Growth Curve Name: Cool season grass
Growth Curve Description: Cool season grass, statewide

Percent Production by Month

<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	10	40	30	10	5	5	0	0	0

Growth Curve Number: SD0005
Growth Curve Name: Warm season grass
Growth Curve Description: Warm season grass, statewide

Percent Production by Month

<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	0	10	40	35	15	0	0	0	0

Soil Limitations

These soils have few limitations to the production of climatically adapted forage crops. Production potential is high. Flooding is a potential hazard to some of these soils. Also, all of these soils receive additional moisture, so the potential exists for soil compaction from grazing or operating machinery on them when wet.

Management Interpretations

Soils in this group that are subject to flooding can have forage production adversely impacted if it occurs during the spring or growing season. Flooding duration, or the time period plants are under water, is more important than flooding frequency for the survival of forage crops. If these soils flood it is generally for only a brief time. Exclude livestock and machinery during extended periods of soil wetness to reduce soil compaction. When establishing new stands or renovating stands select highly productive species and varieties that can make best use of the additional soil moisture this group receives.

FSG Documentation

Similar FSG's:

<u>FSG ID</u>	<u>FSG Narrative</u>
G055CY100SD	Loamy soils do not receive the additional water and are less productive.
G055CY700SD	Subirrigated soils have elevated water tables that come closer to the surface during part of the growing season.

Inventory Data References

- Agriculture Handbook 296-Land Resource Regions and Major Land Resource Areas
- Natural Resources Conservation Service (NRCS) National Water and Climate Center data
- National Soil Survey Information System (NASIS) for soil surveys in South Dakota counties in MLRA 53C
- NRCS South Dakota Technical Guide
- NRCS National Range and Pasture Handbook
- Various Agricultural Research Service, Cooperative Extension Service, and NRCS research trials for plant adaptation and production

State Correlation

This site has been correlated with the following states: South Dakota

Forage Suitability Group Approval

Original Author: Tim Nordquist

Original Date:

Approval by: Dave Schmidt

Approval Date: 2/7/03