

FORAGE SUITABILITY GROUP LOAM

FSG No.: G063AY100SD

Major Land Resource Area: 63A - Northern Rolling Pierre Shale Plains

Physiographic Features

The soils in this group are found on upland slopes, stream terraces, and flood plains.

	<u>Minimum</u>	<u>Maximum</u>
Elevation (feet):	1300	2950
Slope (percent):	0	15
Flooding:		
Frequency:	None	Rare
Duration:	None	Very Brief
Ponding:		
Depth (inches):		
Frequency:	None	None
Duration:	None	None
Runoff Class:	Low	High

Climatic Features

This group occurs in a mid-continental climate characterized by wide seasonal temperature and precipitation fluctuations and extremes.

Annual precipitation varies widely from year to year in MLRA 63A. Average annual precipitation for all climate stations listed below is about 17 inches. About 77 percent of that occurs during the months of April through September. On average, there are about 25 days with greater than .1 inches of precipitation during that same time period. Precipitation is less than needed for optimum forage production and is the single largest factor limiting production from this group on non-irrigated lands.

Average annual snowfall ranges from 24 inches at Midland to 48 inches at Milesville. Snow cover at depths greater than 1 inch range from 27 days at Midland to 82 days at Timber Lake.

Average July temperatures across the MLRA are about 75⁰F, and average January temperatures are about 17⁰F. Recorded temperature extremes in the MLRA during the years 1961 to 1990 are a low of -37 at Kennebec and a high of 114 at both Kennebec and Midland. The MLRA lies in USDA Plant Hardiness Zones 4a, 4b and 5a.

The climate data listed in the tables below represent high and low ranges and averages for the climate stations and dates listed. For additional climate data, access the National Water and Climate Center at <http://www.wcc.nrcs.usda.gov>.

	From	To
Freeze-free period (28 deg)(days): (9 years in 10 at least)	129	162
Last Killing Freeze in Spring (28 deg): (1 year in 10 later than)	May 20	May 04
Last Frost in Spring (32 deg): (1 year in 10 later than)	May 31	May 16
First Frost in Fall (32 deg): (1 year in 10 earlier than)	Sep 09	Sep 24

	From	To
First Killing Freeze in Fall (28 deg): (1 year in 10 earlier than)	Sep 17	Oct 01
Length of Growing Season (32 deg)(days): (9 years in 10 at least)	110	139
Growing Degree Days (40 deg):	4442	5149
Growing Degree Days (50 deg):	2517	3083
Annual Minimum Temperature:	-30	-15
Mean annual precipitation (inches):	16	18

Monthly precipitation (inches) and temperature (F):

2 years in 10:	<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
Precip. Less Than	0.10	0.09	0.31	0.82	1.44	1.55	0.90	0.64	0.41	0.30	0.08	0.16
Precip. More Than	0.60	0.79	2.37	3.46	3.82	4.55	3.58	2.46	1.98	2.06	1.07	0.91
Monthly Average:	0.30	0.42	1.20	1.99	2.86	3.06	2.23	1.80	1.31	1.12	0.48	0.45
Temp. Min.	3.9	9.7	20.0	32.9	44.0	53.9	59.6	57.3	46.4	35.5	20.8	7.9
Temp. Max.	32.8	38.6	48.3	63.1	74.1	83.8	92.2	90.6	79.3	66.4	48.4	35.9
Temp. Avg.	17.1	22.9	33.0	46.7	58.0	68.0	75.0	73.0	61.7	49.6	33.5	20.5

<u>Climate Station</u>	<u>Location</u>	<u>From</u>	<u>To</u>
SD5691	Mobridge SD	1961	1990
SD8307	Timber Lake SD	1961	1990
SD6170	Oahe Dam SD	1961	1990
SD5506	Midland SD	1961	1990
SD5544	Milesville SD	1961	1990
SD6552	Philip SD	1961	1990
SD5891	Murdo SD	1961	1990
SD4516	Kennebec SD	1961	1990

Soil Interpretations

This group consists of very deep, mostly well drained, medium textured soils formed from wind and water deposited materials. Available water capacity is high and permeability is moderately slow to moderate.

Drainage Class:	Moderately well drained	To	Well drained
Permeability Class: (0 - 40 inches)	Moderately slow	To	Moderate
Frost Action Class:	Low	To	Moderate

	<u>Minimum</u>	<u>Maximum</u>
Depth:	72	
Surface Fragments >3" (% Cover):	0	3
Organic Matter (percent): (surface layer)	0.5	4.0
Electrical Conductivity (mmhos/cm): (0 - 24 inches)	0	8
Sodium Absorption Ratio: (0 - 12 inches)	0	10
Soil Reaction (1:1) Water (pH): (0 - 12 inches)	6.1	8.4
Available Water Capacity (inches): (0 - 60 inches)	9	12
Calcium Carbonate Equivalent (percent): (0 - 12 inches)	0	14

Adapted Species List

The following forage species are considered adapted to grow on the soils in this group. Additional information concerning plant characteristics of a number of the listed species as well as individual cultivars of many of those species can be accessed at <http://plants.usda.gov/>.

Cool Season Grasses

Altai wildrye	G
Canada wildrye	F
Crested wheatgrass	G
Green needlegrass	G
Intermediate wheatgrass	G
Meadow bromegrass	G
Newhy hybrid wheatgrass	G
Pubescent wheatgrass	G
Russian wildrye	G
Slender wheatgrass	G
Smooth bromegrass	G
Tall wheatgrass	G
Western wheatgrass	G

Warm Season Grasses

Big bluestem	G
Indiangrass	G
Little bluestem	G
Prairie sandreed	F
Sand bluestem	F
Sideoats grama	G
Switchgrass	G
Legumes	
Alfalfa	G
Birdsfoot trefoil	F
Canada milkvetch	F
Cicer milkvetch	G
Purple prairieclover	F
Sainfoin	F
Sweetclover	G
White prairieclover	F

G - Good adaptation for forage production on this group of soils in this MLRA

F - Fair adaptation but will not produce at its highest potential

Production Estimates

Production estimates listed here should only be used for making general management recommendations. Onsite production information should always be used for making detailed planning and management recommendations.

The high forage production estimates listed below are based on dense, vigorous stands of climatically adapted, superior performing cultivars. They are properly fertilized for high yields and pest infestations are kept below economic thresholds. Mechanical harvests are managed to maintain stand life by cutting at appropriate stages of maturity and harvest intervals. If grazed, optimum beginning and ending grazing heights are adhered to. Adequate time is allowed for plant recovery before entering winter dormancy under both uses.

The production estimates listed below represent total annual above ground plant production on an air-dry-matter basis. Estimates of hay and grazing yields can be calculated from these numbers by multiplying them by a harvest efficiency. A 70 percent harvest efficiency is commonly used when converting to hay yields. Pasture harvest efficiency is highly dependent on the grazing management system applied, ranging from 25 to 50 percent.

Forage Crop	Management Intensity	
	<u>High</u> (lbs/ac)	<u>Low</u> (lbs/ac)
Alfalfa	6600	2700
Alfalfa/Crested wheatgrass	5400	2300
Alfalfa/Intermediate wheatgrass	5400	2300
Alfalfa/smooth bromegrass	5400	2300
Crested wheatgrass	4600	2200
Green needlegrass	4000	1700
Intermediate wheatgrass	4900	2300
Smooth bromegrass	4900	2300
Western wheatgrass	4000	1500

Forage Growth Curves

Growth curves estimate the seasonal distribution of growth of the various forage crops. They indicate when the

forages may be available for grazing or mechanical harvest.

Growth Curve Number: SD0001
Growth Curve Name: Alfalfa
Growth Curve Description: Alfalfa, MLRA's 107, 102B, 63B, 66, 65

Percent Production by Month											
<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	5	30	25	20	15	5	0	0	0

Growth Curve Number: SD0004
Growth Curve Name: Cool season grass
Growth Curve Description: Cool season grass, statewide

Percent Production by Month											
<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	10	40	30	10	5	5	0	0	0

Growth Curve Number: SD0005
Growth Curve Name: Warm season grass
Growth Curve Description: Warm season grass, statewide

Percent Production by Month											
<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	0	10	40	35	15	0	0	0	0

Growth Curve Number: SD0003
Growth Curve Name: Irrigated Alfalfa
Growth Curve Description: Irrigated Alfalfa, statewide

Percent Production by Month											
<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	5	25	25	20	15	10	0	0	0

Soil Limitations

These soils have few limitations to the production of climatically adapted forage crops. On steeper slopes, water erosion is a potential problem during establishment, when renovating stands, and in thin established stands. Livestock trail erosion is a potential problem in established stands.

Management Interpretations

Including sod forming grass species in stands, especially on steeper slopes, will reduce the potential for sheet and rill erosion. Incorporate erosion control practices during the establishment period. Properly locating facilitating practices such as fences, lanes, and water developments can help control livestock movement, reduce trailing perpendicular to steeper slopes, and evenly distribute grazing pressure.

FSG Documentation

Similar FSG's:

FSG ID	FSG Narrative
G063AY120SD	Droughty Loam soils are shallower or coarser textured resulting in lower available water capacity and lower production potential.
G063AY500SD	Overflow soils receive additional moisture due to a favorable landscape position resulting in a higher production potential.

Inventory Data References

Agriculture Handbook 296-Land Resource Regions and Major Land Resource Areas
Natural Resources Conservation Service (NRCS) National Water and Climate Center data
USDA Plant Hardiness Zone Maps,
National Soil Survey Information System (NASIS) for soil surveys in South Dakota counties in MLRA 63A
South Dakota NRCS South Dakota Technical Guides
NRCS National Range and Pasture Handbook
Various South Dakota Agricultural Research Service, Cooperative Extension Service, and NRCS research trials for plant adaptation and production.

State Correlation

This site has been correlated with the following states: South Dakota

Forage Suitability Group Approval

Original Author: Tim Nordquist
Original Date: 4/5/02
Approval by: Dave Schmidt
Approval Date: 7/15/03