


FORAGE SUITABILITY GROUP (FSG) Overflow

FSG No.: G066XY500NE

Major Land Resource Area (MLRA): 66 - Dakota-Nebraska Eroded Tableland



Physiographic Features

The soils in this group are found on flood plains, low stream terraces, and footslopes.

	<u>Minimum</u>	<u>Maximum</u>
Elevation (feet):	1970	2950
Slope (percent):	0	2
Flooding:		
Frequency:	None	Frequent
Duration:	None	Very Brief
Ponding:		
Depth (inches):		
Frequency:	None	None
Duration:	None	None
Runoff Class:	Low	Medium

Climatic Features

This group occurs in a mid-continental climate characterized by wide seasonal temperature and precipitation fluctuations and extremes.

Annual precipitation varies widely from year to year in MLRA 66. Average annual precipitation for all climate stations listed below is about 21 inches. About 77 percent of the annual precipitation occurs during the months of April through September. On average, there are about 29 days with greater than .1 inches of precipitation during that same time period. Precipitation is less than needed for optimum forage production and is the single largest factor limiting production from this group on non-irrigated lands.

Average annual snowfall ranges from 33 inches at O'Neill, Nebraska (NE), to 43 inches at Harrington, South Dakota (SD). Snow cover at depths greater than 1 inch range from 43 days at Springview, NE, to 64 days at Harrington, SD.

Average July temperatures across the MLRA are about 74°F and average January temperatures are about 20°F. Recorded temperature extremes in the MLRA during the years 1961 to 1990 are a low of -38°F at Harrington and a high of 110 both recorded at Mission, Springview, and O'Neill. The MLRA lies in USDA Plant Hardiness Zones 4b and 5a.

The climate data listed in the tables below represent high and low ranges and averages for the climate stations and dates listed. For additional climate data access the National Water and Climate Center at www.wcc.nrcs.usda.gov.

	From	To
Freeze-free period (28 deg)(days): (9 years in 10 at least)	120	150
Last Killing Freeze in Spring (28 deg): (1 year in 10 later than)	May 23	May 08
Last Frost in Spring (32 deg): (1 year in 10 later than)	Jun 01	May 20
First Frost in Fall (32 deg): (1 year in 10 earlier than)	Sep 07	Sep 17

	From	To
First Killing Freeze in Fall (28 deg): (1 year in 10 earlier than)	Sep 11	Sep 26
Length of Growing Season (32 deg)(days): (9 years in 10 at least)	104	130
Growing Degree Days (40 deg):	4580	5148
Growing Degree Days (50 deg):	2615	3038
Annual Minimum Temperature:	-25	-15
Mean annual precipitation (inches):	18	25

Monthly precipitation (inches) and temperature (F):

2 years in 10:	<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
Precip. Less Than	0.08	0.11	0.33	0.62	1.70	1.51	1.54	0.91	0.65	0.53	0.15	0.16
Precip. More Than	0.54	1.24	2.70	3.97	5.70	5.65	4.96	3.94	4.34	2.64	1.49	0.85
Monthly Average:	0.34	0.49	1.42	2.16	3.40	3.46	3.07	2.22	2.15	1.32	0.71	0.52
Temp. Min.	7.1	12.4	20.8	31.7	42.4	52.3	58.5	55.8	45.2	33.5	20.7	10.0
Temp. Max.	30.3	36.2	46.8	62.1	72.9	82.8	88.7	86.5	76.2	64.5	46.4	33.3
Temp. Avg.	19.7	24.8	34.1	47.1	57.9	68.0	74.2	72.1	61.8	49.8	34.7	22.9

<u>Climate Station</u>	<u>Location</u>	<u>From</u>	<u>To</u>
SD3574	Harrington SD	1961	1990
SD5620	Mission SD	1966	1990
SD5638	Mission SD	1961	1990
SD0778	Bonesteel SD	1961	1990
NE8090	Springview NE	1961	1990
NE6290	O'Neill NE	1961	1990

Soil Interpretations

This group consists of very deep, well drained, medium to fine textured soils formed from alluvium. Permeability is slow to moderate, and available water capacity is high.

Drainage Class:	Well drained	To	Well drained
Permeability Class: (0 - 40 inches)	Slow	To	Moderate
Frost Action Class:	Moderate	To	Moderate

	<u>Minimum</u>	<u>Maximum</u>
Depth:	72	
Surface Fragments >3" (% Cover):	0	3
Organic Matter (percent): (surface layer)	1.0	6.0
Electrical Conductivity (mmhos/cm): (0 - 24 inches)	0	2
Sodium Absorption Ratio: (0 - 12 inches)	0	0
Soil Reaction (1:1) Water (pH): (0 - 12 inches)	5.6	8.4
Available Water Capacity (inches): (0 - 60 inches)	9	12
Calcium Carbonate Equivalent (percent): (0 - 12 inches)	0	8

Soil Component List (Some phases of these soils may also occur in other FSG's)

Bridgeport Onita

Adapted Species List

The following forage species are considered adapted to grow on the soils in this group. Additional information concerning plant characteristics of a number of the listed species as well as individual cultivars of many of those species can be accessed on the web at <http://plants.usda.gov/>.

<u>Cool Season Grasses</u>	<u>Symbol</u>	<u>Dry</u>	<u>Irrig</u>	<u>Legumes</u>	<u>Symbol</u>	<u>Dry</u>	<u>Irrig</u>
Green needlegrass	NAVI4	G	NS	Alfalfa	MESA	G	G
Intermediate wheatgrass	THIN6	G	G	Birdsfoot trefoil	LOCO6	G	G
Meadow brome	BRBI2	G	G	Canada milkvetch	ASCAC6	G	NS
Orchardgrass	DAGL	G	G	Cicer milkvetch	ASCI4	G	F
Pubescent wheatgrass	THIN6	G	G	Illinois bundleflower	DEIL	G	NS
Smooth brome	BRINI2	G	G	Purple prairieclover	DAPUP	F	NS
Tall wheatgrass	THPO7	G	NS	Red clover	TRPR2	G	G
Western wheatgrass	PASM	G	NS	White prairieclover	DACAC	F	NS
<u>Warm Season Grasses</u>	<u>Symbol</u>	<u>Dry</u>	<u>Irrig</u>				
Big bluestem	ANGE	G	G				
Indiangrass	SONU2	G	G				
Little bluestem	SCSC	G	NS				
Prairie sandreed	CALO	F	NS				
Sand bluestem	ANHA	F	NS				
Sideoats grama	BOCU	F	NS				
Switchgrass	PAVIV	G	G				

G - Good adaptation for forage production on this group of soils in this MLRA
 F - Fair adaptation but will not produce at its highest potential
 NS - Species is not adapted to the site and should not be planted

Production Estimates

Production estimates listed here should only be used for making general management recommendations. Onsite production information should always be used for making detailed planning and management recommendations.

The high forage production estimates listed below are based on dense, vigorous stands of climatically adapted, superior performing cultivars. They are properly fertilized for high yields, and pest infestations are kept below economic thresholds. Mechanical harvests are managed to maintain stand life by cutting at appropriate stages of maturity and harvest intervals. If grazed, optimum beginning and ending grazing heights are adhered to. Adequate time is allowed for plant recovery before entering winter dormancy under both uses.

The production estimates listed below represent total annual above ground plant production on an air-dry-matter basis. Estimates of hay and grazing yields can be calculated from these numbers by multiplying them by a harvest efficiency. A 70 percent harvest efficiency is commonly used when converting to hay yields. Pasture harvest efficiency is highly dependent on the grazing management system applied, ranging from 25 to 50 percent.

Forage Crop	<u>Dryland</u>		<u>Irrigated</u>	
	Management Intensity		Management Intensity	
	<u>High</u> (lbs/ac)	<u>Low</u> (lbs/ac)	<u>High</u> (lbs/ac)	<u>Low</u> (lbs/ac)
Alfalfa	10000	4000		
Alfalfa/Intermediate wheatgrass	9700	4000	17100	10300
Alfalfa/Orchardgrass	9700	4000	17100	10300
Alfalfa/Smooth brome grass	9700	4000	17100	10300
Big bluestem	7400	4000		
Intermediate wheatgrass	6300	3100	14300	8600
Orchardgrass	5700	3100	14300	8600
Smooth brome grass	5700	3100	14300	8600
Switchgrass	7400	4000		

Forage Growth Curves

Growth curves estimate the seasonal distribution of growth of the various forage crops. They indicate when the forages may be available for grazing or mechanical harvest.

Growth Curve Number: SD0001
Growth Curve Name: Alfalfa
Growth Curve Description: Alfalfa, MLRA's 107, 102B, 63B, 66, 65

Percent Production by Month

<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	5	30	25	20	15	5	0	0	0

Growth Curve Number: SD0004
Growth Curve Name: Cool season grass
Growth Curve Description: Cool season grass, statewide

Percent Production by Month

<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	10	40	30	10	5	5	0	0	0

Growth Curve Number: SD0005
Growth Curve Name: Warm season grass
Growth Curve Description: Warm season grass, statewide

Percent Production by Month

<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	0	10	40	35	15	0	0	0	0

Growth Curve Number: SD0003
Growth Curve Name: Irrigated Alfalfa
Growth Curve Description: Irrigated Alfalfa, statewide

Percent Production by Month

<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	5	25	25	20	15	10	0	0	0

Soil Limitations

These soils have few limitations to the production of climatically adapted forage crops. Production potential is high. Flooding is a potential hazard to some of these soils. Also, all of these soils receive additional moisture, so the potential exists for soil compaction from grazing or operating machinery on them when wet.

Management Interpretations

Soils in this group that are subject to flooding can have forage production adversely impacted if it occurs during the spring or growing season. Flooding duration or the time period plants are under water is more important than flooding frequency for the survival of forage crops. If these soils flood it is generally for only a very brief time. Exclude livestock and machinery during extended periods of soil wetness to reduce soil compaction. When establishing new stands or renovating stands select species and varieties that can make best use of the additional soil moisture this group receives.

FSG Documentation

Similar FSG's:

FSG ID

G066XY100N

FSG Narrative

Loamy soils do not receive the additional water and are less productive.

G066XY700N

Subirrigated soils have elevated watertables that come closer to the surface during part of the growing season.

Inventory Data References:

Agriculture Handbook 296-Land Resource Regions and Major Land Resource Areas

Natural Resources Conservation Service (NRCS) National Water and Climate Center data

USDA Plant Hardiness Zone Maps

National Soil Survey Information System for soil surveys in South Dakota and Nebraska counties in MLRA 66

NRCS Nebraska Field Office Technical Guide and South Dakota Technical Guide

NRCS National Range and Pasture Handbook

Various South Dakota and Nebraska Agricultural Research Service, Cooperative Extension Service, and NRCS research trials for plant adaptation and production.

State Correlation:

This site has been correlated with the following states: Nebraska and South Dakota

Forage Suitability Group Approval:

Original Author: Tim Nordquist

Original Date: 4/4/02

Approval by: Dave Schmidt

Approval Date: 9/20/04