

FORAGE SUITABILITY GROUP

Saline

FSG No.: G102AY895SD

Major Land Resource Area: 102A - Rolling Till Prairie

Physiographic Features

These soils are found on level and nearly level glacial lake plains, flood plains, and till plains.

	<u>Minimum</u>	<u>Maximum</u>
Elevation (feet):	980	1970
Slope (percent):	0	2
Flooding:		
Frequency:	Rare	Frequent
Duration:	Very Brief	Long
Ponding:		
Depth (inches):		
Frequency:	None	None
Duration:	None	None
Runoff Class:	Negligible	Medium

Climatic Features

This group occurs in a mid-continental climate characterized by wide seasonal temperature and precipitation fluctuations and extremes.

Annual precipitation varies widely from year to year in MLRA 102A. Average annual precipitation for all climate stations listed below is about 23 inches. About 75 percent of that occurs during the months of April through September. On average, there are about 31 days with greater than .1 inches of precipitation during the same timeframe.

Average annual snowfall ranges from 36 inches at Britton to 48 inches at Tracy. Snow cover at depths greater than 1 inch range from 56 days at Milbank to 105 days at Morris.

Average July temperatures are about 72⁰F and average January temperatures are about 11⁰F. Recorded temperature extremes in the MLRA during the years 1961 to 1990 are a low of -40 at Brookings and a high of 108 recorded at both Britton and Milbank. The MLRA lies in USDA Plant Hardiness Zones 4a and 4b.

Average annual wind speeds range from about 8 mph in the eastern part of the MLRA to about 11 mph in the west. The highest wind speeds occur during March through May. It is cloudy about 154 days a year in the west and 166 days in the east. Average morning relative humidity in June is about 86 percent and average afternoon humidity is 59 percent.

The climate data listed in the tables below represent high and low ranges and averages for the climate stations and dates listed. For additional climate data, access the National Water and Climate Center at <http://www.wcc.nrcs.usda.gov>.

	From	To
Freeze-free period (28 deg)(days): (9 years in 10 at least)	127	145
Last Killing Freeze in Spring (28 deg): (1 year in 10 later than)	May 22	May 11
Last Frost in Spring (32 deg): (1 year in 10 later than)	May 31	May 17

	From	To
First Frost in Fall (32 deg): (1 year in 10 earlier than)	Sep 08	Sep 19
First Killing Freeze in Fall (28 deg): (1 year in 10 earlier than)	Sep 17	Sep 26
Length of Growing Season (32 deg)(days): (9 years in 10 at least)	109	134
Growing Degree Days (40 deg):	4066	4515
Growing Degree Days (50 deg):	2441	2698
Annual Minimum Temperature:	-30	-20
Mean annual precipitation (inches):	19	26

Monthly precipitation (inches) and temperature (F):

2 years in 10:	<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
Precip. Less Than	0.13	0.19	0.28	0.66	1.29	1.83	1.54	0.91	0.68	0.45	0.17	0.10
Precip. More Than	0.97	1.08	2.70	3.68	4.83	4.92	5.21	3.75	4.63	3.32	2.19	1.19
Monthly Average:	0.54	0.59	1.37	2.20	2.88	3.67	3.21	2.77	2.32	1.83	0.96	0.54
Temp. Min.	-2.8	3.1	17.6	32.8	44.6	54.6	59.3	56.2	45.7	34.2	20.5	4.6
Temp. Max.	21.4	26.5	39.3	56.5	70.4	80.5	85.5	82.9	73.2	61.0	42.0	26.6
Temp. Avg.	10.1	15.9	29.0	44.6	57.2	66.8	72.0	69.5	59.3	47.5	30.8	15.6

<u>Climate Station</u>	<u>Location</u>	<u>From</u>	<u>To</u>
SD1049	Britton, SD	1961	1990
SD1076	Brookings, SD	1961	1990
SD1739	Clark, SD	1961	1990
SD1777	Clear Lake, SD	1961	1990
SD5536	Milbank, SD	1961	1990
MN5400	Milan, MN	1961	1990
MN5638	Morris, MN	1961	1990
MN8323	Tracy, MN	1961	1990

Soil Interpretations

This group consists of poorly drained soils with elevated salinity levels.

Drainage Class:	Poorly drained	To	Poorly drained
Permeability Class: (0 - 40 inches)	Moderate	To	Slow
Frost Action Class:	High	To	High

	<u>Minimum</u>	<u>Maximum</u>
Depth:	72	
Surface Fragments >3" (% Cover):	0	3
Organic Matter (percent): (surface layer)	4.0	9.0
Electrical Conductivity (mmhos/cm): (0 - 24 inches)	8	16
Sodium Absorption Ratio: (0 - 12 inches)	2	7
Soil Reaction (1:1) Water (pH): (0 - 12 inches)	6.1	9
Available Water Capacity (inches): (0 - 60 inches)	5	10
Calcium Carbonate Equivalent (percent): (0 - 12 inches)	0	23

Adapted Species List

The following forage species are considered adapted to grow on the soils in this group. Additional information concerning plant characteristics of a number of the listed species as well as individual cultivars of many those species can be accessed at <http://plants.usda.gov>.

Cool Season Grasses

Beardless wildrye	G
Creeping foxtail	F
Intermediate wheatgrass	F
Newhy hybrid wheatgrass	G
Pubescent wheatgrass	F
Reed canarygrass	F
Slender wheatgrass	G
Tall fescue	G
Tall wheatgrass	G
Western wheatgrass	G

Warm Season Grasses

Alkali sacaton	F
Switchgrass	F

Legumes

Alfalfa	F
Alsike clover	F
Birdsfoot trefoil	F
Cicer milkvetch	F

G - Good adaptation for forage production on this group of soils in this MLRA
F - Fair adaptation but will not produce at its highest potential

Production Estimates

Production estimates listed here should only be used for making general management recommendations. Onsite production information should always be used for making detailed planning and management recommendations.

The high forage production estimates listed below are based on dense, vigorous stands of climatically adapted, superior performing cultivars. They are properly fertilized for high yields and pest infestations are kept below economic thresholds. Mechanical harvests are managed to maintain stand life by cutting at appropriate stages of maturity and harvest intervals. If grazed, optimum beginning and ending grazing heights are adhered to. Adequate time is allowed for plant recovery before entering winter dormancy under both uses.

The production estimates listed below represent total annual above ground plant production on an air-dry-matter basis. Estimates of hay and grazing yields can be calculated from these numbers by multiplying them by a harvest efficiency. A 70 percent harvest efficiency is commonly used when converting to hay yields. Pasture harvest efficiency is highly dependent on the grazing management system applied, ranging from 25 to 50 percent.

Forage Crop	Management Intensity	
	<u>High</u> (lbs/ac)	<u>Low</u> (lbs/ac)
Switchgrass	3400	2000
Tall wheatgrass	5700	3400
Western wheatgrass	3400	2000

Forage Growth Curves

Growth curves estimate the seasonal distribution of growth of the various forage crops. They indicate when the forages may be available for grazing or mechanical harvest.

Growth Curve Number: ND0001
Growth Curve Name: Alfalfa
Growth Curve Description: Alfalfa

Percent Production by Month											
<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	5	25	30	20	15	5	0	0	0

Growth Curve Number: ND0002
Growth Curve Name: Cool season grass
Growth Curve Description: Cool season grass

Percent Production by Month

<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	5	40	35	10	5	5	0	0	0

Growth Curve Number: ND0003
Growth Curve Name: Warm season grass
Growth Curve Description: Warm season grass

Percent Production by Month

<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	0	10	40	35	15	0	0	0	0

Soil Limitations

These soils have severe limitations to the production of climatically adapted forage species. Species selection and productivity are severely limited by the high salinity levels. Also, these soils are poorly drained and will experience periods when trafficability will be difficult or impossible. These soils are subject to compaction if grazed or machinery is operated on them when wet. Drainage also limits species selection.

Management Interpretations

When establishing new stands or renovating stands select species that are tolerant of elevated salinity and sodium levels and that are tolerant of poorly drained soils. Exclude livestock and machinery during extended periods of soil wetness to reduce soil compaction.

FSG Documentation

Similar FSGs:

<u>FSG ID</u>	<u>FSG Narrative</u>
G102AY700S	Subirrigated soils do not have restrictive levels of salinity and/or sodium.
G102AY800S	Claypan soils have better drainage.
G102AY900S	Wet soils do not have restrictive levels of salinity and/or sodium.

Inventory Data References

Agriculture Handbook 296-Land Resource Regions and Major Land Resource Areas
 Natural Resources Conservation Service (NRCS) National Water and Climate Center data
 USDA Plant Hardiness Zone Maps
 National Soil Survey Information System (NASIS) for soil surveys in South Dakota and Minnesota counties in MLRA 102A
 South Dakota NRCS SDTG and Minnesota NRCS FOTG
 NRCS National Range and Pasture Handbook
 Various Agricultural Research Service, Cooperative Extension Service, and NRCS research trials for plant adaptation and production.

State Correlation

This site has been correlated with the following states: Minnesota and South Dakota

Forage Suitability Group Approval

Original Author: Tim Nordquist
Original Date: 1/31/02
Approval By: Dave Schmidt
Approval Date: 1/7/03