FORAGE SUITABILITY GROUP (FSG)

Very Droughty Loam



FSG No.: G102CY130NE

Major Land Resource Area (MLRA): 102C - Loess Uplands

Physiographic Features

The soils in this group are found on uplands, terraces, flood plains, and outwash plains.

	Minimum	Maximum
Elevation (feet):	980	1640
Slope (percent):	0	17
Flooding:		
Frequency:	None	Occasional
Duration:	None	Very Brief
Ponding:		
Depth (inches):		
Frequency:	None	None
Duration:	None	None
Runoff Class:	Very low	High

Climatic Features

Annual precipitation varies widely from year to year in MLRA 102C. Average annual precipitation for all climate stations listed below is about 27 inches. About 73 percent of the annual precipitation occurs during the months of April through September. On average, there are about 33 days with greater than .1 inches of precipitation during the same time period. Annual precipitation and temperature increase from the northwest to the southeast in the MLRA.

Average annual snowfall ranges from 19 inches at Creighton, to 36 inches at Wakefield. Days with snow cover at depths greater than 1 inch range from 9 days at Creighton to 55 days at Wakefield.

Average July temperatures are about 76°F and average January temperatures are about 20°F. Recorded temperature extremes in the MLRA during the years 1961 to 1990 are a low of -36 at Creighton and a high of 108 recorded at both Columbus and Wakefield. The MLRA lies mostly in USDA Plant Hardiness Zone 4b with some small areas of warmer 5a.

At Norfolk, Nebraska (NE), the average annual wind speeds are about 11.2 MPH. The highest wind speeds occur during March and April. It is cloudy about 146 days a year. Average morning relative humidity in June is about 82 percent and average afternoon humidity is 55 percent.

The climate data listed in the tables below represent high and low ranges and averages for the climate stations and dates listed. For additional climate data, access the National Water and Climate Center at http://www.wcc.nrcs.usda.gov.

	From	To
Freeze-free period (28 deg) (days): **(9 years in 10 at least)**	138	168
Last Killing Freeze in Spring (28 deg): **(1 year in 10 later than)**	May 12	Apr 25
Last Frost in Spring (32 deg): **(1 year in 10 later than)**	May 20	May 10
First Frost in Fall (32 deg): **(1 year in 10 earlier than)**	Sep 12	Sep 23

	From	To
First Killing Freeze in Fall (28 deg):	Sep 20	Oct 04
(1 year in 10 earlier than)		
Length of Growing Season (32 deg) (days):	126	146
(9 years in 10 at least)		
Growing Degree Days (40 deg):	4833	5730
Growing Degree Days (50 deg):	2815	3551
Annual Minimum Temperature:	-25	-15
Mean annual precipitation (inches):	23	30

Monthly precipitation (inches) and temperature (F):

2 years in 10:	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Precip. Less Than	0.15	0.17	0.51	0.83	1.92	1.98	1.71	1.35	1.07	0.44	0.15	0.30
Precip. More Than	1.13	1.20	3.39	3.93	5.93	6.63	4.33	5.39	5.53	3.85	2.16	1.47
Monthly Average:	0.55	0.76	2.04	2.53	4.03	4.20	3.09	3.03	3.00	2.02	1.18	0.86
Temp. Min.	5.3	10.6	23.0	35.4	46.5	56.8	61.9	58.7	48.4	35.6	23.8	10.7
Temp. Max.	32.4	38.2	50.2	65.2	75.6	84.8	88.7	86.1	77.7	66.5	49.7	35.3
Temp. Avg.	19.7	25.3	36.8	50.6	61.6	71.3	75.9	73.1	63.9	52.3	37.1	23.5

Climate Station	Location	From	To
NE1825	Columbus, NE	1961	1990
NE1990	Creighton, NE	1961	1990
NE3050	Fremont, NE	1961	1990
NE6018	NE Nebraska Experiment	1964	1990
NE8110	Stanton, NE	1961	1990
NE8480	Tekamah, NE	1961	1990
NE8915	Wakefield, NE	1961	1990
NE8935	Walthill, NE	1961	1990

Soil Interpretations

This group consists of well to excessively drained, medium to coarse textured soils. They have low available water capacity due to moderate to shallow depths to sand and gravel.

Drainage Class:	Well drained	To	Excessively drained
Permeability Class:	Moderate	To	Rapid
(0 - 40 inches)			
Frost Action Class:	Low	To	Moderate

	Minimum	Maximum
Depth:	60	
Surface Fragments >3" (% Cover):		
Organic Matter (percent):	0.5	3.0
(surface layer)		
Electrical Conductivity (mmhos/cm):	0	0
(0 - 24 inches)		
Sodium Absorption Ratio:	0	0
(0 - 12 inches)		
Soil Reaction (1:1) Water (pH):	5.6	7.8
(0 - 12 inches)		
Available Water Capacity (inches):	3	6
(0 - 60 inches)		
Calcium Carbonate Equivalent (percent):	0	15
(0 - 12 inches)		

Soil Component List **(some phases of these soils may also occur in other FSG's)**

Delmont	O'neill	Talmo
Inavale	Simeon	Thurman

Adapted Species List

The following forage species are considered adapted to grow on the soils in this group. Additional information concerning plant characteristics of a number of the listed species as well as individual cultivars of many of those species can be accessed on the web at http://plants.usda.gov.

Cool Season Grasses	Symbol	Dry	Irrig
Green needlegrass	NAVI4	G	NS
Intermediate wheatgrass	THIN6	F	G
Meadow bromegrass	BRBI2	F	G
Orchardgrass	DAGL	F	G
Pubescent wheatgrass	THIN6	G	G
Smooth bromegrass	BRINI2	F	G
Tall fescue	LOAR10	NS	F
Western wheatgrass	PASM	G	NS

Warm Season Grasses	Symbol	Dry	Irrig
Big bluestem	ANGE	F	G
Indiangrass	SONU2	NS	G
Little bluestem	SCSC	G	NS
Sideoats grama	BOCU	G	NS
Switchgrass	PAVIV	NS	G
Legumes	**Symbol**	**Dry**	**Irrig**
Alfalfa	MESA	G	G
Birdsfoot trefoil	LOCO6	NS	G
Cicer milkvetch	ASCI4	G	F
Purple prairieclover	DAPUP	G	NS
Red clover	TRPR2	NS	G
White prairieclover	DACAC	F	NS

G - Good adaptation for forage production on this group of soils in this MLRA
F - Fair adaptation but will not produce at its highest potential
NS - Species is not adapted to the site and should not be planted

Production Estimates

Production estimates listed here should only be used for making general management recommendations. Onsite production information should always be used for making detailed planning and management

Forage Crop	Dryland Management Intensity		Irrigated Management Intensity	
	Low (lbs/ac)	High (lbs/ac)	Low (lbs/ac)	High (lbs/ac)
Alfalfa	2900	5100		
Alfalfa/Intermediate wheatgrass	2600	4300	8600	14300
Alfalfa/Smooth bromegrass	2600	4300	8600	14300
Intermediate wheatgrass	2000	3700	6900	11400
Smooth bromegrass	2000	3700	6900	11400

Production Estimates

Production estimates listed here should only be used for making general management recommendations. Onsite production information should always be used for making detailed planning and management recommendations.

The high forage production estimates listed below are based on dense, vigorous stands of climatically adapted, superior performing cultivars. They are properly fertilized for high yields and pest infestations are kept below economic thresholds. Mechanical harvests are managed to maintain stand life by cutting at appropriate stages of maturity and harvest intervals. If grazed, optimum beginning and ending grazing heights are adhered to. Adequate time is allowed for plant recovery before entering winter dormancy under both uses.

The production estimates listed below represent total annual above ground plant production on an air-dry-matter basis. Estimates of hay and grazing yields can be calculated from these numbers by multiplying them by a harvest efficiency. Seventy percent harvest efficiency is commonly used when converting to hay yields. Pasture harvest efficiency is highly dependent on the grazing management system applied, ranging from 25 to 50 percent.

Growth Curve Number: SD0001
Growth Curve Name: Alfalfa
Growth Curve Description: Alfalfa, MLRA's 102B, 102C, 63B, 66, 65

Percent Production by Month

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	0	5	30	25	20	15	5	0	0	0

Growth Curve Number: SD0004
Growth Curve Name: Cool season grass
Growth Curve Description: Cool season grass, statewide

Percent Production by Month

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	0	10	40	30	10	5	5	0	0	0

Growth Curve Number: SD0005
Growth Curve Name: Warm season grass
Growth Curve Description: Warm season grass, statewide

Percent Production by Month

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	0	0	10	40	35	15	0	0	0	0

Growth Curve Number: SD0003
Growth Curve Name: Irrigated Alfalfa
Growth Curve Description: Irrigated Alfalfa, statewide

Percent Production by Month

Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
0	0	0	5	25	25	20	15	10	0	0	0

Soil Limitations

The primary limitation for these soils is their low available water capacity which limits species selection and production potential. On steeper slopes, water erosion is a potential problem during stand establishment and livestock trail erosion is a potential problem on established stands. Also, wind erosion is a potential problem during stand establishment on moderately coarse and coarse textured soils.

Management Interpretations

The impact on yields of the low available water capacity of these soils can be reduced by selecting forage species that are highly tolerant to periods of drought and inadequate soil moisture. Including sod forming grass species in stands, especially on steeper slopes, will reduce the potential for sheet and rill erosion. Incorporate both wind and water erosion control practices during the establishment period. Properly locating facilitating practices such as fences, lanes, and water developments can help control livestock movement, reduce trailing perpendicular to steeper slopes, and evenly distribute grazing pressure.

FSG Documentation

Similar FSG's:

FSG ID	FSG Narrative
G102CY120NE	Droughty Loam soils have higher available water capacity and greater production potential.

Inventory Data References:

Agriculture Handbook 296-Land Resource Regions and Major Land Resource Areas
Natural Resources Conservation Service (NRCS) National Water and Climate Center data
USDA Plant Hardiness Zone maps
National Soil Survey Information System for soil surveys in Nebraska and South Dakota counties in MLRA 102C
NRCS Nebraska Field Office Technical Guide and South Dakota Technical Guide
NRCS National Range and Pasture Handbook
Various Agricultural Research Service, Cooperative Extension Service, and NRCS research trials for plant adaptation and production.

State Correlation:

This site has been correlated with the following states: Nebraska and South Dakota

Forage Suitability Group Approval:

Original Author:	Tim Nordquist
Original Date:	6/8/2001
Approval by:	Dave Schmidt
Approval Date:	1/13/05