

FORAGE SUITABILITY GROUP (FSG)

Sand


FSG No.: G102CY300NE

Major Land Resource Area (MLRA): 102C - Loess Uplands

Physiographic Features

The soils in this group are found on uplands, stream terraces, and flood plains.

	<u>Minimum</u>	<u>Maximum</u>
Elevation (feet):	980	1640
Slope (percent):	0	17
Flooding:		
Frequency:	None	Frequent
Duration:	None	Long
Ponding:		
Depth (inches):		
Frequency:	None	None
Duration:	None	None
Runoff Class:	Very low	Medium



Climatic Features

Annual precipitation varies widely from year to year in MLRA 102C. Average annual precipitation for all climate stations listed below is about 27 inches. About 73 percent of the annual precipitation occurs during the months of April through September. On average, there are about 33 days with greater than .1 inches of precipitation during the same time period. Annual precipitation and temperature increase from the northwest to the southeast in the MLRA. Average annual snowfall ranges from 19 inches at Creighton, to 36 inches at Wakefield. Days with snow cover at depths greater than 1 inch range from 9 days at Creighton to 55 days at Wakefield.

Average July temperatures are about 76°F and average January temperatures are about 20°F. Recorded temperature extremes in the MLRA during the years 1961 to 1990 are a low of -36 at Creighton and a high of 108 recorded at both Columbus and Wakefield. The MLRA lies mostly in USDA Plant Hardiness Zone 4b with some small areas of warmer 5a.

At Norfolk, Nebraska (NE), the average annual wind speeds are about 11.2 MPH. The highest wind speeds occur during March and April. It is cloudy about 146 days a year. Average morning relative humidity in June is about 82 percent and average afternoon humidity is 55 percent.

The climate data listed in the tables below represent high and low ranges and averages for the climate stations and dates listed. For additional climate data, access the National Water and Climate Center at <http://www.wcc.nrcs.usda.gov>.

	From	To
Freeze-free period (28 deg) (days): (9 years in 10 at least)	138	168
Last Killing Freeze in Spring (28 deg): (1 year in 10 later than)	May 12	Apr 25
Last Frost in Spring (32 deg): (1 year in 10 later than)	May 20	May 10
First Frost in Fall (32 deg): (1 year in 10 earlier than)	Sep 12	Sep 23

	From	To
First Killing Freeze in Fall (28 deg): (1 year in 10 earlier than)	Sep 20	Oct 04
Length of Growing Season (32 deg) (days): (9 years in 10 at least)	126	146
Growing Degree Days (40 deg):	4833	5730
Growing Degree Days (50 deg):	2815	3551
Annual Minimum Temperature:	-25	-15
Mean annual precipitation (inches):	23	30

Monthly precipitation (inches) and temperature (F):

2 years in 10:	<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
Precip. Less Than	0.15	0.17	0.51	0.83	1.92	1.98	1.71	1.35	1.07	0.44	0.15	0.30
Precip. More Than	1.13	1.20	3.39	3.93	5.93	6.63	4.33	5.39	5.53	3.85	2.16	1.47
Monthly Average:	0.55	0.76	2.04	2.53	4.03	4.20	3.09	3.03	3.00	2.02	1.18	0.86
Temp. Min.	5.3	10.6	23.0	35.4	46.5	56.8	61.9	58.7	48.4	35.6	23.8	10.7
Temp. Max.	32.4	38.2	50.2	65.2	75.6	84.8	88.7	86.1	77.7	66.5	49.7	35.3
Temp. Avg.	19.7	25.3	36.8	50.6	61.6	71.3	75.9	73.1	63.9	52.3	37.1	23.5

<u>Climate Station</u>	<u>Location</u>	<u>From</u>	<u>To</u>
NE1825	Columbus, NE	1961	1990
NE1990	Creighton, NE	1961	1990
NE3050	Fremont, NE	1961	1990
NE6018	NE Nebraska Experiment	1964	1990
NE8110	Stanton, NE	1961	1990
NE8480	Tekamah, NE	1961	1990
NE8915	Wakefield, NE	1961	1990
NE8935	Walthill, NE	1961	1990

Soil Interpretations

This group consists of moderately well to excessively drained, coarse textured soils formed in sandy eolian and alluvial materials. Permeability is moderately slow to rapid. Those that are moderately slowly and moderately permeable are sand overlying finer textured subsoil.

Drainage Class:	Moderately well drained	To	Excessively drained
Permeability Class: (0 - 40 inches)	Moderately slow	To	Rapid
Frost Action Class:	Low	To	High

	<u>Minimum</u>	<u>Maximum</u>
Depth:	60	
Surface Fragments >3" (% Cover):		
Organic Matter (percent): (surface layer)	0.5	3.0
Electrical Conductivity (mmhos/cm): (0 - 24 inches)	0	0
Sodium Absorption Ratio: (0 - 12 inches)	0	0
Soil Reaction (1:1) Water (pH): (0 - 12 inches)	5.1	8.4
Available Water Capacity (inches): (0 - 60 inches)	3	10
Calcium Carbonate Equivalent (percent): (0 - 12 inches)	0	10

Soil Component List (some phases of these soils may also occur in other FSG's)

Bazile	Ipage	Thurman
Boelus	Sardak	Ticonic
Inavale	Sarpy	Valentine

Adapted Species List

The following forage species are considered adapted to grow on the soils in this group. Additional information concerning plant characteristics of a number of the listed species as well as individual cultivars of many of those species can be accessed on the web at <http://plants.usda.gov>.

Cool Season Grasses	<u>Symbol</u>	<u>Dry</u>	<u>Irrig</u>	Legumes	<u>Symbol</u>	<u>Dry</u>	<u>Irrig</u>
Intermediate wheatgrass	THIN6	F	G	Alfalfa	MESA	G	G
Meadow bromegrass	BRBI2	NS	G	Cicer milkvetch	ASCI4	F	F
Orchardgrass	DAGL	NS	G	Purple prairieclover	DAPUP	F	NS
Pubescent wheatgrass	THIN6	F	G	Red clover	TRPR2	NS	G
Smooth bromegrass	BRINI2	F	G				
Tall fescue	LOAR10	F	F				
Western wheatgrass	PASM	F	NS				
Warm Season Grasses	<u>Symbol</u>	<u>Dry</u>	<u>Irrig</u>				
Big bluestem	ANGE	G	G				
Indiangrass	SONU2	G	G				
Little bluestem	SCSC	G	NS				
Prairie sandreed	CALO	G	NS				
Sand bluestem	ANHA	G	NS				
Sand lovegrass	ERTR3	F	NS				
Sideoats grama	BOCU	F	NS				
Switchgrass	PAVIV	G	G				

G - Good adaptation for forage production on this group of soils in this MLRA

F - Fair adaptation but will not produce at its highest potential

NS - Species is not adapted to the site and should not be planted

Production Estimates

Production estimates listed here should only be used for making general management recommendations. Onsite production information should always be used for making detailed planning and management recommendations.

The high forage production estimates listed below are based on dense, vigorous stands of climatically adapted, superior performing cultivars. They are properly fertilized for high yields and pest infestations are kept below economic thresholds. Mechanical harvests are managed to maintain stand life by cutting at appropriate stages of maturity and harvest intervals. If grazed, optimum beginning and ending grazing heights are adhered to. Adequate time is allowed for plant recovery before entering winter dormancy under both uses.

The production estimates listed below represent total annual above ground plant production on an air-dry-matter basis. Estimates of hay and grazing yields can be calculated from these numbers by multiplying them by a harvest efficiency. Seventy percent harvest efficiency is commonly used when converting to hay yields. Pasture harvest efficiency is highly dependent on the grazing management system applied, ranging from 25 to 50 percent.

Forage Crop	<u>Dryland</u>		<u>Irrigated</u>	
	<u>Management Intensity</u>	<u>Management Intensity</u>	<u>Management Intensity</u>	<u>Management Intensity</u>
	<u>Low</u>	<u>High</u>	<u>Low</u>	<u>High</u>
	(lbs/ac)	(lbs/ac)	(lbs/ac)	(lbs/ac)
Alfalfa	4300	6600		
Alfalfa/Intermediate wheatgrass	3100	5100	8600	14300
Alfalfa/Smooth bromegrass	3100	5100	8600	14300
Indiangrass	2900	4800		
Intermediate wheatgrass	2600	4300	6900	11400
Sand bluestem	3700	6300		
Smooth bromegrass	2600	4300	6900	11400

Forage Growth Curves

Growth curves estimate the seasonal distribution of growth of the various forage crops. They indicate when the forages may be available for grazing or mechanical harvest.

Growth Curve Number: SD0001
Growth Curve Name: Alfalfa
Growth Curve Description: Alfalfa, MLRA's 102B, 102C, 63B, 66, 65

Percent Production by Month

<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	5	30	25	20	15	5	0	0	0

Growth Curve Number: SD0004
Growth Curve Name: Cool season grass
Growth Curve Description: Cool season grass, statewide

Percent Production by Month

<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	10	40	30	10	5	5	0	0	0

Growth Curve Number: SD0005
Growth Curve Name: Warm season grass
Growth Curve Description: Warm season grass, statewide

Percent Production by Month

<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	0	10	40	35	15	0	0	0	0

Growth Curve Number: SD0003
Growth Curve Name: Irrigated Alfalfa
Growth Curve Description: Irrigated Alfalfa, statewide

Percent Production by Month

<u>Jan</u>	<u>Feb</u>	<u>Mar</u>	<u>Apr</u>	<u>May</u>	<u>Jun</u>	<u>Jul</u>	<u>Aug</u>	<u>Sep</u>	<u>Oct</u>	<u>Nov</u>	<u>Dec</u>
0	0	0	5	25	25	20	15	10	0	0	0

Soil Limitations

Soil blowing is a severe hazard during stand establishment or renovation of forage stands on the soils of this group. Bare areas where livestock concentrate are also susceptible. Production potential is low to moderate due to the low available water capacity and droughtiness of most of these soils. Species choices are also somewhat limited for pasture and hayland for the same reasons.

Management Interpretations

The impact on yields of the low available water capacity of most these soils can be reduced by selecting forage species that are highly tolerant to periods of drought and inadequate soil moisture and can grow on coarse soils. Incorporate wind erosion control practices during stand establishment. Properly locating facilitating practices such as fences, lanes, and water developments can help control livestock movement, reduce trailing perpendicular to steeper slopes, evenly distribute grazing pressure, and reduce bare areas.

FSG Documentation

Similar FSG's:

FSG ID

G102CY130NE

FSG Narrative

Very Droughty Loam soils have finer textures than sands.

Inventory Data References:

Agriculture Handbook 296-Land Resource Regions and Major Land Resource Areas

Natural Resources Conservation Service (NRCS) National Water and Climate Center data

USDA Plant Hardiness Zone maps

National Soil Survey Information System for soil surveys in Nebraska and South Dakota counties in MLRA 102C

NRCS Nebraska Field Office Technical Guide and South Dakota Technical Guide

NRCS National Range and Pasture Handbook

Various Agricultural Research Service, Cooperative Extension Service, and NRCS research trials for plant adaptation and production.

State Correlation:

This site has been correlated with the following states: Nebraska and South Dakota

Forage Suitability Group Approval:

Original Author: Tim Nordquist

Original Date: 6/8/2001

Approval by: Dave Schmidt

Approval Date: 1/13/05