

United States Department of Agriculture Natural Resources Conservation Service

Ecological Site Description

Site Type: Rangeland

Site Name: Loamy

Site ID: R102BY010SD


Major Land Resource Area (MLRA): 102B – Till Plains

Physiographic Features

This site occurs on nearly level to moderately sloping uplands.

Landform: till plain, moraine, outwash terrace **Aspect:** N/A

	<u>Minimum</u>	<u>Maximum</u>
Elevation (feet):	1,100	1,900
Slope (percent):	1	25
Water Table Depth (inches):	48	80
Flooding:		
Frequency:	None	Rare
Duration:	None	Brief
Ponding:		
Depth (inches):	None	None
Frequency:	None	None
Duration:	None	None
Runoff Class:	Medium	Very high



Climatic Features

MLRA 102B is considered to have a continental climate – cold winters and relatively hot summers, low to moderate humidity, light rainfall, and much sunshine. Extremes in temperature may also abound. The climate is the result of this MLRA's location near the geographic center of North America. There are few natural barriers on the Northern Great Plains and air masses move freely across the plains and account for rapid changes in temperature.

Annual precipitation typically ranges from 24 to 26 inches per year. The average annual temperature is about 46°F. January is the coldest month with average temperatures ranging from about 14°F (Wentworth 2 WNW, South Dakota (SD)), to about 18°F (Canton 4 WNW, SD). July is the warmest month with temperatures averaging from about 72°F (Wentworth 2 WNW, SD), to about 73°F (Canton 4 WNW, SD). The range of normal average monthly temperatures between the coldest and warmest months is about 57°F. This large annual range attests to the continental nature of this area's climate. Hourly winds are estimated to average about 11 miles per hour (mph) annually, ranging from about 13 mph during the spring to about 10 mph during the summer. Daytime winds are generally stronger than nighttime and occasional strong storms may bring brief periods of high winds with gusts to more than 50 mph.

Growth of cool-season plants begins in early to mid-March, slowing or ceasing in late June. Warm-season plants begin growth about mid-May and continue to early or mid-September. Greenup of cool-season plants may occur in September and October when adequate soil moisture is present.

	<u>Minimum</u>	<u>Maximum</u>
Frost-free period (days):	145	147
Freeze-free period (days):	162	165
Mean Annual Precipitation (inches):	24	26

Average Monthly Precipitation (inches) and Temperature (°F):

	Precip. Min.	Precip. Max	Temp. Min.	Temp. Max.
January	0.51	0.52	4.0	28.3
February	0.58	0.69	7.0	31.6
March	1.13	1.71	20.1	45.0
April	2.36	3.05	33.2	60.3
May	3.25	3.85	44.4	72.2
June	3.86	4.28	54.5	81.1
July	2.97	3.16	59.2	85.8
August	3.02	3.03	57.2	84.4
September	2.55	2.77	48.4	76.2
October	1.63	2.19	36.0	63.8
November	0.96	1.33	21.6	44.9
December	0.48	0.54	9.7	31.3

Climate Stations		Period	
Station ID	Location or Name	From	To
SD1392	Canton 4 WNW	1896	2009
SD1579	Centerville 6 SE	1897	2009
SD7667	Sioux Falls WSFO	1948	2009
SD9042	Wentworth 2 WNW	1893	2006

For local climate stations that may be more representative, refer to <http://www.wcc.nrcs.usda.gov>.

Riparian and Wetland Features

No riparian areas or wetland features are directly associated with this site.

Representative Soil Features

The common features of soils in this site are fine sandy loam to clay loam textured subsurface soils, with slopes ranging from 1 to 25 percent. Some soils are underlain with sand and/or gravel at about two to three feet in depth. The soils in this site are well-drained and typically formed in till, drift, and/or loess. The loam to silty clay loam surface layer is typically 6 to 20 inches thick. The soils have a slow to moderately slow infiltration rate. This site typically should show slight to no evidence of rills, wind scoured areas, or pedestalled plants. If present, water flow paths are broken, irregular in appearance, or discontinuous. The soil surface is stable and intact. Subsurface soil layers are nonrestrictive to water movement and root penetration. These soils are mainly susceptible to water erosion. The hazard of water erosion increases on slopes greater than about nine percent. Loss of 50 percent or more of the surface layer of the soils on this site can result in a shift in species composition and/or production. Access Web Soil Survey (<http://websoilsurvey.nrcs.usda.gov/app/HomePage.htm>) for specific local soils information.

Parent Material Kind: till, unspecified, and loess
Parent Material Origin:
Surface Texture: loam, silt loam, silty clay loam
Surface Texture Modifier: none
Subsurface Texture Group: loamy
Surface Fragments ≤3” (% Cover): 0-5
Surface Fragments >3” (%Cover): 0-10
Subsurface Fragments ≤3” (% Volume): 0-30
Subsurface Fragments >3” (% Volume): 0-4

	<u>Minimum</u>	<u>Maximum</u>
Drainage Class:	moderately well	well
Permeability Class:	slow	moderately slow
Depth (inches):	40	80
Electrical Conductivity (mmhos/cm)*:	0	4
Sodium Absorption Ratio*:	0	2
Soil Reaction (1:1 Water)*:	5.6	8.4
Soil Reaction (0.1M CaCl2)*:	NA	NA
Available Water Capacity (inches)*:	6	8
Calcium Carbonate Equivalent (percent)*:	0	30

*These attributes represent from 0-40 inches or to the first restrictive layer.

Plant Communities

Ecological Dynamics of the Site


This site developed under Northern Great Plains climatic conditions, light to severe grazing by bison and other large herbivores, sporadic natural or man-caused wildfire (often of light intensities), and other biotic and abiotic factors that typically influence soil/site development. Changes will occur in the plant communities due to short-term weather variations, impacts of native and/or exotic plant and animal species, and management actions. While the following plant community descriptions describe more typical transitions that will occur, severe disturbances, such as periods of well below average precipitation, can cause significant shifts in plant communities and/or species composition that may not be described within this document.

Continuous season-long grazing (during the typical growing season of May through October) and/or repeated seasonal grazing (e.g., every spring, every summer) without adequate recovery periods following grazing events causes departure from the 3.1 Bluestem/Needlegrass Plant Community Phase. Little bluestem, wheatgrass, sideoats grama, and blue grama will increase. Eventually, blue grama, quackgrass, and Kentucky bluegrass may develop into a sod. Indiangrass, big bluestem, porcupine grass, green needlegrass, sideoats grama, and little bluestem will decrease in frequency and production. Extended periods of nonuse and/or lack of fire will result in excessive litter and a plant community dominated by cool-season grasses such as Kentucky bluegrass, smooth brome grass, and green needlegrass.

Interpretations are primarily based on the 1.1 Bluestem/Needlegrass Plant Community Phase. It has been determined by study of rangeland relic areas, areas protected from excessive disturbance, and areas under long-term rotational grazing regimes. Trends in plant community dynamics ranging from heavily grazed to lightly grazed areas, seasonal use pastures, and historical accounts also have been used. Plant community phases, states, transitional pathways, and thresholds have been determined through similar studies and experience.

The following is a diagram that illustrates the common plant community phases that can occur on the site and the transition pathways between communities. These are the most common plant community phases based on current knowledge and experience and changes may be made as more data is collected. Narratives following the diagram contain more detail pertaining to the ecological processes.

Plant Communities and Transitional Pathways



Refer to narrative for details on pathways: **C** – Cropped, abandoned; **CSLG** – Continuous season-long grazing; **E** – Encroachment of introduced species; **F** – Fire; **GPF** – Grazing, precipitation, and/or fire returning to more normal disturbance regime levels and frequencies; **HCG** – Heavy continuous grazing; **HCSG** – Heavy continuous seasonal grazing; **HCSLG** – Heavy continuous season-long grazing; **LTPG** – Long-term prescribed grazing; **NU, NF** – Non-use, no fire; **PG** – Prescribed grazing; **PM** – Pest management (herbicide); **S** – Seeding.

Any Plant Community

Plant Community Composition and Group Annual Production

			1.1 Bluestem/Needlegrass		
COMMON/GROUP NAME	SCIENTIFIC NAME	SYMBOL	Group	lbs./acre	% Comp
GRASSES & GRASS-LIKES				2960 - 3330	80 - 90
TALL WARM-SEASON GRASSES			1	740 - 1665	20 - 45
big bluestem	Andropogon gerardii	ANGE	1	370 - 1295	10 - 35
Indiangrass	Sorghastrum nutans	SONU2	1	185 - 925	5 - 25
switchgrass	Panicum virgatum	PAVI2	1	74 - 370	2 - 10
tall dropseed	Sporobolus compositus var. compositus	SPCOC2	1	0 - 111	0 - 3
COOL-SEASON BUNCHGRASSES			2	370 - 740	10 - 20
green needlegrass	Nassella viridula	NAVI4	2	185 - 740	5 - 20
porcupine grass	Hesperostipa spartea	HESP11	2	185 - 740	5 - 20
Canada wildrye	Elymus canadensis	ELCA4	2	0 - 111	0 - 3
MID WARM-SEASON GRASSES			3	370 - 740	10 - 20
little bluestem	Schizachyrium scoparium	SCSC	3	185 - 555	5 - 15
sideoats grama	Bouteloua curtipendula	BOCU	3	74 - 370	2 - 10
prairie dropseed	Sporobolus heterolepis	SPHE	3	74 - 370	2 - 10
WHEATGRASS			4	185 - 370	5 - 10
slender wheatgrass	Elymus trachycaulus	ELTR7	4	74 - 370	2 - 10
western wheatgrass	Pascopyrum smithii	PASM	4	0 - 185	0 - 5
OTHER NATIVE GRASSES			5	111 - 259	3 - 7
blue grama	Bouteloua gracilis	BOGR2	5	37 - 185	1 - 5
prairie junegrass	Koeleria macrantha	KOMA	5	37 - 74	1 - 2
Scribner panicum	Dichanthelium oligosanthes var. scribnerianum	DIOLS	5	0 - 37	0 - 1
Wilcox panicum	Dichanthelium wilcoxianum	DIWI5	5	0 - 37	0 - 1
other grasses		2GRAM	5	37 - 185	1 - 5
GRASS-LIKES			6	37 - 185	1 - 5
sedge	Carex spp.	CAREX	6	37 - 185	1 - 5
other grass-likes		2GL	6	0 - 111	0 - 3
FORBS			8	185 - 555	5 - 15
American pasqueflower	Pulsatilla patens ssp. multifida	PUPAM	8	0 - 37	0 - 1
American vetch	Vicia americana	VIAM	8	37 - 74	1 - 2
aromatic aster	Symphyotrichum oblongifolium	SYOB	8	0 - 74	0 - 2
Canada goldenrod	Solidago canadensis	SOCA6	8	37 - 74	1 - 2
catclaw sensitive briar	Mimosa nuttallii	MINU6	8	0 - 37	0 - 1
compassplant	Silphium laciniatum	SILA3	8	0 - 74	0 - 2
cudweed sagewort	Artemisia ludoviciana	ARLU	8	37 - 74	1 - 2
false boneset	Brickellia eupatorioides	BREU	8	37 - 74	1 - 2
false gromwell	Onosmodium bejariense var. occidentale	ONBEO	8	37 - 74	1 - 2
gayfeather	Liatris spp.	LIATR	8	37 - 111	1 - 3
green sagewort	Artemisia campestris	ARCA12	8	37 - 74	1 - 2
heath aster	Symphyotrichum ericoides	SYER	8	37 - 74	1 - 2
Missouri goldenrod	Solidago missouriensis	SOMI2	8	0 - 37	0 - 1
prairie coneflower	Ratibida columnifera	RACO3	8	37 - 74	1 - 2
purple coneflower	Echinacea angustifolia	ECAN2	8	37 - 74	1 - 2
purple prairie clover	Dalea purpurea	DAPU5	8	37 - 74	1 - 2
rush skeletonweed	Lygodesmia juncea	LYJU	8	0 - 37	0 - 1
scurfpea	Psoraleidum spp.	PSORA2	8	37 - 111	1 - 3
stiff goldenrod	Oligoneuron rigidum	OLRI	8	37 - 74	1 - 2
stiff sunflower	Helianthus pauciflorus	HEPA19	8	37 - 74	1 - 2
wavyleaf thistle	Cirsium undulatum	CIUN	8	0 - 74	0 - 2
western ragweed	Ambrosia psilostachya	AMPS	8	37 - 74	1 - 2
western yarrow	Achillea millefolium var. occidentalis	ACMIO	8	37 - 74	1 - 2
woolly verbena	Verbena stricta	VEST	8	0 - 37	0 - 1
native forbs		2FN	8	37 - 185	1 - 5
SHRUBS			9	74 - 296	2 - 8
fringed sagewort	Artemisia frigida	ARFR4	9	0 - 37	0 - 1
leadplant	Amorpha canescens	AMCA6	9	37 - 148	1 - 4
rose	Rosa spp.	ROSA5	9	37 - 111	1 - 3
smooth sumac	Rhus glabra	RHGL	9	0 - 74	0 - 2
snowberry	Symphoricarpos spp.	SYMPH	9	37 - 111	1 - 3
other shrubs		2SHRUB	9	0 - 74	0 - 2

Annual Production lbs./acre	LOW	RV	HIGH
GRASSES & GRASS-LIKES	2465 -	3145	-3695
FORBS	165 -	370	-665
SHRUBS	70 -	185	-340
TOTAL	2700 -	3700	-4700

This list of plants and their relative proportions are based on near normal years. Fluctuations in species composition and relative production may change from year to year dependent upon precipitation or other climatic factors. RV = Representative value.

Plant Community Composition and Group Annual Production

COMMON/GROUP NAME	SYMBOL	1.1 Bluestem/Needlegrass			3.2 Needlegrass/Little Blue-stem/Kentucky Bluegrass			4.1 Smooth Bromegrass/Kentucky Bluegrass			4.2 Kentucky Bluegrass/Quackgrass		
		Grp	lbs./acre	% Comp	Grp	lbs./acre	% Comp	Grp	lbs./acre	% Comp	Grp	lbs./acre	% Comp
GRASSES & GRASS-LIKES			2960 - 3330	80 - 90		2550 - 2850	85 - 95		2720 - 3230	80 - 95		1920 - 2280	80 - 95
TALL WARM-SEASON GRASSES		1	740 - 1665	20 - 45	1	150 - 450	5 - 15	1	0 - 170	0 - 5	1	0 - 120	0 - 5
big bluestem	ANGE	1	370 - 1295	10 - 35	1	60 - 450	2 - 15	1	0 - 102	0 - 3	1	0 - 72	0 - 3
Indiangrass	SONU2	1	185 - 925	5 - 25	1	0 - 150	0 - 5						
switchgrass	PAV12	1	74 - 370	2 - 10	1	0 - 150	0 - 5				1	0 - 72	0 - 3
tall dropseed	SPCOC2	1	0 - 111	0 - 3	1	0 - 150	0 - 5	1	0 - 170	0 - 5	1	0 - 120	0 - 5
COOL-SEASON BUNCHGRASSES		2	370 - 740	10 - 20	2	300 - 900	10 - 30	2	0 - 340	0 - 10	2	0 - 120	0 - 5
green needlegrass	NAVI4	2	185 - 740	5 - 20	2	150 - 750	5 - 25	2	0 - 340	0 - 10	2	0 - 120	0 - 5
porcupine grass	HESP11	2	185 - 740	5 - 20	2	60 - 300	2 - 10	2	0 - 68	0 - 2			
Canada wildrye	ELCA4	2	0 - 111	0 - 3	2	0 - 30	0 - 1	2	0 - 68	0 - 2			
MID WARM-SEASON GRASSES		3	370 - 740	10 - 20	3	450 - 750	15 - 25	3	0 - 170	0 - 5	3	0 - 120	0 - 5
little bluestem	SCSC	3	185 - 555	5 - 15	3	300 - 750	10 - 25	3	0 - 170	0 - 5	3	0 - 72	0 - 3
sideoats grama	BOCU	3	74 - 370	2 - 10	3	60 - 450	2 - 15	3	0 - 170	0 - 5	3	0 - 72	0 - 3
prairie dropseed	SPHE	3	74 - 370	2 - 10	3	0 - 150	0 - 5						
WHEATGRASS		4	185 - 370	5 - 10	4	150 - 450	5 - 15	4	0 - 170	0 - 5	4	0 - 120	0 - 5
slender wheatgrass	ELTR7	4	74 - 370	2 - 10	4	30 - 240	1 - 8	4	0 - 170	0 - 5	4	0 - 24	0 - 1
western wheatgrass	PASM	4	0 - 185	0 - 5	4	60 - 300	2 - 10	4	0 - 170	0 - 5	4	0 - 120	0 - 5
OTHER NATIVE GRASSES		5	111 - 259	3 - 7	5	150 - 300	5 - 10	5	0 - 170	0 - 5	5	24 - 240	1 - 10
blue grama	BOGR2	5	37 - 185	1 - 5	5	60 - 240	2 - 8	5	0 - 136	0 - 4	5	24 - 240	1 - 10
prairie junegrass	KOMA	5	37 - 74	1 - 2	5	30 - 60	1 - 2	5	0 - 34	0 - 1	5	0 - 24	0 - 1
Scribner panicum	D1OLS	5	0 - 37	0 - 1	5	0 - 30	0 - 1	5	0 - 34	0 - 1	5	0 - 24	0 - 1
Wilcox panicum	DIWI5	5	0 - 37	0 - 1	5	0 - 30	0 - 1	5	0 - 34	0 - 1	5	0 - 24	0 - 1
other grasses	ZGRAM	5	37 - 185	1 - 5	5	30 - 150	1 - 5	5	0 - 170	0 - 5	5	0 - 72	0 - 3
GRASS-LIKES		6	37 - 185	1 - 5	6	30 - 150	1 - 5	6	0 - 170	0 - 5	6	24 - 120	1 - 5
sedge	CAREX	6	37 - 185	1 - 5	6	30 - 150	1 - 5	6	0 - 170	0 - 5	6	24 - 120	1 - 5
other grass-likes	ZGL	6	0 - 111	0 - 3	6	0 - 90	0 - 3	6	0 - 68	0 - 2	6	0 - 24	0 - 1
NON-NATIVE GRASSES		7			7	150 - 450	5 - 15	7	1020 - 2550	30 - 75	7	480 - 1800	20 - 75
Kentucky bluegrass	POPR				7	150 - 450	5 - 15	7	340 - 1700	10 - 50	7	240 - 1440	10 - 60
quackgrass	ELRE4				7	0 - 150	0 - 5	7	0 - 340	0 - 10	7	120 - 960	5 - 40
smooth bromegrass	BRIN2				7	0 - 240	0 - 8	7	680 - 2380	20 - 70	7	120 - 720	5 - 30
FORBS		8	185 - 555	5 - 15	8	150 - 300	5 - 10	8	170 - 510	5 - 15	8	120 - 360	5 - 15
American pasqueflower	PUPAM	8	0 - 37	0 - 1									
American vetch	VIAM	8	37 - 74	1 - 2	8	0 - 30	0 - 1						
aromatic aster	SYOB	8	0 - 74	0 - 2	8	0 - 30	0 - 1						
Canada goldenrod	SOCA6	8	37 - 74	1 - 2	8	30 - 90	1 - 3	8	34 - 238	1 - 7	8	24 - 144	1 - 6
catclaw sensitive briar	MINU6	8	0 - 37	0 - 1									
compassplant	SILA3	8	0 - 74	0 - 2									
cudweed sagewort	ARLU	8	37 - 74	1 - 2	8	30 - 90	1 - 3	8	34 - 238	1 - 7	8	0 - 120	0 - 5
false boneset	BREU	8	37 - 74	1 - 2	8	0 - 30	0 - 1						
false gromwell	ONBEO	8	37 - 74	1 - 2	8	0 - 30	0 - 1						
gayfeather	LIATR	8	37 - 111	1 - 3	8	0 - 30	0 - 1						
green sagewort	ARCA12	8	37 - 74	1 - 2	8	30 - 60	1 - 2	8	34 - 136	1 - 4	8	24 - 96	1 - 4
heath aster	SYER	8	37 - 74	1 - 2	8	30 - 90	1 - 3	8	34 - 170	1 - 5	8	0 - 96	0 - 4
Missouri goldenrod	SOM12	8	0 - 37	0 - 1	8	0 - 30	0 - 1						
prairie coneflower	RAC03	8	37 - 74	1 - 2	8	0 - 30	0 - 1						
purple coneflower	ECAN2	8	37 - 74	1 - 2	8	0 - 30	0 - 1						
purple prairie clover	DAPU5	8	37 - 74	1 - 2	8	0 - 30	0 - 1						
rush skeletonweed	LYJU	8	0 - 37	0 - 1	8	0 - 30	0 - 1	8	0 - 34	0 - 1			
scurfpea	PSORA2	8	37 - 111	1 - 3	8	30 - 90	1 - 3	8	34 - 136	1 - 4	8	0 - 72	0 - 3
stiff goldenrod	OLRI	8	37 - 74	1 - 2	8	30 - 90	1 - 3	8	34 - 136	1 - 4	8	24 - 120	1 - 5
stiff sunflower	HEPA19	8	37 - 74	1 - 2	8	0 - 30	0 - 1						
wavyleaf thistle	CIUN	8	0 - 74	0 - 2	8	0 - 30	0 - 1						
western ragweed	AMPS	8	37 - 74	1 - 2	8	30 - 90	1 - 3	8	34 - 170	1 - 5	8	24 - 120	1 - 5
western yarrow	ACMIO	8	37 - 74	1 - 2	8	30 - 90	1 - 3	8	34 - 136	1 - 4	8	24 - 120	1 - 5
woolly verbena	VEST	8	0 - 37	0 - 1	8	0 - 60	0 - 2	8	0 - 102	0 - 3	8	0 - 72	0 - 3
native forbs	ZFN	8	37 - 185	1 - 5	8	0 - 60	0 - 2	8	0 - 34	0 - 1	8	0 - 24	0 - 1
introduced forbs	ZFI				8	30 - 90	1 - 3	8	34 - 340	1 - 10	8	24 - 240	1 - 10
SHRUBS		9	74 - 296	2 - 8	9	30 - 150	1 - 5	9	34 - 170	1 - 5	9	0 - 120	0 - 5
fringed sagewort	ARFR4	9	0 - 37	0 - 1	9	0 - 60	0 - 2	9	0 - 34	0 - 1	9	0 - 72	0 - 3
leadplant	AMCA6	9	37 - 148	1 - 4	9	0 - 30	0 - 1						
rose	ROSA5	9	37 - 111	1 - 3	9	30 - 60	1 - 2	9	0 - 34	0 - 1			
smooth sumac	RHGL	9	0 - 74	0 - 2	9	0 - 120	0 - 4	9	0 - 170	0 - 5	9	0 - 120	0 - 5
snowberry	SYMPH	9	37 - 111	1 - 3	9	0 - 60	0 - 2	9	34 - 102	1 - 3	9	0 - 48	0 - 2
other shrubs	ZSHRUB	9	0 - 74	0 - 2	9	0 - 30	0 - 1	9	0 - 68	0 - 2	9	0 - 24	0 - 1
Annual Production lbs./acre			LOW RV HIGH		LOW RV HIGH		LOW RV HIGH		LOW RV HIGH		LOW RV HIGH		LOW RV HIGH
GRASSES & GRASS-LIKES			2465 - 3145 - 3695		2040 - 2685 - 3290		2420 - 2958 - 3500		1595 - 2100 - 2450				
FORBS			165 - 370 - 665		135 - 225 - 345		150 - 340 - 610		105 - 240 - 420				
SHRUBS			70 - 185 - 340		25 - 90 - 165		30 - 102 - 190		0 - 60 - 130				
TOTAL			2700 - 3700 - 4700		2200 - 3000 - 3800		2600 - 3400 - 4300		1700 - 2400 - 3000				

This list of plants and their relative proportions are based on near normal years. Fluctuations in species composition and relative production may change from year to year dependent upon precipitation or other climatic factors. RV = Representative value. Refer to PLANTS database for scientific names and codes: <http://plants.usda.gov>

Plant Community and Vegetation State Narratives

Reference State (State 1)

This state represents the natural range of variability that dominated the dynamics of this ecological site (ES). This state was dominated by warm-season grasses, with cool-season grasses being subdominant. In pre-European times, the primary disturbance mechanisms for this site in the reference condition included periods of below and/or above average precipitation, periodic fire, and herbivory by insects and large ungulates. Timing of fires and herbivory coupled with weather events dictated the dynamics that occurred within the natural range of variability. In some locations, this site likely received relatively heavy grazing pressure. Tall warm-season grasses would have declined and cool-season bunchgrasses and short to mid-statured warm-season grasses would have increased. Today, a similar state (State 3) can be found on areas that are properly managed with grazing and/or prescribed burning and sometimes on areas receiving occasional short periods of rest.

1.1 Bluestem/Needlegrass Plant Community Phase

Interpretations are based primarily on the Bluestem/Needlegrass Plant Community Phase (this is also considered to be climax). The potential vegetation was about 80 percent grasses or grass-like plants, 15 percent forbs, and 5 percent shrubs. The community was dominated by warm-season grasses, with cool-season grasses being subdominant. The major grasses included big bluestem, Indiangrass, porcupine grass, green needlegrass, and little bluestem. Other grass or grass-like species included switchgrass, sideoats grama, slender wheatgrass, prairie dropseed, and sedges. This plant community was resilient and well adapted to the Northern Great Plains climatic conditions. The diversity in plant species allowed for high drought tolerance. This was a sustainable plant community in regards to site/soil stability, watershed function, and biologic integrity.

The following growth curve is an estimate of the monthly percentages of total annual growth of the dominant species expected during an average year:

Growth curve number: SD0214

Growth curve name: Till Plains, warm-season dominant, cool-season subdominant.

Growth curve description: Warm-season dominant, cool-season subdominant.

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
0	0	3	7	17	25	25	15	7	1	0	0

Transitions or pathways leading to other plant communities were as follows:

- 1.1a – Heavy continuous grazing which included herbivory at moderate to heavy levels at the same time of year each year without adequate recovery periods, or a combination of disturbances such as extended periods of below average precipitation coupled with periodic or chronic heavy grazing would have shifted this community to the *1.2 Needlegrass/Little Bluestem/Big Bluestem/Grama Plant Community Phase*.
- 1.1b – Fire occurring at relatively frequent intervals and occasional grazing events immediately following early season fire caused a reduction in cool-season grasses and an increase in warm-season grasses. The warm-season grasses were more tolerant of shorter return intervals of fire and would increase in vigor and production leading to a temporary shift to the *1.3 Big Bluestem/Indiangrass/Sideoats Grama Plant Community Phase*.

1.2 Needlegrass/Little Bluestem/Big Bluestem/Grama Plant Community Phase

This plant community evolved under heavy continuous grazing or from over utilization during extended drought periods. The potential plant community was made up of approximately 85 percent grasses and grass-like species, 10 percent forbs, and 5 percent shrubs.

Dominant grasses included green needlegrass, little bluestem, big bluestem, and sideoats grama. Grasses of secondary importance included blue grama, western wheatgrass, porcupine grass, Indiangrass, switchgrass, tall dropseed, prairie dropseed, slender wheatgrass, and sedge. Forbs commonly found in this plant community included Canada goldenrod, cudweed sagewort, heath aster, scurfpea, stiff goldenrod, western ragweed, and western yarrow. This plant community had similar plant composition to the 3.2 Needlegrass/Little Bluestem/Kentucky Bluegrass Plant Community Phase (refer to the plant composition tables). The main difference is that this plant community phase did not have the presence of nonnative invasive species such as Kentucky bluegrass and smooth brome grass. When compared to the Bluestem/Needlegrass Plant Community Phase (1.1), green needlegrass and little bluestem increased. Production of tall warm-season grasses was reduced. This plant community was moderately resistant to change. The herbaceous species present were well adapted to grazing; however, species composition could be altered through long-term overgrazing. If the herbaceous component was intact, it tended to be resilient if the disturbance was not long-term. Most of the components of the ecological processes would have been functioning at optimum levels. However, the vigor and reproductive capability of the tall warm-season grasses would have been reduced due to grazing pressure or a combination of stressors. A reduction of this dominant functional group allowed for an increase in shorter-statured (and shallower rooted) species.

The following growth curve is an estimate of the monthly percentages of total annual growth of the dominant species expected during an average year:

Growth curve number: SD0213

Growth curve name: Till Plains, cool-season/warm-season codominant.

Growth curve description: Cool-season, warm-season codominant.

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
0	0	3	10	20	28	21	10	5	3	0	0

Transitions or pathways leading to other plant communities are as follows:

- T1a – Heavy continuous grazing (stocking levels well above carrying capacity for extended portions of the growing season and often at the same time of year each year) would have converted this plant community to the *2.1 Little Bluestem/Wheatgrass/Grama Plant Community Phase* and the *Degraded State*.
- 1.2a – Grazing, precipitation, and/or fire returned to normal disturbance regime levels and frequencies or periodic light to moderate grazing possibly including periodic rest would have converted this plant community to the *1.1 Bluestem/Needlegrass Plant Community Phase*.

2.1 Little Bluestem/Wheatgrass/Grama Plant Community

This plant community evolved under heavy continuous season grazing or from over utilization during extended drought periods. The potential plant community was made up of approximately 85 percent grasses and grass-like species, 10 percent forbs, and 5 percent shrubs. Dominant grasses included little bluestem, western wheatgrass, sideoats grama, and blue grama. Grasses of secondary importance included big bluestem, switchgrass, green needlegrass, slender wheatgrass, tall dropseed, and sedge. Forbs commonly found in this plant community included cudweed sagewort, green sagewort, and western yarrow. When compared to the Bluestem/Needlegrass Plant Community Phase (1.1), tall warm-season grasses were reduced, and the more grazing tolerant species such as blue grama, little bluestem, and sideoats grama were dominant on this plant community. With the exception of western wheatgrass, cool-season grasses decreased significantly. This vegetation state was very resistant to change especially if the disturbance continued and the short-statured species such as blue grama increased. The herbaceous species present were well adapted to grazing. This plant community was less productive than other phases.

The following growth curve is an estimate of the monthly percentages of total annual growth of the dominant species expected during an average year:

Growth curve number: SD0214

Growth curve name: Till Plains, warm-season dominant, cool-season subdominant.

Growth curve description: Warm-season dominant, cool-season subdominant.

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
0	0	3	7	17	25	25	15	7	1	0	0

1.3 Big Bluestem/Indiangrass/Sideoats Grama Plant Community Phase

This plant community was a result of fire occurring at relatively frequent intervals. This phase could have also resulted from a combination of grazing events immediately following early season fire (i.e., large ungulates attracted to highly nutritious vegetative growth following a fire). These events would have caused a reduction in cool-season grasses and an increase in warm-season grasses. The warm-season grasses were more tolerant of shorter return intervals of fire and would have increased in vigor and production leading to a temporary shift to this phase. Needlegrasses would have decreased most significantly amongst the cool-season grasses. The potential vegetation was about 80 percent grasses or grass-like plants, 15 percent forbs, and 5 percent shrubs. The community was dominated by warm-season grasses. The major grasses included big bluestem, Indiangrass, switchgrass, sideoats grama, little bluestem, and prairie dropseed. Other grass or grass-like species included green needlegrass, porcupine grass, western wheatgrass, blue grama, slender wheatgrass, tall dropseed, and sedge. This plant community was not resistant to change and would have readily shifted back to the 1.1 Bluestem/Needlegrass Plant Community Phase with a return of more normal fire return intervals.

The following growth curve is an estimate of the monthly percentages of total annual growth of the dominant species expected during an average year:

Growth curve number: SD0215

Growth curve name: Till Plains, warm-season dominant.

Growth curve description: Warm-season dominant.

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
0	0	2	5	15	25	30	15	7	1	0	0

Transitions or pathways leading to other plant communities were as follows:

- 1.3a – Grazing, precipitation, and/or fire returned to normal disturbance regime levels and frequencies or periodic light to moderate grazing possibly including periodic rest will convert this plant community to the *1.1 Bluestem/Needlegrass Plant Community Phase*.

Transition from Reference State (State 1) to the Native/Invaded Grass State (State 3)

- T1b – Encroachment of non-native grasses such as Kentucky bluegrass and smooth brome grass, and disruption of natural disturbance regimes (typically as a result of fire suppression following settlement) led this state over a threshold to the *Native/Invaded Grass State (State 3)*.

Native/Invaded Grass State (State 3)

This state represents the more common range of variability that exists with higher levels of grazing management but in the absence of periodic fire due to fire suppression. This state is dominated by cool- and warm-season grasses. It can be found on areas that are properly managed with grazing

and/or prescribed burning, and sometimes on areas receiving occasional short periods of rest. Taller warm-season species can decline and a corresponding increase in short statured grass will occur.

3.1 Bluestem/Needlegrass Plant Community Phase

This plant community phase is similar to the 1.1 Bluestem/Needlegrass Plant Community Phase, but it also contains minor amounts of nonnative invasive grass species such as Kentucky bluegrass and smooth brome grass (up to about 10 percent by air-dry weight). The potential vegetation is about 80 percent grasses or grass-like plants, 15 percent forbs, and 5 percent shrubs. This community is dominated by warm-season grasses, with cool-season grasses being subdominant. The major grasses include big bluestem, Indiangrass, porcupine grass, green needlegrass, and little bluestem. Other grass or grass-like species include switchgrass, sideoats grama, slender wheatgrass, prairie dropseed, and sedges. This plant community is resilient and well adapted to the Northern Great Plains climatic conditions. The diversity in plant species allows for high drought tolerance. This is a sustainable plant community in regards to site/soil stability, watershed function, and biologic integrity.

The following growth curve is an estimate of the monthly percentages of total annual growth of the dominant species expected during an average year:

Growth curve number: SD0214

Growth curve name: Till Plains, warm-season dominant, cool-season subdominant.

Growth curve description: Warm-season dominant, cool-season subdominant.

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
0	0	3	7	17	25	25	15	7	1	0	0

Transitions or pathways leading to other plant communities are as follows:

- 3.1a – Heavy continuous grazing (stocking levels well above carrying capacity for extended portions of the growing season, and often at the same time of year each year), or continuous season-long grazing, or a combination of disturbances such as extended periods of below average precipitation coupled with periodic heavy grazing will shift this community to the 3.2 Needlegrass/Little Bluestem/Kentucky Bluegrass Plant Community Phase.

3.2 Needlegrass/Little Bluestem/Kentucky Bluegrass Plant Community Phase

This plant community is a result of heavy continuous grazing or from over utilization during extended drought periods. The potential plant community is made up of approximately 85 percent grasses and grass-like species, 10 percent forbs, and 5 percent shrubs. Dominant grasses include green needlegrass, little bluestem, big bluestem, sideoats grama, and Kentucky bluegrass. Grasses of secondary importance include blue grama, western wheatgrass, porcupine grass, Indiangrass, switchgrass, tall dropseed, prairie dropseed, slender wheatgrass, and sedge. Forbs commonly found in this plant community include Canada goldenrod, cudweed sagewort, heath aster, scurfpea, stiff goldenrod, western ragweed, and western yarrow.

When compared to the Bluestem/Needlegrass Plant Community Phase (1.1), green needlegrass and little bluestem have increased. Production of tall warm-season grasses is reduced. This plant community is moderately resistant to change. The herbaceous species present are well adapted to grazing; however, species composition can be altered through long-term overgrazing. If the herbaceous component is intact, it tends to be resilient if the disturbance is not long-term. Most of the components of the ecological processes are functioning at optimum levels. However, the vigor and reproductive capability of the tall warm-season grasses are reduced due to grazing pressure or a combination of stressors. A reduction of this dominant functional group allows for an increase in shorter-statured (and shallower rooted) species. The introduction of nonnative invasive species such as Kentucky bluegrass and smooth brome grass results in alterations to the soil profile. Organic

matter levels tend to decrease and begin to be concentrated more in the surface layers and the structure will begin to be modified. These changes favor the shallow-rooted species and hasten their eventual dominance if steps are not taken to reduce these species.

The following growth curve is an estimate of the monthly percentages of total annual growth of the dominant species expected during an average year:

Growth curve number: SD0213

Growth curve name: Till Plains, cool-season/warm-season codominant.

Growth curve description: Cool-season, warm-season codominant.

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
0	0	3	10	20	28	21	10	5	3	0	0

Transitions or pathways leading to other plant communities are as follows:

- T3b – Heavy continuous seasonal grazing (stocking levels well above carrying capacity for extended portions of the growing season, and at the same time of year each year) will cause a shift across a threshold leading to the *4.2 Kentucky Bluegrass/Quackgrass Plant Community Phase* within the *Invaded State (State 4)*.
- 3.2a – Prescribed grazing (alternating season of use and providing adequate recovery periods) or periodic light to moderate grazing possibly including periodic rest will convert this plant community to the *3.1 Bluestem/Needlegrass Plant Community Phase*.

Transition from Native/Invaded Grass State (State 3) to the Invaded State (State 4)

- T3a – Non-use and no fire for extended periods of time (typically for 10 or more years) will likely lead this state over a threshold resulting in the *4.1 Smooth Bromegrass/Kentucky Bluegrass Plant Community Phase* within the *4.0 Invaded State*.

Invaded State (State 4)

This state is a result of encroachment mainly by invasive introduced cool-season grasses. The ecological processes are not functioning, especially the biotic processes and the hydrologic functions. The introduced cool-season grasses cause reduced infiltration and increased runoff. Preliminary studies would tend to indicate this threshold may exist when Kentucky bluegrass exceeds 30 percent of the plant community and native grasses represent less than 40 percent of the plant community composition. The opportunity for high intensity spring burns is severely reduced by early green up and increased moisture and humidity at the soil surface and grazing pressure cannot cause a reduction in sodgrass dominance. Production is limited to the sod forming species. Infiltration continues to decrease and runoff increases and energy capture into the system is restricted to early season low producing species. Nutrient cycling is limited by root depth of the dominant species.

4.1 Smooth Bromegrass/Kentucky Bluegrass Plant Community Phase

This plant community phase is a result of extended periods of nonuse and no fire or occasionally light levels of grazing over several years. It is characterized by dominance of smooth bromegrass and to a lesser extent Kentucky bluegrass. The dominance is at times so complete that other species are difficult to find on the site. A thick duff layer also accumulates at or above the soil surface.

Nutrient cycling is greatly reduced and native plants have great difficulty becoming established. When dominated by smooth bromegrass, infiltration is moderately reduced and runoff is moderate. Production can be equal to or higher than the interpretive plant community. However, when dominated by Kentucky bluegrass, infiltration is greatly reduced and runoff is high. Production in this

case will likely be significantly less. In either case, the period that palatability is high is relatively short, as these cool-season species mature rapidly. Energy capture is also reduced.

The dominance of these introduced species has been shown to alter the biotic component of the soil, as well as, organic matter levels and eventually the soil structure. These alterations perpetuate the dominance of Kentucky bluegrass and smooth bromegrass and tend to make establishment of native species extremely difficult.

The following growth curve is an estimate of the monthly percentages of total annual growth of the dominant species expected during an average year:

Growth curve number: SD0211

Growth curve name: Till Plains, cool-season dominant.

Growth curve description: Cool-season dominant.

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
0	0	4	12	25	36	10	5	4	4	0	0

Transitions or pathways leading to other plant communities are as follows:

- 4.1a – Heavy continuous seasonal grazing (stocking levels well above carrying capacity for extended portions of the growing season, and at the same time of year each year) or heavy continuous season-long grazing will convert this plant community to the *4.2 Kentucky Bluegrass/Quackgrass Plant Community Phase*.

4.2 Kentucky Bluegrass/Quackgrass Plant Community Phase

This plant community phase is a result of heavy, continuous seasonal grazing or heavy, continuous season-long grazing. It is characterized by a dominance of Kentucky bluegrass and quackgrass. The dominance is at times so complete that other species are difficult to find on the site. A relatively thick duff layer can sometimes accumulate at or above the soil surface. Nutrient cycling is greatly reduced, and native plants have great difficulty becoming established. Infiltration is greatly reduced and runoff is high. Production will be significantly reduced when compared to the interpretive plant community. The period that palatability is high is relatively short as Kentucky bluegrass matures rapidly. Energy capture is also reduced. Biological activity in the soil is likely reduced significantly in this phase.

The following growth curve is an estimate of the monthly percentages of total annual growth of the dominant species expected during an average year:

Growth curve number: SD0211

Growth curve name: Till Plains, cool-season dominant.

Growth curve description: Cool-season dominant.

JAN	FEB	MAR	APR	MAY	JUN	JUL	AUG	SEP	OCT	NOV	DEC
0	0	4	12	25	36	10	5	4	4	0	0

Transitions or pathways leading to other plant communities are as follows:

- 4.2a – Prescribed grazing (alternating season of use and providing adequate recovery periods) or periodic light to moderate grazing possibly including periodic rest may convert this plant community to the *4.1 Smooth Bromegrass/Kentucky Bluegrass Plant Community Phase*.

Transition Pathway from Any Plant Community to the 4.3 Annual/Pioneer, Non-Native Perennial Plant Community Phase within the Invaded State (State 4)

- T5 – Encroachment of non-native invasive/noxious species, abandonment of cropping, or seeding of introduced and/or native improved varieties of forage species may lead this plant community phase over a threshold to the *Invaded State (State 4)* and more specifically to the *4.3 Annual/Pioneer, Non-native Perennial Plant Community Phase*. In the case of a seeding, refer to the corresponding Forage Suitability Group (FSG) description for adapted species and expected production (production estimates in the FSG description may be unrealistically high due to the degraded condition of the site at this phase).

4.3 Annual/Pioneer, Non-Native Perennial Plant Community Phase

This plant community developed under continuous heavy grazing or other excessive disturbances. The potential plant community is made up of approximately 40 to 80 percent grasses and grass-like species, 20 to 60 percent forbs, and 0 to 5 percent shrubs. The species present in this phase are highly variable but often include non-native invasive and/or early seral species. Plant diversity is low (plant richness may be high, but areas are often dominated by a few species). The ecological processes are difficult to restore because of the loss of plant diversity and overall soil disturbance. Soil erosion is potentially very high because of the bare ground and shallow rooted herbaceous plant community. Water runoff will increase and infiltration will decrease due to animal related soil compaction and loss of root mass due to low plant diversity and vigor. This plant community will require significant economic inputs and time to move towards another plant community. This movement is highly variable in its succession. This is due to the loss of diversity (including the loss of the seed bank), within the existing plant community, and the plant communities on adjacent sites. This community can be renovated to improve the production capability; however, if management changes are not made the vegetation could revert back to early seral species.

Transitions or pathways leading to other states are as follows:

- T4 – Pest management (herbicides) and often heavy continuous seasonal grazing will likely result in an eventual dominance by Kentucky bluegrass which will lead to the *4.2 Kentucky Bluegrass/Quackgrass Plant Community Phase*.

Restoration Pathway from the Invaded State (State 4) to the Native/Invaded Grass State (State 3)

- R4 - Long-term prescribed grazing (moderate stocking levels coupled with adequate recovery periods, or other grazing systems such as high-density, low-frequency intended to treat specific species dominance, or periodic light to moderate stocking levels possibly including periodic rest) may lead this plant community phase over a threshold to the *Native/Invaded Grass State (State 3)*. Pest management (i.e., herbicide) may also be needed to suppress cool-season invasive grasses. This will likely take a long period of time, possibly up to 10 years or more, and recovery may not be attainable. Success depends on whether native reproductive propagules remain intact on the site.

Ecological Site Interpretations

Animal Community – Wildlife Interpretations

-- Under Development --

Bluestem/Needlegrass Plant Community Phases (1.1 & 3.1):

Needlegrass/Little Bluestem/Big Bluestem/Grama Plant Community Phases (1.2):

Big Bluestem/Indiangrass/Sideoats Grama Plant Community Phase (1.3):

Little Bluestem/Wheatgrass/Grama Plant Community Phase (2.1):

Needlegrass/Little Bluestem/Kentucky Bluegrass Plant Community Phase (3.2):

Smooth Bromegrass/Kentucky Bluegrass Plant Community Phase (4.1):

Kentucky Bluegrass/Quackgrass Plant Community Phase (4.2):

Annual/Pioneer, Non-native Perennial Plant Community Phase (4.3):

Animal Preferences (Quarterly – 1,2,3,4[†])

Common Name	Cattle	Sheep	Horses	Deer	Antelope	Bison	Elk
Grasses and Grass-likes							
big bluestem	U D P D	U D U U	U D P D	U D U U	U D U U	U D P D	U D P D
blue grama	U D P U	D P P D	U D P U	D P P D	D P P D	U D P U	U D P U
Canada wildrye	U D U U	N U N N	U D U U	N U N N	N U N N	U D U U	U D U U
green needlegrass	U P U D	N P N P	U P U D	N P N P	N P N P	U P U D	U P U D
Indiangrass	U D P D	U D U U	U D P D	U D U U	U D U U	U D P D	U D P D
little bluestem	U D D U	N D N N	U D D U	N D N N	N D N N	U D D U	U D D U
porcupine grass	U P U D	N D N U	U P U D	N D N U	N D N U	U P U D	U P U D
prairie dropseed	N U P U	N U D U	N U P U	N U D U	N U D U	N U P U	N U P U
prairie junegrass	U D U D	N D N U	U D U D	N D N U	N D N U	U D U D	U D U D
Scribner panicum	U U D U	N U N N	U U D U	N U N N	N U N N	U U D U	U U D U
sedge	U D U D	U P N D	U D U D	U D U D	U D U D	U D U D	U D U D
sideoats grama	U D P U	U P D U	U D P U	U P D U	U P D U	U D P U	U D P U
slender wheatgrass	U P U U	N D U N	U P U U	N D U N	N D U N	U P U U	U P U U
switchgrass	U D D U	U D U U	U D D U	N N N N	N N N N	U D D U	U D D U
tall dropseed	N U N N	N U N N	N U N N	N U N N	N U N N	N U N N	N U N N
western wheatgrass	U P D U	N D N N	U P D U	N D N N	N D N N	U P D U	U P D U
Wilcox panicum	U U U U	N U N N	U U U U	N U N N	N U N N	U U U U	U U U U
Forbs							
American pasqueflower	N N N N	N U N N	N N N N	N U N N	N U N N	N N N N	N N N N
American vetch	U D P U	U P P U	U D P U	U P P U	U P P U	U D P U	U P P U
aromatic aster	U U U U	N N N N	U U U U	N N N N	N N N N	U U U U	N N N N
Canada goldenrod	N N U N	N U U N	N N U N	N U U N	N U U N	N N U N	N N U N
catclaw sensitive briar	U D P U	U P P U	U D P U	U P P U	U P P U	U D P U	U P P U
compassplant	U U D U	U P P U	U U D U	U P P U	U P P U	U U D U	U P P U
cudweed sagewort	U U U U	U U D U	U U U U	U U D U	U U D U	U U U U	U U D U
false boneset	U U D U	N D U N	U U D U	N D U N	N D U N	U U D U	N D U N
false gromwell	U U U U	N N N N	U U U U	N N N N	N N N N	U U U U	N N N N
gayfeather	U U D U	U P P U	U U D U	U P P U	U P P U	U U D U	U P P U
green sagewort	U U U U	U U U U	U U U U	U U U U	U U U U	U U U U	U U U U
heath aster	U U D U	U U P U	U U D U	U U P U	U U P U	U U D U	U U P U
Missouri goldenrod	U U D U	N U U N	U U D U	N U U N	N U U N	U U D U	N U U N
prairie coneflower	U U D U	U P P U	U U D U	U P P U	U P P U	U U D U	U P P U
purple coneflower	U U D U	U P P U	U U D U	U P P U	U P P U	U U D U	U P P U
purple prairie clover	U D P U	U P P U	U D P U	U P P U	U P P U	U D P U	U P P U
rush skeletonweed	U U U U	N N N N	U U U U	N N N N	N N N N	U U U U	N N N N
scurfpea	U U U U	N U U N	U U U U	N U U N	N U U N	U U U U	N U U N
stiff goldenrod	U U U U	N N U N	U U U U	N N U N	N N U N	U U U U	N N U N
stiff sunflower	U D P U	U D P U	U D P U	U D P U	U D P U	U D P U	U D P U
wavyleaf thistle	U U U U	N N N N	U U U U	N N N N	N N N N	U U U U	N N N N
western ragweed	U U U U	N N N N	U U U U	N N N N	N N N N	U U U U	N N N N
western yarrow	U U U U	N U U N	U U U U	N U U N	N U U N	U U U U	N U U N
woolly verbena	U U U U	N N N N	U U U U	N N N N	N N N N	U U U U	N N N N
Shrubs							
fringed sagewort	U U U U	U U U U	U U U U	U D D U	U P P D	U U U U	U U U D
leadplant	U P D U	U P D U	U P D U	U P D U	U P D U	U P D U	U P D U
rose	U D D U	U D D U	U D D U	U D D U	U D D U	U D D U	U D D U
smooth sumac	D U U D	D U U D	D U U D	D U U D	D U U D	D U U D	D U U D
snowberry	U U U U	U U U U	U U U U	D U D D	U U U U	U U U U	D U U U

N = not used; **U** = undesirable; **D** = desirable; **P** = preferred; **T** = toxic

[†] Quarters: 1 – Jan., Feb., Mar.; 2 – Apr., May, Jun.; 3 – Jul., Aug., Sep.; 4 – Oct., Nov., Dec.

Animal Community – Grazing Interpretations

The following table lists annual, suggested initial stocking rates with average growing conditions. These are conservative estimates that should be used only as guidelines in the initial stages of conservation planning. Often, the current plant composition does not entirely match any particular plant community (as described in this ES description). Because of this, a resource inventory is necessary to document plant composition and production. More accurate carrying capacity estimates should eventually be calculated using the following stocking rate information along with animal preference data and actual stocking records, particularly when grazers other than cattle are involved. With consultation of the land manager, more intensive grazing management may result in improved harvest efficiencies and increased carrying capacity.

Plant Community	Average Annual Production (lbs./acre, air-dry)	Stocking Rate* (AUM/acre)
Bluestem/Needlegrass (1.1 & 3.1)	3,700	1.01
Needlegrass/Little Bluestem/Kentucky Bluegrass (3.2)	3,000	0.82
Smooth Bromegrass/Kentucky Bluegrass (4.1)	3,400	0.93
Kentucky Bluegrass/Quackgrass (4.2)	2,400	0.66
Annual/Pioneer, Non-Native Perennial (4.3)	1,000	0.27

*Based on 912 lbs./acre (air-dry weight) per Animal Unit Month (AUM), and on 25 percent harvest efficiency (refer to United States Department of Agriculture (USDA) Natural Resources Conservation Service (NRCS), National Range and Pasture Handbook).

Grazing by domestic livestock is one of the major income-producing industries in the area. Rangeland, in this area, may provide yearlong forage. During the dormant period, the forage for livestock will likely be lacking protein to meet livestock requirements and added protein will allow ruminants to better utilize the energy stored in grazed plant materials. A forage quality test (either directly or through fecal sampling) should be used to determine the level of supplementation needed.

Hydrology Functions

Water is the principal factor limiting forage production on this site. This site is dominated by soils in hydrologic group B, with some soils in hydrologic group C. Infiltration and runoff potential for this site varies from moderate to high depending on soil hydrologic group, slope and ground cover. In many cases, areas with greater than 75 percent ground cover have the greatest potential for high infiltration and lower runoff. An example of an exception would be where shortgrasses form a strong sod and dominate the site. Dominance by blue grama, bluegrass, and/or smooth bromegrass will result in reduced infiltration and increased runoff. Areas where ground cover is less than 50 percent have the greatest potential to have reduced infiltration and higher runoff (refer to Section 4, NRCS National Engineering Handbook for runoff quantities and hydrologic curves).

Recreational Uses

This site provides hunting, hiking, photography, bird watching and other opportunities. The wide varieties of plants that bloom from spring until fall have an esthetic value that appeals to visitors.

Wood Products

No appreciable wood products are typically present on this site.

Other Products

Seed harvest of native plant species can provide additional income on this site.

Supporting Information

Associated Sites

Thin Upland (R102BY012SD), Loamy Overflow (R102BY020SD)

Similar Sites

(R102BY020SD) – Loamy Overflow [more big bluestem; higher production]
(R102BY011SD) – Clayey [more green needlegrass; less big bluestem]

Inventory Data References

Information presented here has been derived from NRCS clipping data and other inventory data. Field observations from range-trained personnel were also used. Those involved in developing this site include: Stan Boltz, Range Management Specialist, NRCS.

State Correlation

This site occurs entirely within SD.

Field Offices/Counties

Brookings, SD	Brookings	Howard, SD	Miner	Salem, SD	McCook
Canton, SD	Lincoln	Madison, SD	Lake	Sioux Falls, SD	Minnehaha
Elk Point, SD	Union	Parker, SD	Turner	Vermillion, SD	Clay
Flandreau, SD	Moody	Parkston, SD	Hutchinson	Yankton, SD	Yankton

Relationship to Other Established Classifications

Level IV Ecoregions of the Conterminous United States: 46k – Prairie Coteau; 46n – James River Lowland.

Other References

High Plains Regional Climate Center, University of Nebraska, 830728 Chase Hall, Lincoln, NE 68583-0728. (<http://hpccsun.unl.edu>).

USDA, NRCS. National Water and Climate Center, 101 SW Main, Suite 1600, Portland, OR 97204-3224. (<http://www.wcc.nrcs.usda.gov>).

USDA, NRCS. National Range and Pasture Handbook, September 1997.

USDA, NRCS. National Soil Information System, Information Technology Center, 2150 Centre Avenue, Building A, Fort Collins, CO 80526. (<http://nasis.nrcs.usda.gov>).

USDA, NRCS. 2001. The PLANTS Database, Version 3.1 (<http://plants.usda.gov>). National Plant Data Center, Baton Rouge, LA 70874-4490 USA.

Site Description Approval

SD, State Range Management Specialist

Date