

NATURAL RESOURCES CONSERVATION SERVICE
CONSERVATION PRACTICE STANDARD

FORAGE HARVEST MANAGEMENT

(Ac.)

CODE 511

DEFINITION

The timely cutting and removal of forages from the field as hay, green-chop or ensilage.

PURPOSE

- Optimize yield and quality of forage at the desired levels
- Promote vigorous plant re-growth
- Manage for the desired species composition
- Use forage plant biomass as a soil nutrient uptake tool
- Control insects, diseases and weeds
- Maintain and/or improve wildlife habitat

CONDITIONS WHERE PRACTICE APPLIES

This practice applies to all land uses where machine harvested forage crops are grown.

CRITERIA

General Criteria Applicable to All Purposes

Forage will be harvested at a frequency and height that optimizes the desired forage stand, plant community, and stand life. Follow AgriLIFE EXTENSION recommendations for forage harvest based on stage of maturity, moisture content, length of cut, stubble height and harvest interval. The following criteria must be met:

Stage of Maturity. Harvest forage at the stage of maturity that provides the desired quality and quantity without compromising plant vigor and stand longevity. Delay harvest if prolonged or heavy precipitation is forecast that would seriously damage cut forage.

Moisture Content. Harvest silage/haylage crops within the optimum moisture range for the type of storage method(s) or structure(s) being utilized.

AgriLIFE EXTENSION recommendations must be followed for optimum moisture content and levels as well as methods and techniques to monitor and/or determine moisture content and levels. Refer to AgriLIFE EXTENSION publication "[Hay Production in Texas](#)".

Avoid fermentation and seepage losses of digestible dry matter from direct cut hay crop silage (moisture content >70%) by treatment with chemical preservatives or add dry feedstuffs.

For optimal dry hay quality, rake hay at 30 to 40 percent moisture and ted or invert swaths when moisture is above 40 percent.

To preserve forage quality and quantity, moisture content of hay at baling for large round bales should

not exceed 18 percent; for small square bales, moisture content should not exceed 20 percent. Bale forced air dried hay at 20 – 35 percent moisture.

Length of Cut. When harvested for ensilage, forage will be chopped to a size appropriate for type of storage structure used and optimal effective fiber. The length of chop selected will allow adequate packing to produce the anaerobic conditions necessary to ensure the proper ensiling process.

A shorter chop length on very dry silage may help to ensure good packing and adequate silage density.

Stubble Height. Cut forage plants at a height that will promote the vigor and health of the desired species. Cutting heights will provide adequate residual leaf area; adequate numbers of terminal, basal or auxiliary tillers or buds; insulation from extreme heat or cold; and/or unsevered stem bases that store food reserves needed for full, vigorous recovery. For introduced sod forming grasses a minimum cutting height of 3 inches and a minimum cutting height of 4 to 8 inches for introduced and native bunch type grasses will help insure longevity of the stand, see Prescribed Grazing Standard (528) for specific guidance on minimum use heights. The final warm season hay harvest should be made 4 to 6 weeks before the first killing frost.

Contaminants. Forage shall not contain contaminants that can cause illness or death to the animal being fed or rejection of the offered forage.

Additional Criteria to Improve or Maintain Stand Life, Plant Vigor and Forage Species Mix

Stage of Maturity and Harvest Interval. Cut forage plants at a stage of maturity or harvest interval range that will provide adequate food reserves and/or basal or auxiliary tillers or buds for regrowth and/or reproduction to occur without loss of plant vigor.

Cut reseeding annuals at a stage of maturity and frequency that ensures the production of ample viable seed or carryover of hard seed to maintain desired stand density.

If plants show signs of short-term environmental stress, harvests will be adjusted in a manner that encourages the continued health and vigor of the stand.

Manipulate timing and cutting heights of harvest to ensure germination and establishment of reseeding or seeded annuals.

Additional Criteria for Use as a Nutrient Uptake Tool

Employ a harvest regime that utilizes the maximum amount of available or targeted nutrients. Using this practice for this purpose may require more frequent harvests to increase uptake instead of managing for stand longevity. Expected nutrient uptake values are located in the Texas NRCS 590-633 Spreadsheet/Planning tool ([S_Crops](#)).

Additional Criteria to Control Disease, Insect, Weed and Invasive Plant Infestations

Follow AgriLIFE EXTENSION guidelines when available for control of disease, insect and weed infestations to forage. For insect control guidelines refer to "[Managing Insect and Mite Pests of Texas Forage Crops](#)". For weed control guidelines refer to "[Suggestions for Weed Control in Pastures and Forages](#)".

Schedule harvest periods to control disease, insect, and weed infestations. When a pesticide is used to control disease, insects or weeds, adhere to the specified days to harvest period stated on the pesticide label. Evaluate pest management options by planning conservation practice standard 595 Integrated Pest Management for all forage areas to be harvested.

Plan and schedule removal of invasive plants and noxious weeds as needed.

Lessen incidence of disease, insect damage, and weed infestation by managing harvests to maintain a full, vigorous, dense forage stand.

Additional Criteria to Improve Wildlife Habitat Values

If client objectives include providing suitable habitat for desired wildlife specie(s) then appropriate harvest schedule(s), cover patterns, and minimum plant heights to provide suitable habitat for the desired specie(s) should be implemented and maintained.

Time harvests to benefit the desired wildlife species by following state guidelines. Avoid forage harvest during primary nesting season for ground-nesting wildlife ([March 1st through July 1st](#)).

Coordinate this practice with conservation practice standard Upland Wildlife Habitat Management (645) and accompanying job sheets.

CONSIDERATIONS

Where applicable, coordinate this practice with NRCS practice standard Prescribed Grazing (528).

When nutrients or other soil amendments are applied coordinate forage harvests with NRCS practice standard Nutrient Management (590) and/or Waste Utilization (633) as appropriate. An excess or improper balance of nutrients such as nitrogen can produce plant material that causes toxicity in some animals.

Produce stored forages of the quality needed for optimum performance of the animal being fed. Legume forages too low in fiber can lead to metabolic disorders in ruminants and an economic loss to the producer due to lowered animal performance. Consider analyzing harvested forages for feed quality. Coordinate this practice with NRCS practice standard Feed Management (592).

In animal feeding operations, harvesting and feeding high quality forage such as silage and green chop recycles nutrients on the farm and reduces import of nutrients from purchased forage, which in turn reduces the potential of excessive nutrient build-up in the soil. Forage testing and accurate yield monitoring are encouraged when planning Forage and Biomass Harvest for this purpose.

Hay should be marketed based on forage analysis and weight to better assure production costs are recovered. Hay should be evaluated based on Relative Feed Value (RFV), protein content and physical characteristics. Relative feed value is calculated as follows: (% Digestible Dry Matter) x (% Dry Matter Intake) x (0.775). DDM and DMI are derived from Acid Detergent Fiber and Neutral Detergent Fiber.

Stockpiling forage in the fall for use in early winter should be considered to reduce the costs of hay production and feeding. AgriLIFE EXTENSION and others have shown that stockpiled bermudagrass and bahiagrass maintain quality as standing forage into mid-January in most years.

To control forage plant diseases, insects, and movement of weeds, clean harvesting equipment after harvest and before storing. Cut forages after dew, rain, or irrigation water on leaves has evaporated.

Consider delaying harvest if prolonged or heavy precipitation is forecast that would reduce forage quality.

Direct cut grass and legume silage can create silage leachate (seepage) in storage. Consider use of practice standards Runoff Management System (570) and Waste Storage Facility (313).

In conjunction with harvest options, consider storage and feeding options that will retain acceptable forage quality and minimize digestible dry matter loss. Consider storage location for large square or round bales with regards to inside vs. outside, along hedgerows, winter/mud considerations etc. Hay feeding areas should be moved periodically to avoid damage and over use of the pasture.

Where weather conditions make it difficult to harvest the desired quality of forage consider use of mechanical or chemical conditioners, forced air barn curing and/or ensile.

In regions where rainfall and/or humidity levels cause unacceptable forage quality losses consider green chopping or ensiling the forage to reduce or eliminate field drying time. Other options are: the use of desiccants, preservatives, or macerating implements to reduce field-drying time. These techniques can improve the timeliness of harvest and preserve forage quality.

To reduce safety hazards, avoid operating harvesting and hauling equipment on field slopes over 25 percent, particularly on cross slope traffic patterns.

Consider Harvesting Forages in the afternoon to optimize water soluble carbohydrates and nutritional quality.

To minimize haying operation impact on wildlife, mowing or swathing operation should begin in the middle of the field and work outward.

PLANS AND SPECIFICATIONS

Place the detailed specifications in a site-specific job or design sheet or in the practice narrative in the conservation plan.

Plans and Specifications must include as minimum for the forage harvest operations:

1. Production / yield goals, objectives, specific purpose (such as high forage quantity and quality or nutrient uptake, etc.)
2. Forage species to be harvested
3. For each dominant forage species harvested show:
 - a. Method of harvest
 - b. Stage of maturity
 - c. Optimal harvest moisture content
 - d. Length of cut
 - e. Stubble height to be left
 - f. Harvest interval including late harvest if applicable
 - g. Contaminant avoidance recommendations.

These plans and specifications shall be consistent with this standard and shall describe the requirement for applying the practice to achieve its intended purpose.

OPERATION AND MAINTENANCE

Before forage harvest, clear fields of debris that could damage machinery or if ingested by livestock, lead to sickness (for example, hardware disease) or death.

Operate all forage harvesting equipment at the optimum settings and speeds to minimize loss of leaves.

To control forage plant diseases, insects, and movement of weeds, clean harvesting equipment after harvest and before storing.

Set shear-plate on forage chopper to the proper theoretical cut for the crop being harvested. Keep knives well sharpened. Do not use re-cutters or screens unless forage moisture levels fall below recommended levels for optimum chopping action.

Follow all agricultural equipment manufacturer's safety measures when operating forage harvesting equipment.

Regardless of silage/haylage storage method, ensure good compaction and an airtight seal to exclude oxygen and mold or bacterial formations.

Dispose of the plastic wrap or bags used to store forage in an environmentally sound manner.

REFERENCES:

- Ball, D. M., C. S. Hoveland, & G. D. Lacefield. 2002. Southern Forages. (Third Ed.). Potash & Phosphate Institute.
- Barnes, R. F., C. J. Nelson., K.J. Moore & M. Collins. 2007. Forages, The Science of Grassland Agriculture, Sixth Edition.
- Hanson, A. A., D. K. Barnes, & R. R. Hill, Jr. 1988. Alfalfa and Alfalfa Improvement.
- Ishler, V. A., etal. 1991. Harvesting and Utilizing Silage. Pennsylvania State University Circular 396.
- Matches, A. G. 1973. Anti-Quality Components of Forages. Crop Science Society of America Special Pub. No. 4.
- Pitt, R. E. 1990. Silage and Hay Preservation. Northeast Regional Agricultural Engineering Service.
- Leep R., J. Andresen, D.H. Min & A. Pollyea. 2006. Late Summer and Fall Harvest Management of Alfalfa. First Alfalfa Technology Conference. Michigan State University, East Lansing, MI.
- Henning, J.C., M. Collins, D. Ditsch, G.D. Lacefield. 1998. Baling Forage Crops for Silage. Univ. of Ky., Ext. Pub. AGR-173, Lexington, KY.
- Serotkin, N., Ed. The Penn State Agronomy Guide, 1995-1996. Pennsylvania State University. 1994. University Park, PA.
- Smith, D. 1975. Forage Management in the North, Third Edition. Kendall/Hunt Publishing Company.
- Taylor, N. L. 1985. Clover Science and Technology. American Society of Agronomy..

