

NATURAL RESOURCES CONSERVATION SERVICE
CONSERVATION PRACTICE GENERAL SPECIFICATION

HERBACEOUS WEED CONTROL

(Ac.) CODE 315

General Use

Herbaceous weed control will be applied in a manner to achieve the desired control of the target species and protection of desired species. This will be accomplished by mechanical, chemical, burning or biological methods either alone or in combination. When burning is used as a method, the Prescribed Burning standard (338) will also be applied.

NRCS will not develop biological or chemical treatment recommendations except for biological control utilizing grazing animals. Prescribed Grazing (528) is used to ensure desired results are achieved and maintained.

NRCS may provide clients with acceptable biological and/or chemical control references. Chemical references must be from Chemical Weed and Brush Control Suggestions for Rangeland RM-1466 or <http://pestman.tamu.edu/>

Livestock and people access will be controlled based on management methods applied and restrictions as listed on the chemical labels.

Consider using Integrated Pest Management (595) in support of herbaceous weed control. Consider soil erosion potential and difficulty of vegetation establishment when choosing a method of control that causes soil disturbance.

Consider the appropriate time period for treatment. Some herbaceous weed control activities can be effective when applied within a single year; others may require multiple years of treatment(s) to achieve desired objectives.

Consider impacts to wildlife species, in general, treatments that create a mosaic pattern may be the most desirable.

Consider impacts to wildlife food supplies, space, and cover availability when planning the method and amount of herbaceous weed control.

State issued licenses may be required when using chemical pesticide treatments.

For air quality purposes, consider using chemical methods of herbaceous weed control that minimize chemical drift and excessive chemical usage and consider mechanical methods of herbaceous weed control that minimize the entrainment of particulate matter.

Adjacent land uses must be considered before chemicals are used.

Herbaceous Weed Control Plan:

Prepare plans and specifications for each field or treatment unit according to the criteria included in this specification.

At a minimum, a herbaceous weed control plan shall include:

1. Goals and objectives statement.
2. Plan map and soil map for the site.
3. Pre-treatment cover or density of the target plant(s) and the planned post-treatment cover or density and desired efficacy.

4. Maps, drawings, and/or narratives detailing or identifying areas to be treated, pattern of treatment (if applicable), and areas that will not be disturbed.
5. A monitoring plan that identifies what shall be measured (including timing and frequency) and the changes in the plant community (compare with objectives) that will be achieved.

For Mechanical Treatment Methods. Plans and specifications will include items 1 through 5 above, plus the following:

- Type of equipment to use for management
- Dates of treatment for effective management.
- Operating instructions (if applicable)
- Techniques and procedures to be followed.

For Chemical Treatment Methods. Plans and specifications will include items 1 through 5, above, plus the following:

- Acceptable chemical treatment references for containment and management of target species
- Document techniques to be used, planned dates and rates of application
- Evaluation and interpretation of herbicide risks associated with the selected treatment(s) using WIN-PST or other approved tools.
- Any special mitigation, timing considerations or other factors (such as soil texture and organic matter content) that must be considered to ensure the safest, most effective application of the herbicide
- Reference to product label instructions

For Biological Treatment Methods. Plans and specifications will include items 1 through 5, above, plus the following:

- Acceptable biological treatment references for the selected biological agent used to contain and manage the target species
- Document release date, kind, and number of livestock
- Timing, frequency, duration and intensity of grazing or browsing
- Desired degree of grazing or browsing use for effective management of target species
- Maximum allowable degree of use on desirable non-target species
- Special mitigation, precautions, or requirements associated with the selected treatment(s)

VARIANCES

- Any requests for variances are to be submitted to the State Rangeland Management Specialist.

The Following methods may be utilized after Range Planting (550) and Forage and Biomass Planting (512), during establishment:

Herbicides

- Chemicals used must be federally and locally registered and must be applied in accordance with authorized registered uses, directions on label, and other federal or state policies and requirements.

- Seeded species should have 3 to 5 leaves per plant before herbicides are applied. Generally, when 3 weeds per square foot or a 50% canopy are observed, weed control should be considered.

Mowing

- Weeds should be mowed when they reach a height of 6 to 8 inches.
- Mowing will be above the height of seeded plants. The cover crop should also be maintained. Mowing will not be done when daily maximum air temperature exceeds 95° and the humidity falls below 30% to prevent dehydration of the seedlings. Generally, mowing should not be done after July 15.

Grazing

- Flash grazing by livestock may be used to control annual grasses and forbs and a time when they are small and palatable. This method will not be used later than July 15, except when abnormal summer moisture promotes excessive weed production. Flash grazing will not be used when the soils are wet and hoof action will damage seedlings.
- Flash grazing is using high concentrations of livestock to harvest palatable competitive plants in a short period of time. Should there be significant use or damage to seeded plants, the grazing should cease immediately. In cases where additional applications are needed, the procedure should be repeated soon enough to prevent the weedy vegetation from becoming tough or unpalatable.

Enhance Accessibility, Quantity, and Quality of Forage and/or Browse

- Herbaceous weed control will be applied in a manner to minimize negative impact to forage and/or other non-targeted plants. Timing and sequence of control shall be planned in coordination with specifications developed for Prescribed Grazing (528) or Forage Harvest Management (511).

Restore or Release Native or Create Desired Plant Communities and Wildlife Habitats Consistent with the Ecological Site

- Apply herbaceous weed control in a manner to protect the health and vigor of native or desired plant species.
- Use applicable Ecological Site Description (ESD) State and Transition models, to develop specifications that are ecologically sound and defensible. Treatments must be congruent with dynamics of the ecological site(s) and keyed to states and plant community phases that have the potential and capability to support the desired plant community. If an ESD is not available, base specifications on the best approximation of the desired plant community composition, structure, and function.
- Treatments will be conducted during periods of the year when weed species are most vulnerable and will promote restoration of the native or desired plant communities.
- Apply herbaceous weed control in a manner that maintain or enhance important wildlife habitat requirements.
- Treatments will be conducted during periods of the year that accommodate reproduction and other life cycle requirements of target wildlife and pollinator species.
- Apply treatments that maintain or enhance plant community composition and structure to meet the requirements of target wildlife species.

Protect Soils and Control Erosion

- Apply herbaceous weed control to minimize soil disturbance and soil erosion.
- Additional treatment will be applied to protect soils and prevent erosion.

Reduce Fine-Fuels Fire Hazard and Improve Air Quality

- Treat weed species in a manner that creates a native or desired plant community which reduces the potential for accumulating excessive fuel loads and increased wildfire hazards.
- Apply treatment methods in a manner that minimize the potential for unintended impacts to air resources, e.g., smoke, chemical drift etc.

OPERATION AND MAINTENANCE

Operation. Herbaceous weed control practices shall be applied using approved materials and procedures. Operations will comply with all local, state, and federal laws and ordinances.

Success of the practice shall be determined by evaluating regrowth or reoccurrence of target species after sufficient time has passed to monitor the situation and gather reliable data. Length of evaluation periods will depend on the herbaceous weeds species being monitored, proximity of propagules (seeds, plant materials and roots) to the site, transport mode of seeds (wind or animals) and methods and materials used.

The operator will develop a safety plan for individuals exposed to chemicals, including telephone numbers and addresses of emergency treatment centers and the telephone number for the nearest poison control center. The National Pesticide Information Center (NPIC) telephone number in Corvallis, Oregon, may also be given for non-emergency information: **1-800-858-7384**

Monday to Friday 6:30 a.m. to 4:30 p.m. Pacific Time

The national Chemical Transportation Emergency Center (CHEMTRAC) telephone number is: 1-800-424-9300

- Follow label requirements for mixing/loading setbacks from wells, intermittent streams and rivers, natural or impounded ponds and lakes, and reservoirs.
- Post signs, according to label directions and/or federal, state, tribal, and local laws, around fields that have been treated. Follow restricted entry intervals.
- Dispose of herbicide and herbicide containers in accordance with label directions and adhere to federal, state, tribal, and local regulations.
- Read and follow label directions and maintain appropriate Material Safety Data Sheets (MSDS). MSDS and herbicide labels may be accessed on the Internet at: <http://www.greenbook.net/>
- Calibrate application equipment according to recommendations before each seasonal use and with each major chemical and site change.
- Replace worn nozzle tips, cracked hoses, and faulty gauges on spray equipment.
- Maintain records of plant management for at least two years. Herbicide application records shall be in accordance with USDA Agricultural Marketing Service's Pesticide Recordkeeping Program and state-specific requirements.

Maintenance. Following initial application, some regrowth, re-sprouting, or reoccurrence of herbaceous weeds may be expected. Spot treatment of individual plants or areas needing re-treatment should be completed as needed when weed vegetation is most vulnerable to desired treatment procedures.

Review and update the plan periodically in order to incorporate new IPM technology; response to grazing management and complex weed population changes; and avoid the development of weed resistance to herbicide chemicals.

REFERENCES

- AgriLIFE Extension. 2014. Chemical Weed and Brush Control Suggestions for Rangeland. Publ. RM-1466, Texas A&M University, College Station, TX
- Alex, J.F., and C.M. Switer. 1982. Ontario weeds. Publ. 505, University of Guelph – Ontario Agricultural College, Guelph, Ontario, Canada.
- American Sheep Industry, A. Peischel and D.D. Henry, Jr., 2006. Targeted Grazing: a Natural Approach to Vegetation Management and Landscape Enhancement.
- Cheney, M. Common poisonous plants of western Washington which affect livestock. [Online] Available at http://www.piercecountycd.org/tip_toxicplants_p.html. (Accessed 15 October 2008).
- Ciba-Geigy Corp. Plants that poison livestock: Information chart.
- Cornell University Department of Animal Science. Plants Poisonous to Livestock and Other Animals. [Online]. Available at: <http://www.ansci.cornell.edu/plants/>
- DeWolf , G. and M. Hondalus. 1988. Common Massachusetts plants poisonous to horses. University of Massachusetts Cooperative Extension Service, Amherst, MA.
- Ensminger, M.E. 1992. The stockman's handbook. (7th Ed.) The Interstate Printers and Publishers, Inc. Danville, Il.
- Evers, R.A., and R.P. Link. 1972. Poison plants of the Midwest and their effects on livestock. Special Publication 24, University of Illinois – College of Agriculture, Urbana, Il.
- Hamilton, G.W., and J.R. Mitchell. 2001. [Online] Poisonous plants in pastures. Univ. of New Hampshire Coop. Ext. Serv., Durham, NH. Available at http://extension.unh.edu/resources/representation/Resource000623_Rep645.pdf. (Accessed 15 October 2008).
- Hill, R.J., and D. Folland. 1986. Poisonous plants of Pennsylvania. Pennsylvania Department of Agriculture, Harrisburg, PA.
- Reed, C.F. 1970. Selected weeds of the United States. Agriculture Handbook No. 366, U.S. Government Printing Office, Washington, D.C.
- USDA-ARS. 2006. Bulletin 415 - Plants poisonous to livestock in the Western states. [Online]. Available at <http://www.ars.usda.gov/Services/docs.htm?docid=12140> (Updated 08 February 2006, accessed 15 October 2008).

