



**NATURAL RESOURCES CONSERVATION SERVICE
CONSERVATION PRACTICE STANDARD**

FORAGE HARVEST MANAGEMENT

**CODE 511
(ACRE)**

DEFINITION

The timely cutting and removal of forages from the field as hay, green-chop or ensilage.

PURPOSE

- Optimize yield and quality of forage at the desired levels
- Promote vigorous plant re-growth
- Manage for the desired species composition
- Use forage plant biomass as a soil nutrient uptake tool
- Control insects, diseases and weeds
- Maintain and/or improve wildlife habitat

CONDITIONS WHERE PRACTICE APPLIES

This practice applies to all land uses where machine harvested forage crops are grown.

CRITERIA

General Criteria Applicable to All Purposes

Forage will be harvested at a frequency and height that optimizes the desired forage stand, plant community, and stand life. Follow UW Extension recommendations for forage harvest based on stage of maturity, moisture content, length of cut, stubble height and harvest interval. The following criteria must be met:

Stage of Maturity. Harvest forage at the stage of maturity that provides the desired quality and quantity without compromising plant vigor and stand longevity.

Moisture Content. Harvest silage/haylage crops within the optimum moisture range for the type of storage method(s) or structure(s) being utilized.

UW Extension recommendations must be followed for optimum moisture content and levels as well as methods and techniques to monitor and/or determine moisture content and levels.

Avoid fermentation and seepage losses of digestible dry matter from direct cut hay crop silage (moisture content >70 percent) by treatment with chemical preservatives or add dry feedstuffs.

For optimal dry hay quality, rake hay at 30 to 40 percent moisture and ted or invert swaths when moisture is above 40 percent.

To preserve forage quality and quantity, bale field cured hay at 15–20 percent moisture and bale force air-dried hay and 20–35 percent moisture.

Length of Cut. When harvested for ensilage forage will be chopped to a size appropriate for type of storage structure used and optimal effective fiber. The length of chop selected will allow adequate packing to produce the anaerobic conditions necessary to ensure the proper ensiling process.

A shorter chop length on very dry silage may help to ensure good packing and adequate silage density.

Stubble Height. Cut forage plants at a height that will promote the vigor and health of the desired species. Cutting heights will provide adequate residual leaf area; adequate numbers of terminal, basal or auxiliary tillers or buds; insulation from extreme heat or cold; and/or unsevered stem bases that store food reserves needed for full, vigorous recovery. Follow UW Extension recommendations for proper stubble heights and harvest dates to avoid winterkill of forage species in cold climates.

Contaminants. Forage shall not contain contaminants that can cause illness or death to the animal being fed or rejection of the offered forage. Check UW Extension contaminant notices, cautions, and recommendations for the specific harvest information.

Additional Criteria to Improve or Maintain Stand Life, Plant Vigor and Forage Species Mix

Stage of Maturity and Harvest Interval. Cut forage plants at a stage of maturity or harvest interval range that will provide adequate food reserves and/or basal or auxiliary tillers or buds for regrowth and/or reproduction to occur without loss of plant vigor.

Cut reseeding annuals at a stage of maturity and frequency that ensures the production of ample viable seed or carryover of hard seed to maintain desired stand density.

If plants show signs of short-term environmental stress, harvests will be adjusted in a manner that encourages the continued health and vigor of the stand. Follow UW Extension recommendations in these cases.

Manipulate timing and cutting heights of harvest to ensure germination and establishment of reseeding or seeded annuals.

Additional Criteria for Use as a Nutrient Uptake Tool

Employ a harvest regime that utilizes the maximum amount of available or targeted nutrients. Using this practice for this purpose may require more frequent harvests to increase uptake.

Additional Criteria to Control Disease, Insect, Weed and Invasive Plant Infestations

Follow UW Extension guidelines when available for control of disease, insect, weed and invasive plant infestations to forage crops.

Schedule harvest periods to control disease, insect, and weed infestations. When a pesticide is used to control disease, insects or weeds, adhere to the specified days to harvest period stated on the pesticide label. Evaluate pest management options by planning Wisconsin Conservation Practice Standard (WI NRCS CPS), Pest Management (Code 595) for all forage areas to be harvested. Also plan and schedule removal of invasive plants and noxious weeds.

Lessen incidence of disease, insect damage, and weed infestation by managing harvests to maintain a full, vigorous, dense forage stand.

Cut forages after dew, rain, or irrigation water on the leaves has evaporated.

Additional Criteria to Improve Wildlife Habitat Values

If client objectives include providing suitable habitat for desired wildlife species then appropriate harvest schedule(s), cover patterns, and minimum plant heights to provide suitable habitat for the desired species should be implemented and maintained.

Time harvests to benefit the desired wildlife species by following state guidelines.

Coordinate this practice with WI NRCS CPS, Upland Wildlife Habitat Management (Code 645) and accompanying job sheets.

CONSIDERATIONS

Where applicable coordinate this practice with WI NRCS CPS, Prescribed Grazing (Code 528).

When nutrients or other soil amendments are applied coordinate forage harvests with WI NRCS CPS, Nutrient Management (Code 590) and/or Waste Utilization (Code 633) as appropriate. An excess or improper balance of nutrients such as nitrogen can produce plant material that causes toxicity in some animals.

Produce stored forages of the quality needed for optimum performance of the animal being fed. Legume forages too low in fiber and lead to metabolic disorders in ruminants and an economic loss to the producer due to lowered animal performance. Consider analyzing harvested forages for feed quality. Coordinate this practice with WI NRCS CPS, Feed Management (Code 592).

Direct cut grass and legume silage can create silage leachate (seepage) in storage. Consider use of WI NRCS CPS, Runoff Management System (Code 570) and Waste Storage Facility (Code 313).

In conjunction with harvest options, consider storage and feeding options that will retain acceptable forage quality and minimize digestible dry matter loss.

Where weather conditions make it difficult to harvest the desired quality of forage consider use of mechanical or chemical conditioners, forced air barn curing and/or ensile.

Consider delaying harvest if prolonged or heavy precipitation is forecast that would reduce forage quality.

When rainfall and/or humidity levels cause unacceptable forage quality losses consider green chopping or ensiling the forage to reduce or eliminate field drying time. Other options are: the use of desiccants, preservatives, or macerating implements to reduce field-drying time.

To reduce safety hazards, avoid operating harvesting and hauling equipment on field slopes over 25 percent, particularly on cross slope traffic patterns.

Consider Harvesting Forages in the afternoon to optimize water soluble carbohydrates and nutritional quality.

Select cultivars that are suitable for the harvest regime, species mix, and forage quality desired. For specific nutrient uptake, select species that can maximize uptake. See WI NRCS CPS, Pasture and Hay Planting (Code 512).

When insect and disease outbreaks exceed economic thresholds and are uncontrollable by harvest management pesticide applications may be needed. Another option is to select a resistant cultivar when the stand is replaced. See WI NRCS CPS, Pest Management (Code 595).

To control forage plant diseases, insects, and weeds, clean harvesting equipment after harvest and before storing. Do not cut forages until dew, rain, or irrigation water on leaves has evaporated.

When weed infestation exceeds the economic threshold and is uncontrollable by forage harvest management alone, weed management should be planned and applied. See WI NRCS CPS, Pest Management (Code 595).

Take care not to produce stored forages whose quality is not that needed for optimum performance of the animal being fed. For instance, immature legume forages can be too low in fiber and lead to metabolic disorders in ruminants and an economic loss to the producer due to lowered animal performance.

PLANS AND SPECIFICATIONS

Place the detailed specifications in a site-specific job or design sheet or in the practice narrative in the conservation plan.

Plans and Specifications must include as minimum for the forage harvest operations:

- Goals, objectives, specific purpose (such as high forage quantity and quality or nutrient uptake, etc.)
- Forage species to be harvested

By each dominant forage species harvested show:

- Method of harvest
- Stage of maturity
- Optimal harvest moisture content
- Length of cut
- Stubble height to be left
- Harvest interval including late harvest if applicable
- Contaminant avoidance recommendations.

Document any additional factors to consider prior to harvest:

- Wildlife nesting dates
- Late season harvest dates
- Strategies to maximize nutrient uptake
- Strategies to address insect or weed problems

These plans and specifications shall be available through appropriate job sheets and other materials for applying the practice to achieve its intended purpose.

OPERATION AND MAINTENANCE

Before forage harvest, clear fields of debris that could damage machinery or if ingested by livestock, lead to sickness (for example, hardware disease) or death.

Operate all forage harvesting equipment at the optimum settings and speeds to minimize loss of leaves.

To control forage plant diseases, insects, and movement of weeds, clean harvesting equipment after harvest and before storing.

Set shear-plate on forage chopper to the proper theoretical cut for the crop being harvested. Keep knives well sharpened. Do not use re-cutters or screens unless forage moisture levels fall below recommended levels for optimum chopping action.

Follow all agricultural equipment manufacturer's safety measures when operating forage harvesting equipment.

Regardless of silage/haylage storage method, ensure good compaction and an airtight seal to exclude oxygen and mold or bacterial formations.

Dispose of the plastic wrap or bags used to store forage in an environmentally sound manner.

Regardless of silage/haylage storage method, ensure good compaction and an air-tight seal to exclude oxygen and mold formation.

FEDERAL, TRIBAL, STATE AND LOCAL LAWS

Users of this standard should be aware of potentially applicable federal, tribal, state and local laws, rules, regulations or permit requirements governing cover crops. This standard does not contain the text of federal, tribal, state or local laws.

REFERENCES:

Ball, D. M., C. S. Hoveland, & G. D. Lacefield. Southern Forages. 1991. Potash & Phosphate Institute, Norcross, GA.

Barnes, R. F., D. A. Miller, & C. J. Nelson. Forages, The Science of Grassland Agriculture, Fifth Edition. 1995. Iowa State University Press, Ames, IA.

Hanson, A. A., D. K. Barnes, & R. R. Hill, Jr. Alfalfa and Alfalfa Improvement. 1988. American Society of Agronomy, Madison, WI.

Ishler, V. A. Et al. Harvesting and Utilizing Silage. 1991. Penn State University Circular 396. University Park, PA.

Matches, A. G. Anti-Quality Components of Forages. 1973. Crop Science Society of America Special Pub. No. 4, Madison, WI

Pitt, R. E. Silage and Hay Preservation. 1990. Northeast Regional Agricultural Engineering Service. Ithaca, NY.

Serotkin, N., Ed. The Penn State Agronomy Guide, 1995-1996. Pennsylvania State University. 1994. University Park, PA.

Smith, D. Forage Management in the North, Third Edition. 1975. Kendall/Hunt Publishing Company, Dubuque, IA.

Taylor, N. L. Clover Science and Technology. 1985. American Society of Agronomy, Madison, WI
Alfalfa Management Guide NCR547.

UWEX Publication A3529, Pastures for Profit.

UWEX Publication A3492, Red Clover Establishment, Management, and Utilization.

USDA, NRCS Wisconsin Field Office Technical Guide (FOTG), Section IV, Practice Standards and Specifications.

In accordance with Federal civil rights law and U.S. Department of Agriculture (USDA) civil rights regulations and policies, the USDA, its Agencies, offices, and employees, and institutions participating in or administering USDA programs are prohibited from discriminating based on race, color, national origin, religion, sex, gender identity (including gender expression), sexual orientation, disability, age, marital status, family/parental status, income derived from a public assistance program, political beliefs, or reprisal or retaliation for prior civil rights activity, in any program or activity conducted or funded by USDA (not all bases apply to all programs). Remedies and complaint filing deadlines vary by program or incident. Persons with disabilities who require alternative means of communication for program information (e.g., Braille, large print, audiotape, American Sign Language, etc.) should contact the responsible Agency or USDA's TARGET Center at (202) 720-2600 (voice and TTY) or contact USDA through the Federal Relay Service at (800) 877-8339. Additionally, program information may be made available in languages other than English. To file a program discrimination complaint, complete the USDA Program Discrimination Complaint Form, AD-3027, found online at [How to File a Program Discrimination Complaint](#) and at any USDA office or write a letter addressed to USDA and provide in the letter all of the information requested in the form. To request a copy of the complaint form, call (866) 632-9992. Submit your completed form or letter to USDA by: (1) mail: U.S. Department of Agriculture, Office of the Assistant Secretary for Civil Rights, 1400 Independence Avenue, SW, Washington, D.C. 20250-9410; (2) fax: (202) 690-7442; or (3) email: program.intake@usda.gov. USDA is an equal opportunity provider, employer, and lender.